

May 1988

A.E.Res. 88-7

# **THE ECONOMICS OF HATCHERY PRODUCED ALGAE AND BIVALVE SEED**

**by**

**Julia A. Myers**

**and**

**Richard N. Boisvert**

Department of Agricultural Economics  
Cornell University Agricultural Experiment Station  
New York State College of Agriculture and Life Sciences  
A Statutory College of the State University  
Cornell University, Ithaca, New York 14853-7801

**It is the policy of Cornell University actively to support equality of educational and employment opportunity. No person shall be denied admission to any educational program or activity or be denied employment on the basis of any legally prohibited discrimination involving, but not limited to, such factors as race, color, creed, religion, national or ethnic origin, sex, age or handicap. The University is committed to the maintenance of affirmative action programs which will assure the continuation of such equality of opportunity.**

## THE ECONOMICS OF HATCHERY PRODUCED ALGAE AND BIVALVE SEED

by

Julia A. Myers and Richard N. Boisvert

### *ABSTRACT*

The bivalve aquacultural industry in the northeastern United States has been strengthened by declining and fluctuating natural harvests in the surrounding coastal region. Hatcheries in which the bivalves are grown to a small seed size have gained acceptance as a link in the aquacultural process. These firms are potentially able to provide a consistent, reliable source of seed to the bivalve aquaculturist.

This research contributes to an understanding of the hatchery industry by examining factors that influence production costs. The research is based on observations of a working hatchery located in the northeast and on current literature. The data were used to develop the framework for a computer program that can estimate variable production costs. This program required the development of an algorithm that simulates the production process. The model sums costs at different stages of animal growth, which are further broken down into smaller units of time.

There was sufficient information available from the literature and hatchery observations to simulate the overall production system, but expenditure records did not separate costs between stages or between hatchery and open-water field operations. The empirical analysis focuses on estimating costs of algae production, because data on this process were more readily isolated from the rest of the hatchery and because feeding costs are recognized as a large proportion of total costs.

The FORTRAN computer program based on the model focuses on the cost of producing algae feed, but as more details become available about other hatchery production processes it would be a simple matter to incorporate them into the existing FORTRAN code. The computer program combines estimates of the cost of algae production with survival and growth rates and feeding efficiencies to determine the costs of feeding a batch of bivalves to various sizes.

The model is applied to raising Atlantic Oyster seed. Twelve initial simulations were made, assuming different values for important parameters. To explore the implications of these feed cost simulations for the marketing of bivalve seed, some rough estimates of total costs of production per bivalve were developed on the basis of others' estimates of the proportion that feed costs are of total costs. Under the most optimistic assumptions, bivalve seed can be produced and marketed profitably. These conclusions must be qualified by the fact that sufficient data were not available from the hatchery under study to estimate all components of total cost. Collection of these types of data would help in further testing of the algorithm as well as contributing to an overall understanding of hatchery costs at each stage of production. Equally important from a research perspective is the need for better data on hatchery survival rates and the size-age and feeding rates.

## Table of Contents

	<u>Page</u>
<i>I - INTRODUCTION.....</i>	1
<i>II - FORMATION OF THE HATCHERY INDUSTRY .....</i>	2
Background.....	2
Population Dynamics .....	2
Demand Considerations .....	3
Alternatives .....	5
<i>III - HATCHERY PRODUCTION OF BIVALVE SEED.....</i>	7
The Hatchery.....	7
The Hatchery's Production System.....	7
The Hatchery's Physical Systems .....	13
Costs and Input Considerations.....	15
<i>IV - SIMULATING BIVALVE PRODUCTION COSTS.....</i>	17
Structure of the Model.....	17
Algorithms .....	19
Algae Price and Quantity.....	22
<i>V - EMPIRICAL TESTS OF THE SIMULATION MODEL .....</i>	25
The Data.....	25
Empirical Results.....	27
<i>VI - SUMMARY AND CONCLUSIONS .....</i>	35
References.....	37
<i>APPENDIX A.....</i>	39
<i>APPENDIX B.....</i>	83

## List of Tables

1. Daily Consumption of Algae Cells by Oysters and Clams.....	10
2. Optimum Algae Concentrations for Oyster Larvae.....	11
3. Inputs to Bivalve Seed Production in the Hatchery .....	16
4. Production Costs in Various Production Stages in Bivalve Hatcheries of Two Sizes.....	18
5. Estimated Equations for Use in Simulation.....	26
6. Description of Parameters for Alternative Simulation Runs .....	28
7. Summary of Simulation Output for Initial Runs.....	29
8. Algae Consumption for Selected Simulations .....	30
9. Summary of Simulations to Test Sensitivity of Survival Parameter .....	31
10. Sensitivity of Simulation Results to the Age-Size and Feed-Size Relationships.....	32
11. Alternative Estimates of Total Costs of Production (\$ Per Oyster).....	34

List of Figures

	<u>Page</u>
1. Map of Long Island Coastal Region.....	3
2. Commercial Landings from New York Marine District Waters.....	4
3. Long Island Clam Harvest, 1966 - 1982.....	4
4. Production Locations in the Hatchery.....	8
5. Flow Between Hatchery Production Stages.....	9
6. Energy and Water Flow Within the Hatchery.....	14
7. Bivalve Code Flowchart .....	19
8. Algae Price Subroutine .....	23
9. Algae Quantity Subroutine.....	24

# THE ECONOMICS OF HATCHERY PRODUCED ALGAE AND BIVALVE SEED

by

Julia A. Myers and Richard N. Boisvert\*

## I - INTRODUCTION\*

Over the past 25 years, bivalve aquaculture in the northeastern United States has become a reliable supplier of marketable oysters and clams. However, to initiate production, the industry requires adequate supplies of bivalve seed. Seed can be supplied from either natural sets or artificial spawning in a hatchery.

A hatchery operator's ability to produce and market seed as an efficient alternative to naturally-spawned seed requires accurate information about the market, the production process and production costs. As with most managers in a new, rapidly developing industry, hatchery operators have difficulty assembling adequate information to make effective investment and production decisions.

The research reported here contributes to an understanding of the economics of artificially spawned bivalve seed by: a) describing the hatchery industry's development and its production processes and b) formulating a model, associated algorithms and computer code by which operators may calculate variable production costs.

Much of the information about production processes, input requirements and

prices was obtained in 1983 from one hatchery on the northeast coast. That being its first full year of operation, the firm's record keeping system was new and not well enough developed to facilitate accurate measurement of the quantities of inputs used at every stage of a bivalve seed's development. However, feed contributes importantly to the total variable cost of bivalve seed (Im *et al.* 1976; Gates *et al.* 1974 and Bolton 1982). Data on algae production were quite good. Thus, the algorithms that calculate algae production and feeding cost are developed in detail. The algorithms can also be used to examine the sensitivity of results with respect to different assumptions about survival rates, growth rates and feeding efficiency.

The remainder of the bulletin begins, in Section *II*, by placing the hatchery industry within the context of the historical decline in the natural bivalve harvest. Various aspects of the hatchery industry affecting the firm and its costs, such as the industry's information network, the issue of property rights and market structure are also discussed. Section *III* describes the production processes using one hatchery as an illustration. This section also outlines choices that affect quality, quantity and timing of inputs and outputs and contains a general discussion of production costs. Section *IV* describes algorithms that simulate the cost of production for each of the system's components. Section *V* presents the empirical application of the program to the American Atlantic Oyster, including tests using data obtained from the hatchery and sensitivity analyses for the program's important variables. Preliminary estimates of total production costs per bivalve are compared with bivalve seed prices to determine at what age (size) the bivalve seed should be sold. Section *VI* summarizes the major findings and their implications for a hatchery's production.

\*J. A. Myers is a Research Associate at the University of California, San Diego, Laboratory for Mathematics and Statistics, and R. N. Boisvert is a professor in the Department of Agricultural Economics, Cornell University. The research draws extensively on Myers' thesis (1985) sponsored by the New York Sea Grant Institute under a grant from the Office of Sea Grant, NOAA. The cooperation from individuals at the hatchery where data were collected is gratefully acknowledged. The opinions expressed in the report are those of the authors and not necessarily those of the funding agency nor other cooperators in the research.

## II - FORMATION OF THE HATCHERY INDUSTRY

In the northeastern United States, the long-standing clam and oyster fishing industry has given rise to the development of the bivalve hatchery industry. Over harvests and increased urbanization, with the consequent decline in the once-abundant natural bivalve population, have forced harvesters to look for alternatives. Among these alternatives are government regulation and incentives, but perhaps the most important alternative is the aquacultural industry, which is replacing traditional methods of harvest. Recently, hatcheries producing bivalves to seed size only have emerged as an alternative to natural bivalve spawnings and also as a link in the aquacultural production of marketable bivalves.

### *Background*

Clams and oysters are harvested in all 14 states along the Atlantic Seaboard (U.S. Dept. of Commerce 1979). The most common oyster is the American Atlantic Oyster (*Crassostrea virginica*). Four species of clam-hard, soft, surf, and ocean quahog--constitute 99% of total U.S. landings by weight. The northern hard shell clam (*Mercenaria mercenaria*) accounts for 53% of the landings, although it constitutes only 17% of their value (Dressel and Fitzgibbon 1978).

Long Island, New England, and the Chesapeake Bay have historically been centers of clam and oyster fisheries (Ritchie 1977; Manzi *et al.* 1982b). Long Island Sound (the waters between Long Island and Connecticut), and Great South Bay, (the waters between Long Island and Fire Island), are the most productive of the New York areas (Figure 1). The seabeds there are hard, shallow and scattered with culch material to which the young animals attach (Korringa 1976). Additionally, the strong tidal currents carry abundant food into the bay (Bardach *et al.* 1972) and the bivalves are protected from many predators because the salinity is lower than in the surrounding ocean water.

Bivalves, especially oysters, have been an important food source for the New York

Eight coastal region. Oysters were used by Native Americans for food and trade and later by the colonialists (Terry 1977). Clams were fewer in number than oysters, but by the 1900's the rise of the cannery industry encouraged the growth of the clamping industry (Dressel and Fitzgibbon 1978).

The rise in clam production was interrupted after red meat shortages of the 1940's put increased pressure on the shellfish industry. Landings in the New York Marine District decreased from 6.8 million to 3.3 million pounds between 1943 and 1945 (Figure 2). Although landings later increased, the industry continued a pattern of boom and bust. In 1976 Long Island harvests reached a peak of 8.4 million pounds, worth about \$16.8 million (Freedman and Morris 1983). Great South Bay was considered the most important hard-shell clam industry in the world. It harvested 80% of Long Island's total landings, which represented 50% of the U.S. total (U.S. Dept. of Commerce 1974). Since then, the Long Island clam harvest has declined; by 1982, harvests were down by 60% from 1976 levels (Figure 3).


After a peak in 1904, oyster harvests in New York remained fairly stable between 1921 and 1952 but experienced a near-collapse in the 1960's (Figure 2). Revival of the industry through aquaculture has increased landings. Today, oyster production in the New York region is based entirely on aquaculture (Terry 1977).

### *Population Dynamics*

To understand the dramatic changes in the Long Island clam and oyster industries, one must examine the interaction among competing factors that determine the cyclic pattern of bivalve populations. Although population fluctuations occur naturally, they have been exaggerated through over harvest and destruction of natural habitat.

An inherent problem of common property fisheries is overuse (Hardin 1968), which leaves too few animals unharvested for the species to repopulate. This problem is acute around Long Island because bivalves in

**FIGURE 1. MAP OF LONG ISLAND COASTAL REGION**



Source: Terry (1977)

the cool waters require a long maturation period and because a large proportion of the population must be left unharvested in order to sustain the population. Thus, one major reason for the periodic decline of the clam and oyster populations is the likelihood that harvests will occasionally exceed the maximum sustainable yield.

Competition with man has decreased the natural habitat of clams and oysters and adversely affected bivalve populations. As urbanization and shoreline use in the New York area intensified, natural spawning grounds were reduced. Siltation, which is particularly harmful to young spat, increased; and saltwater intrusion caused by the removal of sandbars decreased protection against natural predators (Freedman and Morris 1983). Because toxins accumulate in the flesh of oysters and clams, the increased sewage effluent and agricultural runoff in the area have

increased mortality rates as well as decreased the marketability of the bivalve. The actual numbers of oysters and clams available for harvest have been reduced further through the closure of contaminated natural spawning beds (Figure 1).

#### *Demand Considerations*

While many factors affecting the bivalve population dynamics have resulted in a reduction in their supply, the industry has also been affected by demand considerations. For example, although the income elasticity of demand for clams is positive, it is relatively low, giving rise to only a slow growth in demand for clams over time due to rising per capita income and increasing human population (Gates *et al.* 1974). The income elasticity of demand for oysters is positive as well, but per capita consumption has actually decreased

FIGURE 2. COMMERCIAL LANDINGS FROM NEW YORK MARINE DISTRICT WATERS





FIGURE 3. LONG ISLAND CLAM HARVEST



Sources: McHugh and Williams (1976); Terry (1977);  
Regional Marine Resources Council (1974)

Source: Freedman and Morris (1983)

(Im and Langmo 1977). This decrease might have been caused by either the decrease in supply (Gates *et al.* 1974) and the associated rise in price, or by a change in consumers' tastes due to the fear of eating contaminated animals.

#### *Alternatives*

These problems have led to a disappearance of, or change in, the traditional shellfish industry. The scarcity of the hard clam has caused a shift to other species, such as surf clams (Dressel and Fitzgibbon 1978). Production has shifted southward to less-populated regions where competition for habitat is not so keen (Manzi *et al.* 1982a). Also, oyster imports have increased (Im and Langmo 1977).

Options to halt the gradual disappearance of the existing bivalve industry do exist. One option is a comprehensive government management plan to stop the decrease in, or even increase, the natural bivalve populations. Aquacultural production of marketable bivalves and hatchery production of bivalve seed are also viable alternatives, but they both require radical change in traditional harvesting techniques.

#### Regulation

Because most clam and oyster beds are common-property resources, they have suffered, as most common-property resources have, from over-use and abuse. Although size, time and method of catch are regulated, no consistent policy for predator control or return of culch material to the beds has been in force (Matthiessen 1970). Furthermore, these regulations have often been ineffective because of inadequate knowledge, failure to enforce existing laws, lack of cooperation and an absence of comprehensive planning.

The Regional Marine Resources Council's Shellfish Committee, created in the 1970's, partially filled the void by identifying 18 guidelines for improving the industry (Regional Marine Resources Council 1974). A Long Island clam union exists to aid members of the industry. On the consumer side, iden-

tification of potentially harmful pathogens, public health inspections and enforcement of closed beds will help maintain consumer confidence and increase demand.

To protect the industry, some Long Island governments have purchased young shellfish seed in order to seed public grounds (Smith 1982-83; Goldstein 1983). The population could also be increased by controlling predators, working the bottom for better spawning success, transferring animals to non-polluted grounds and controlling pollution, saltwater intrusion and siltation of habitat.

Unfortunately, implementation of such non-traditional resource policies is politically arduous and requires placation of special-interest groups and cooperation among affected parties. These measures are often expensive to establish and to enforce. The overall benefits to society from an increased bivalve population and the resulting strengthened harvesting industry must be measured against the costs of implementing any resource-regulating mechanisms.

#### Aquaculture

Aquaculture, the growing of organisms in water under controlled conditions, offers an alternative to traditional bivalve harvesting techniques. The primary benefits of controlling the environment are increased product supplies, although factors adversely affecting quality, such as predators, crowding and parasites, may be reduced.

The viability of an aquaculture industry may depend on the granting of exclusive rights to harvest or to develop a section of the ocean bottom aquaculturally. Marine resources have traditionally been common property (Terry 1977), but sedentary marine life on the bottom may be considered private property, although this right is difficult to substantiate (Hanson 1974; Terry 1977).

The states around the New York Bight area and towns on Long Island have a history of granting bottom leases to private individuals (Kochiss 1974; Terry 1977; Gates *et al.* 1974). In New York, the Department of En-

vironmental Conservation grants leases on the condition that the water quality is good and that no naturally-productive shellfish beds exist (Terry 1977). Great South Bay is held in trust by local townships; harvesters have full bottom rights to certain areas.

In general, economists support private ownership to increase efficiency and productivity (Agnello and Donnelley 1976; Terry 1977). Private beds experience five to ten times greater productivity (Dressel and Fitzgibbon 1978), probably because private harvesters can use less labor-intensive methods than can the public harvesters, who are required by the government to use inefficient harvesting techniques to control harvest quantity. Furthermore, without the assurance of exclusive rights to a harvest, there is no incentive for private harvesters to return culch, to control predators, to establish seed beds or to purchase seed to supplement the natural sets.

As stated above, the viability of the industry is demonstrated by the fact that all oysters on Long Island are currently produced by aquaculture (Terry 1977; Pillay 1976), and all oyster aquaculturists produce some hard clams (Korringa 1976). However, there has been increasing opposition led by local harvesters to the leasing of marine bottom, even of non-productive beds. Few leases have been granted recently (Terry 1977), and if these beds continue to remain in public control and open to all harvesters, the community must either exert better management controls or expect diminishing supply. On the other hand, private ownership might lead to an increase of marketable species but a decrease or disappearance of other naturally-occurring species.

### Hatcheries

Potential advantages to hatchery-produced seed include greater reliability of production and better quality control. The development of the hatchery industry (which grows clam and other larvae into bottom-dwelling seed large enough to be planted in open waters for later cultivation or harvest-

ing) has come in response to natural reproduction failure in open water (Loosanoff and Davis 1963a; Henderson 1978). Hatcheries can potentially achieve better quality control, decreased mortality and faster growth (Bardach *et al.* 1972; Donohue *et al.* 1981), thus offering aquaculturists a reliable source of bivalve seed. Furthermore, local governments can supplement declining natural spawns with hatchery-produced seed, a practice local harvesters generally favor as an alternative to direct governmental support of open-water growout in areas where a few individuals can affect the local bivalve market (Miller 1977; Rhodes 1974).

Worldwide demand for oyster and clam seed is growing at an increasing rate because of decreasing natural stocks and improving hatchery techniques. Henderson (1978) has estimated the annual world market for hatchery seed to be \$2 billion, or 25% of the annual U.S. whole clam and oyster harvests. In the U.S., hatcheries exist in the Southeast, the Northwest and in parts of the Northeast. The U.S. leads in seed production, having pioneered many clam hatchery techniques, but its oyster techniques are still considered "primitive" (Bardach *et al.* 1972).

Opinions differ as to the sufficiency of seed availability. Clam seed is more readily available than oyster seed (Terry 1977), except for seed larger than 10 mm (Castagna and Kraeuter 1977; Manzi *et al.* 1982a). Supplies of oyster seed have proven unreliable and insufficient to meet demand at current price (Donohue *et al.* 1981; Im and Langmo 1977). The demand for hatchery-produced seed is cyclical and increases during natural set failures (Henderson 1978). If hatcheries could produce a reliable, inexpensive product that is more attractive to growers than natural seed, they could capture a larger share of the seed market.

Hatcheries currently operate with less than complete information for two reasons: a) lack of cooperation between affected parties in the industry (Henderson 1978); and b) inasmuch as no two hatcheries are operated exactly alike (Terry 1977), such information as is available from one is not entirely appli-

cable to another. Partly due to this lack of information, hatcheries are only marginally cost-effective (Terry 1977). Cooperation should be improved between hatchery operators and buyers, researchers and the government (Henderson 1978). This in turn should affect the flow of information and help industry development.

Despite the problems faced by hatcheries, many in the field believe that they will continue as a viable industry (Terry 1977). Genetic breeding and development of domestic bivalve strains are future potentials of the industry (Terry 1977). The development and success of the hatchery industry depends on the continued demand for seed, encouragement by government and the scientific community and the success of hatchery operators. The success of the individual operator depends in large part on an ability to compete in the market place. To be competitive, the hatchery operator must be able to make the best production decisions based on an accurate assessment of production costs. This is the subject of the following sections.

### *III - HATCHERY PRODUCTION OF BIVALVE SEED*

Raising oyster and clam seed under controlled conditions requires technically-sophisticated, delicately-monitored production processes. The purpose of this section is to describe these processes, using the hatchery from which data were collected as a model. Although production systems vary among hatcheries, especially in algae and culch preparation and in facility and system designs, most operations are universal to all hatcheries. Techniques are varied primarily to take advantage of available natural resources at the site.

#### *The Hatchery*

The bivalve seed hatchery, from which data were collected for this study, was the outgrowth of an experimental plant begun in the late 1970's. This model facility was used

to help train staff and develop techniques utilizing the available natural resources in the surrounding area.

Construction of the permanent hatchery was completed in 1982. The building used for seed production has approximately 500 square meters of indoor floor space. There is an ample outdoor work area, as well as room available for expansion. The hatchery began partial production in 1982, and the first full production year was 1983. Both oysters and clams are grown, but production is now concentrated on oysters.

The hatchery is located a few hundred yards from a bay. The ownership of the bottom rights to part of the bay led to the development of a field operation ("outgrow") where some of the seed produced in the hatchery is grown to maturity. Most of these outgrow operations are conducted separately from those of the hatchery.

#### *The Hatchery's Production System*


The production process can be divided into six life-cycle stages for the bivalve:<sup>1</sup> 1) broodstock maintenance and conditioning; 2) spawning; 3) larvae development; 4) setting; 5) juvenile development and 6) field (or out-grow); and two support stages: 1) algae production and 2) culch production. Figure 4 shows the location of each production stage within the hatchery.

To maintain a low mortality rate, young bivalves must be treated with care during all phases of production, including inspection, cleaning and transfer. Exposure to severe environmental conditions or fouling by foreign organisms, such as fungus, bacteria and predators, can cause death or deformation of the animals or a poisoned product. Thus, particular attention must be given to cleanliness at each stage of production and to selec-

---

<sup>1</sup> Because the differences between clam and oyster production are minimal, the two processes will be treated as one throughout the remainder of this analysis.

**FIGURE 4. PRODUCTION LOCATIONS IN THE HATCHERY**



tion of non-toxic materials for all equipment that comes in direct contact with the animals. The quantity and quality of inputs used, such as energy, algae, culch and other materials, depend on the biological requirements of the

bivalve, the existing environmental conditions and on the hatchery's ability to purchase or produce those inputs. Inadequate or poor quality inputs will affect the survival of the bivalves.

As with most biological production processes, there is a trade-off between achieving maximum production and the cost of inputs. For instance, higher densities improve efficiency in the use of inputs such as water and labor; however, a high density decreases the survival and growth rates of the existing animals. The hatchery must weigh the marginal cost of providing ideal environmental conditions against the resulting increase in the value of the marginal output.


Proper timing of production is critical in two respects. First, if the hatchery begins production too early in the season, the animals will be ready for transfer to the juvenile tanks or outgrow facility before the water has reached a suitable temperature. Extending production too late may leave the hatchery with unsold inventory. Second, new spawnings and transfers between stages must be timed so as not to exceed the system's capacity in terms of space or ability to produce sufficient inputs. The optimal age of transfer

depends in part on the cost of maintaining the animal. The best way to understand the production system is to examine each of the components separately. (Figure 5 shows the flow between the production stages.)

#### Algae

Production of algae for feed in sufficient quantity and quality is important to the bivalve at all stages of growth and is one of the principal problems in profitably growing animals to market size entirely within a closed system (Gates *et al.* 1974; Epifanio 1975). Not much is known about the specific food requirements of the bivalve (Galtsoff 1964), especially about the adult bivalve (Epifanio 1975), but in the work that has been done, bivalve algae consumption is measured either by the number of cells consumed per animal or by the algae concentration made available to the animal. Existing estimates of the number of cells consumed by

FIGURE 5. FLOW BETWEEN HATCHERY PRODUCTION STAGES



oysters and clams at different stages of animal growth are in Table 1. Using the second and most common method, optimum algae concentrations for oyster larvae are given in Table 2.

The amount of algae assimilated, i.e., actually used for growth by the animal, is estimated at between 70% and 90% of the amount consumed (Epifanio 1975). Consumption is less than the filtering rate (the cells filtered per volume of water pumped), and the filtering rate in turn is less than the pumping rate. Filtering efficiency is affected by environmental conditions and was measured at between 34% and 58% by Baab and Associates (1973).

For bivalve feed, a mix of algae species is preferable (Ukeles 1975). The digestibility and nutrient value of the food depends on the chemical composition, size, texture, taste (Epifanio 1976), cell wall and toxicity (Ukeles 1975) of the algae species. Juveniles are less sensitive to algae quality and

size than are larvae, which require cells sized from 6-10 microns in diameter (Breese and Malouf 1975). Clams are generally less particular than are oysters about the quality of algae fed (Ukeles 1975).

The quantity of algae that the hatchery must produce to feed the bivalves depends on whether or not sea-water containing naturally occurring algae is made available to the animals. The proportion of its diet an animal is able to receive in this way depends on the concentration of algae in the seawater, on the pumping and filtering abilities of the animal, on the seasonal variation of that concentration and on the amount of water to which the animal is exposed. Also, some additional algae must be produced and held in reserve in case main supplies are contaminated. Producing excessive amounts can be costly, because algae can be stored only a few days.

The hatchery's algae stage is managed by a single employee, who spends more than one-half the total working hours exclusively

Table 1. Daily Consumption of Algae Cells by Oysters and Clams

Stage/Age of Oyster	Equivalent Stage at the Hatchery	Algae Cells Consumed Per Day	
		Oysters	Clams <sup>a</sup>
<b>Larvae</b>			
2 days		$1.0 \times 10^4$	
12-15 days	Larval	$4.0 \times 10^4$	
set size		$5.0 \times 10^4$	
<b>Spat</b>			
3 weeks	Setting	$2.0 \times 10^5$	$1.0 \times 10^5$
8 weeks		$3.0 \times 10^6$	$2.4 \times 10^7$
<b>Seed</b>			
	Juvenile	$8.0 \times 10^7$	$6.0 \times 10^5$
<b>Adult</b>			
3 inches	Juvenile/Outgrow		$1.0 \times 10^7$
		$1.1 \times 10^8$	

Sources: Claus and Adler (1970); Matthiessen and Toner (1966); Gates et al. (1974).

<sup>a</sup>The three consumption levels for clams refer to animals of 0.2, 2.0 and 8.0 mm in size, respectively. The corresponding stages for the oysters listed at the left are probably not exactly comparable to these sizes.

Table 2. Optimum Algae Concentrations for Oyster Larvae

Initial Larvae Shell Length ( $\mu$ )	Food Concentration for Highest Growth Rate ( $\mu\text{l}$ of packed cells/1)
77	2.5
78	2.5
104	10.0
104	10.0
139	10.0
146	20.0
201	40.0

Source: Rhodes and Landers (1973).

on algae production. The batch method, in which algae is cultured to a maximum density using successively larger containers, is used. Not all algae at a maximum age and density are used exclusively for food; instead, some are used to inoculate a larger tank.<sup>2</sup> This "modified batch" strategy requires additional labor (Matthiessen and Smith 1979).

The second-floor laboratory is where algae samples are analyzed for contamination and proper densities, nutrient solution is prepared to supplement the algae cultures, and initial algae stock is stored. Four different varieties are cultured: *Thalassiosira pseudoma* (diatom), *Pavlova lutheri*, *Pavlova lutherii Hoptaphyceae* and *Isochrysis galbana*. The latter represents 50% of the algae grown and 50% of the animal's diet. Because young animals require small algae cells, the diatom is used only during the juvenile stage.

<sup>2</sup> In contrast to techniques that centrifuge large quantities of seawater to concentrate naturally-occurring algae, this method selects only the best species of algae. The centrifuge process, which is commonly used on Long Island, is less expensive (Matthiessen and Smith 1979), but it requires an abundant and reliable source of naturally-occurring, desirable algae. The batch or modified batch process offers certain advantages for feeding efficiency because higher densities of algae can be obtained. With these processes, algae can be produced at densities of  $10^4$  cells/ml, as compared to densities in seawater of  $10^3$  cells/ml (Gates et al. 1974).

The primary algae production stage begins with tube cultures. Algae, in nutrient-rich well water, are allowed to mature two weeks to a maximum density of  $3 \times 10^5$  cells/ml. The culture is then transferred to sterilized flask containers. After one week, this culture matures and is transferred to carboys. Water must remain cool and sufficiently aerated. Cool fluorescent lighting provides the algae with artificial sunlight 24 hours each day.

The secondary stage, located in the algae room next to the laboratory, consists of seven 757-litre tanks. Each tank is inoculated with 16 litres of primary algae culture; nutrients and well water are added. The number of tanks used depends on near-term food requirements. Optimum algae density of  $4 \times 10^5$  cells/ml is reached in one week.

The secondary culture flows to the ground floor to innoculate the 3,785-litre tertiary algae tanks. The requirements at this stage are similar to those for previous cultures; and when a maximum density of 5 to  $8 \times 10^5$  cells/ml is reached, the tertiary, or "direct-feed", stage is used to feed the bivalve broodstock and the animals at the larvae and set stages. It is also used to innoculate the next culture stage. After one week, any algae remaining are discarded.

The final algae tank is a sunken 49,205-litre tank in the juvenile room. This "solar" tank is innoculated from the tertiary tanks and allowed to bloom under natural sunlight provided by large, sloping windows over the tank. Maintenance is similar to that in other stages. When a suitable density is reached, the entire amount is transferred to the algae holding tank located in the same room; the solar tank is then reinnoculated. The juvenile bivalves are fed directly from the holding tank. After one week, the remaining algae are discarded.

#### Broodstock Maintenance and Conditioning

The quality of the bivalve produced depends to a degree on the genetic quality of the parental stock, which is selected on the

basis of size, shape and growth potential. The broodstock is kept in rectangular 825-litre tanks of aerated, filtered water. Cool water temperatures are maintained to prevent spontaneous reproduction.

Before spawning can occur, the mature adults must be "conditioned"; the water temperature is increased to 26-29°C, and the food supply is also increased. The conditioning time varies according to the temperature and the amount of food used, but oysters require approximately 35 days (Gates *et al.* 1974).

#### Spawning

Once the male and female bivalves have been conditioned, the spawning process takes two employees one to three hours to complete. Conditioned adults are induced to spawn by either introducing a sharp temperature increase or by adding previously-collected sperm (Breese and Malouf 1975). One female clam can produce as many as 24 million eggs at a time (Loosanoff and Davis 1963b). Within 24 to 48 hours, the gametes become free-swimming larvae.

#### Larvae Development

During the larvae stage, algae-rich seawater is circulated about the mobile animals ("veligers"). In the larvae room are 38 conical, 340-litre larvae tanks and two 700-litre, flat-bottom larvae tanks. These tanks are filled with warm, filtered seawater, which must be heated when seawater temperature is below 25°C. Approximately 14 litres of algae are added to each tank/day.

Oyster and clam larvae are maintained close to ideal densities for fastest growth and survival, viz., densities from 0.4 to 15 animals/ml and 5 animals/ml of water, respectively (Gates *et al.* 1974; Matthiessen and Toner 1966). Growth and survival rates at this and subsequent stages are affected not only by density and by food quantity and quality, but also by the temperature and available oxygen. Thus, water is maintained at the ideal temperature, and aeration is supplied continually to all tanks.

Maintenance of the larvae is labor-intensive. Besides daily feeding and monitoring, every 48 hours the tanks are cleaned, the larvae are sorted by size, and the dead animals are removed. This takes two employees nearly a full day to complete, depending on the number of tanks in use. Animals of similar size are collected together from different tanks and are then returned to a newly-filled tank, and algae are added. This culling process is an essential feature of the production process because it, in large part, determines the quality and the quantity of the final output.

The most critical period for the larvae occurs at metamorphosis, when the bivalves change from free-swimming larvae to bottom-dwelling spat. Setting occurs at 7 to 10 days for clams (Bardach *et al.* 1972; Gates *et al.* 1974) and 10 to 20 days for oysters (Gates *et al.* 1974). The clams at this stage measure 200 to 215 microns (Bardach *et al.* 1972; Gates *et al.* 1974) and oysters 300 to 325 microns. Prior to this metamorphosis, the animals must be transferred to the set tanks, which have been filled with culch.

#### Culch

Culch is the material onto which the oyster larvae set and attach permanently. Thus, before the larvae can be transferred to the setting tanks, culch is prepared. The choice of culch material depends on the type of grow-out anticipated rather than the relative cost of the materials (Matthiessen 1970; Gates *et al.* 1974).<sup>3</sup> Before 1960, whole oyster or clam shell was used exclusively for bottom culch material (Matthiessen and Smith 1979); other techniques, such as the suspended-string method, could be used in lieu of bottom culch. Today, choices for culch include a variety of shell sizes, artificial materials and "culchless" oysters.

The culchless oyster is an oyster with no attached culch, that is, one that has been induced to set on a material, such as plastic

---

<sup>3</sup> Clams do not require culch since they do not attach firmly.

sheeting, from which it can be removed easily. Culchless oysters can be grown at greater densities than the culched variety inasmuch as clumping, which crowds and deforms the spat on the culch, is less of a problem. Culchless oysters may be less expensive to produce because culch production costs are avoided, and because shipping and handling of the culchless oyster is less costly. However, once transplanted, the culched oyster has a better chance of surviving, especially in waters with high siltation (Matthiessen 1970). Thus, culchless oysters (and clams) should be raised to larger sizes in the hatchery, potentially offsetting any other savings.

The preparation of culch is tedious, taking about 12 minutes to produce one litre of sorted, crushed shell. The prepared culch is then cleaned and spread on the bottom of the set tanks at densities of 200 to 350 ml/square foot.

#### Setting

Prior to setting, the larvae are transferred to eight 825-litre tanks filled with warm, filtered, aerated water at densities of 2 to 18 million animals per tank. Maintenance is similar to that for larvae; the same algae mix is used, although cleaning is less frequent (every 72 hours). The animals are fed 70 litres of algae/day/tank during the first week; this is increased to 140 litres by the second week.

Because the animals are delicate at this stage of their life, their transfer too early to the juvenile tanks could reduce the overall survival rate. The hatchery does not heat the juvenile tanks and must wait for the bay water to heat sufficiently by natural means before young spat can be transferred. However, if the animals remain too long in the set tanks, they cannot receive the large quantities of flowing water and algae necessary for rapid growth, nor can the tanks be used for subsequent batches. Animals as small as 300 microns and as large as 1.9 mm have been transferred out of the set tanks in the hatchery.

#### Juvenile Development

From the set tanks, the young spat are transferred into either the juvenile tanks or the upwell system, or they are sold. Most oyster and clam seed are sold at about 3 mm; smaller animals than this would experience too low of a survival rate.

The juvenile tank has a capacity of 26,500 litres. The culched oysters are placed in trays that measure 0.5 x 0.6 m; these trays are then placed on the bottom of the juvenile tank. The tank is filled with enough sieved bay water to cover the trays and is aerated. Approximately 75 litres of algae are added per day to achieve a cell count of  $2 \times 10^6$  cells/ml. The tank is cleaned twice per week, at which time the animals are washed and culled for death, fouling, and deformation. By the time the animals are ready for sale or transfer to the field, there has been a significant reduction in their numbers.

The upwell system is used only for clams and culchless or micro-culched oysters. The spat are held in screened-bottomed buckets, which are suspended over 825-litre rectangular tanks. Water from the bay holding tank is siphoned into these tanks, creating a continuous flow of water that aerates the spat. This sieved water is at bay temperature: 17°C in May and 28°C in August. Algae from the holding tank are added at a rate of about 7,600 litres per day. Animals have been kept in this system until they have reached 2.3 to 8 mm in diameter, and then they have been sold or transferred to the field.

#### *The Hatchery's Physical Systems*

The facility was designed for efficiency and flexibility to allow for changes in production techniques in response to new scientific information, environmental conditions or market changes. Simple changes in internal arrangements, made possible by movable tanks that can be used for a variety of functions, can streamline production. The hatchery was designed to facilitate easy cleaning. Energy, water and air systems were coordinated to maximize efficiency.

### Energy System

The amount and the type of energy used in the hatchery depend on the production method and on the prevailing environmental conditions. Natural lighting and heating are used wherever possible. Because the bay water temperature is cool, the hatchery must heat large quantities of water to fill the brood, larvae and set tanks.


The energy flow is complex (Figure 6). Heat inputs include electricity for lighting and machinery and fuel oil for running the boilers to heat water and ambient air. Cooling inputs include air conditioners, bay water and ambient air. The relative contribution required from each of these sources to maintain ideal water and air temperatures is a function of the season. It is possible to determine the amount of electricity and fuel oil required if initial temperatures, final temperatures, and heating and cooling capacities are known.

### Water System

Water is used in all phases of production: for all animal stages, for algae production and for cleaning. Water, the quality of which is critical to all hatchery operations (Bardach *et al.* 1972), can be sterilized with ultraviolet light or with chemicals. Fresh water for cleaning is obtained from a reservoir at no cost to the hatchery, and three pumps bring water from the bay (Figure 6). The first pump is for deep-well water, which is used exclusively for algae production. It is the purest of the three bay waters and is never heated. Pump 2 filters out particles greater than 1 to 5 microns; this water is used for larvae, brood and set tanks. The third pump delivers water to the juvenile room and sieves out particles greater than 700 microns.

The facility is designed to minimize the electrical costs of pumping. By using a series of different levels, water can be siphoned for various uses once it is within the

FIGURE 6. ENERGY AND WATER FLOW WITHIN THE HATCHERY



facility. Gravity is also used for the transfer of algae from the secondary to the tertiary stages and from the tertiary to the solar stage.

#### Air Flow System

If water is not kept in constant flow, the hatchery must provide aeration. Although the hatchery's air system is not elaborate, it is essential to algae and animal growth. Depending on size, all tanks are provided with a certain number of air tubes, except for the upwell system, which receives a constant flow of water. Air is pumped through these tubes from an air compressor that operates 24 hours per day. To reduce electricity consumption, the hatchery replaced its 5-HP air compressor with a 1-HP compressor, which has proven sufficient for the hatchery's needs.

#### *Costs and Input Considerations*

Before the decision to build a hatchery is made, a complete financial assessment should be conducted, including estimates of the initial investment cost, operating expenses and expected earnings. To date, existing hatcheries were estimated to have an average initial capital investment for building and equipment greater than \$100,000 (Matthiessen and Smith 1979), but technically-advanced hatcheries, such as this one, may cost even more. Using land with direct access to the ocean can add considerably to the total price, although in this case, the land was already owned.

Hatchery operating costs were estimated at between \$2,700 and \$10,000 per month for most facilities (Matthiessen and Smith 1979). The actual amount depends on the hatchery's output and design, on the existing environmental conditions, and on unforeseen production risk beyond the control of the operator.

To date, there are only a handful of published estimates on the cost of bivalve seed, and the average costs per animal range from \$0.0005 (Im and Langmo 1977) to \$0.01 (Matthiessen 1979) for smaller seed and from \$.029 to \$.25 (Gates *et al.* 1974) for larger seed. The most comprehensive estimates have been made in recent years by Im and Langmo

(1977), Im *et al.* (1976) and Bolton (1982). They have attempted to estimate costs at each stage of production (e.g., brood, larvae etc.) using an economic engineering approach based on small, experimental hatcheries both on the Pacific and Atlantic coasts.

These studies are an important first step, but they fail to determine the cost of producing animals to different sizes in the same stage. This information is particularly critical at the final stage, that is, just prior to sale or transfer to outgrow facilities, because an aquaculturist purchases seed from a hatchery at a size in accordance with his production method, locality and production goals. It appears that 3 mm is a common size at which to sell the spat. Those who purchase larger seed face significantly less risk of mortality in open waters since they are more resilient to predation, siltation and climatic changes (Matthiessen and Smith 1979).

When larger seed are produced, it is usually because a) the hatchery cannot find buyers for seed not previously contracted for sale (usually the hatchery overproduces some animals to leave a margin of error), b) environmental conditions are such that the hatchery cannot transfer seed to their own "growout" facility or c) it is experimenting with growth techniques for larger animals. In other words, production to a larger size is not necessarily desirable but may be preferable to disposal of the animals if the hatchery can cover its costs at the margin.

#### Classifying Productive Inputs

An important first step in estimating the cost of raising bivalve seed to different sizes is to identify the input requirements for each stage of production. As outlined in Table 3, production of bivalve seed can be separated into five production stages and two support stages: 1) broodstock conditioning, 2) spawning, 3) larval development, 4) setting, and 5) juvenile development; and 1) culch and 2) algae preparation. Because input quantities change over time, they will also change over the duration of most stages. Therefore, in this analysis, the stages are broken down into smaller units of time, or periods. At the end of a period, there is an associated size and age

Table 3. Inputs to Bivalve Seed Production in the Hatchery

Inputs	Condition	Stages <sup>a</sup>						Inputs	Stages <sup>a</sup>					
		Production					Support		Production					Support
		1	2	3	4	5	1	2	1	2	3	4	5	1
ELECTRICITY														
Water pumping														
Wellwater														
Filtered water	x	x	x	x			x							
Sieved water	x				x	x								
Air compressor	x	x	x	x	x	x		x						
Air cooling														
Lab room							x							
Algae room							x							
Air heating														
Office								x						
Algae room								x	x					
Larvae room		x	x	x				x						
Back room	x			x				x						
Equipment														
Office							x							
Laboratory							x							
Autoclave							x							
Crusher					x									
Sorter					x									
Lighting														
Office							x							
Laboratory						x	x							
Algae room					x									
Larvae room						x	x							
Back room						x								
Juvenile room						x	x							
FUEL OIL														
Water heating														
Heat exch.	x	x	x	x										
Filtered water	x	x	x	x										
Sieved water					x	x								
Space heating														
Office							x							
Algae room							x							
Larvae room		x	x	x										
Back room	x		x											

<u>LABOR</u>														
Tank maintenance	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Culling			x	x	x	x								
Feeding	x		x	x	x	x								
Transfers		x	x	x	x	x								
Cleaning							x						x	x
Management								x					x	x
Laboratory work	x	x	x	x	x	x	x		x	x	x	x	x	x
Equipment maintenance														
<u>MATERIALS</u>														
Antibiotics											x	x	x	x
Cleaning								x	x	x	x	x	x	x
Glassware									x	x	x	x	x	x
Lightbulbs									x	x	x	x	x	x
Screening									x	x	x	x	x	x
Office supplies									x	x	x	x	x	x
Fresh water									x	x	x	x	x	x
Nutrient solutions									x	x	x	x	x	x
<u>ALGAE</u>														
Direct-feed tanks	x		x	x	x	x	x	x	x	x	x	x	x	x
Solar tanks							x	x	x	x	x	x	x	x
From natural seawater								x	x	x	x	x	x	x
<u>CULCH</u>														
Micro									x	x	x	x	x	x
Large									x	x	x	x	x	x
<u>MISCELLANEOUS</u>														
Travel and communications											x	x	x	x
Insurance											x	x	x	x
Permits											x	x	x	x
Shipping							x	x	x	x	x	x	x	x
Loan payments											x	x	x	x
Taxes											x	x	x	x

<sup>a</sup>See text and Figure 5 for identification of stages. General hatchery inputs are independent of stage. Intermediate inputs are produced in support stages.

of animal, as determined by the growth rate experienced in the hatchery. The cost per animal at any age can be estimated by measuring the inputs that are used in a period.

As with most production systems, inputs can be classified as either fixed or variable. Fixed inputs are those whose quantity cannot be changed during the length of time under consideration since the cost of quick

variation in their amount is large as to make such variations impractical (Mansfield 1979). Here, we are interested in smaller seed (produced in less than 4 months) as well as in larger animals that must be left in the hatchery longer. For convenience, the length of time under consideration is taken to be one year. Thus, fixed costs are assumed to include depreciation on building and equipment, taxes, and rental charges.

Some inputs are also fixed once a batch is started and so either do not vary with output or vary insignificantly. These inputs include general hatchery maintenance, such as for machinery and general hatchery cleaning. It is difficult to isolate these costs in terms of their applicability to any particular stage or period of the bivalve's life.

Variable costs differ in how much they are affected by the level of production output. First, because of the nature of hatchery production, which uses a modular-style (e.g., separate tanks), many inputs are only secondarily related to the number of animals within the system. For example, a larvae tank requires a certain amount of labor for cleaning and a certain amount of heated water no matter how many animals are in each tank even though the number of tanks used does depend on the total number of animals. Second, some inputs are related to the area where the animals are located and must be used no matter how many tanks are in that area, such as space heating. Third, some events take place only once and are not significantly affected by the number of animals, for example, sampling. Finally, there are those inputs, such as algae consumption, that vary directly with output.

It is the inputs that vary according to the number of animals in the system that are of greatest interest to this analysis, since it is these inputs that vary most with animal size. Inputs that do not vary directly with output are shared according to the relative length of time that the animal is retained in the hatchery, rather than the total output or the animal's size.

For this study, inputs are divided into six primary categories: electricity, fuel oil, labor, materials, algae and culch (Table 3). Although both algae and culch are support stages in the system, they are intermediate inputs that could be produced outside the hatchery and purchased.

In most hatcheries, it is not difficult to separate the total monthly operating costs into the total cost of each primary input. The real difficulty is in allocating a share of these primary input costs to different stages of pro-

duction. Im, Johnston and Langmo (1976) made some progress on such a breakdown in costs. A summary of their results is presented in Table 4. These data highlight the importance of algae production as a component in the variable costs of bivalve production and serve as a benchmark by which to calibrate the results from the bivalve cost simulation program described in the next section.

#### ***IV - SIMULATING BIVALVE PRODUCTION COSTS***

This section describes a simulation model by which one may calculate input requirements, net output and the cost of growing bivalve seed. Within the model, primary input costs are allocated to different stages of production, thus facilitating the comparison of the marginal cost of growing bivalve seed to a larger size with the expected marginal revenue of doing so. The model was designed to facilitate its coding as an interactive computer program that could be used by hatchery operators.

##### *Structure of the Model*

The model can be used to calculate production costs, inputs and output for an entire batch of bivalves. Even though most hatcheries would have concurrent batches at different stages of development, focusing on a single batch makes it possible to allocate input costs to the different production stages and to generate the information necessary to determine the optimal timing of production. It is also assumed that only one production pathway is followed for each application of the model; that is, all animals in all stages are treated similarly regarding the choice of culch, timing of transfers and all other aspects of the production processes.

To compute inputs and costs for animals of different ages, each batch is initially divided into the production stages discussed above. These stages are subdivided into shorter periods so that costs can be determined at various points in time within a stage. This is particularly useful for long stages, such as the juvenile stage, where a more precise estimate of the change in cost with age is required. All periods within a

Table 4. Production Costs in Various Production Stages in Bivalve Hatcheries of Two Sizes<sup>a</sup>

Stage	Percentage of Total Cost		Percentage of Variable Cost	
	Plant 1	Plant 2	Plant 1	Plant 2
Conditioning	2.7	0.2	3.9	0.2
Spawning	5.0	0.3	8.7	0.3
Larvae Rearing	15.0	27.8	21.4	25.1
Larvae Setting	7.2	18.8	7.6	13.5
Algae Production	38.9	33.3	29.4	39.0
Culch Preparation	3.7	9.3	7.2	14.7
Miscellaneous	27.5	10.3	21.8	7.1

<sup>a</sup> The source of the data is Im, Johnston and Langmo (1976). Plant 1 is an experimental plant with an output of 15 bushels/week. Plant 2 is scaled up to a projected 800 bushels/week.

stage are of equal length, but the length and number of periods may differ amongst stages.

The number of bivalves surviving and their size are calculated at the end of each period, along with the per day averages over the period. The quantities and costs are calculated for each period and are averaged over each period as well. The costs incurred in the current period are added to those of previous periods to accumulate total costs up to the end of the period. This type of structure, in which there is a succession of repeated computations, lends itself easily to FORTRAN, the computer language in which the model is coded.<sup>4</sup>

As shown in Figure 7, the options to run the code are chosen in step 1a, and data, including price of inputs, are entered or computed in steps 1b-1g. After all data are entered, a batch is begun on the first day of the first month of production (step 2). Each stage *i*, beginning with the conditioning stage, is looped through steps 3-14, and within this stage loop, period *j* is incremented (steps 5-13). The passage of time during the course of a batch is accumulated within *j* (step 6), and thus the age of the bivalves is advanced.


With this nested looping structure, any computation within the *j* loop will depend not only on *j*, but also on time, stage and the month the batch is begun.

Survival rates for bivalves (step 7), size of bivalves (step 8) and input quantities (step 11) are computed within the loop *j*, calling function subroutines, which are placed outside of the loop when used by more than one subroutine. Cumulative and average costs, based on information from previous steps, then are computed (step 12). The program output contains information generated about the batch beginning at month *M* (step 15); a new batch is begun on the first day of month *M+1* (step 16). By performing computations for each month, information about batches begun at different times during the production season can be generated.

The options included in the code allow the model to be adapted to fit a variety of different hatcheries. With little effort, the internal parameters of the code can be re-specified to change the number, units and name of inputs, and the number and name of stages, thus adapting the code to a hatchery's specific production methods. When running the program, the user may specify the length of each stage and the number of periods within that stage, thus determining in part the level of detail contained in the information generated. It is possible to skip entirely stages for which information is not required or for which data may be insufficient.

<sup>4</sup> Preliminary development of the code, which concentrated on specific subroutines, is shown in Myers (1985). A listing of the completed code is given in Appendix A. It is being run on an IBM mainframe computer. Data can be entered entirely by file, entirely interactively, or by a combination of both.

FIGURE 7. BIVALVE CODE FLOWCHART



The user may also choose the way in which input prices and quantities are determined. If the code contains the necessary algorithms, the user may choose either to allow the program to compute input price and/or quantities (*steps 1f, 11b*) or enter these as fixed parameters (*steps 1d, 1f*). The modular design of the code facilitates the addition or

deletion of these price and quantity algorithms.

#### Algorithms

The algorithms for bivalve survival rates, bivalve size and production costs are

required in the model and are described here.<sup>5</sup> In addition, the optional algorithms for algae price and quantity, which constitute the major focus of this analysis, are outlined. These algorithms, except those for price, are contained within the period loop and thus depend on time, period and stage. The prices for inputs used in a period are functions of the elapsed time up to period  $j$  and on the month during which that period occurs.

#### Survival Rate

The output from a batch of bivalves, as well as the quantity of inputs and production costs, depends on the initial number of bivalves in the batch and on the number of animals surviving from period to period. Death of a proportion of the population in a hatchery is unavoidable even though the mortality rate is lower than in the wild. Survival rates are affected by environmental conditions maintained in the tanks, but little is known about the trade-off between increasing the survival rates and cost of improving the conditions in the tanks. Survival rates will also vary by hatchery, although in all hatcheries survival should increase at an increasing rate because the animals become more resilient with age and culling rates go down with increasing age.

Hatcheries often overproduce at early stages to account for the declining numbers of animals over time, which is a result of natural mortality and the culling of inferior animals from the batch. Culling assures a faster overall growth rate and a superior final harvest. However, it is difficult to distinguish between the effects of mortality and culling on the number of remaining animals because both dead and inferior animals are removed in the same sorting process.

It is reasonable to assume that the number will decline over time. There are, however, few, if any, data on actual survival

rates under these controlled conditions. At the time data were collected for this study, the hatchery had not been in production long enough to have reliable data on survival rates. For this reason, a general exponential function reflecting a constant percentage mortality rate over time was written into the code. This allows the user to specify particular survival rates and facilitates testing the sensitivity of costs to changes in survival rates.

The number surviving at the end of each period is computed at step 7 of the code by calling a function that relates survival at time  $t$  to the number of animals initially in the batch, i.e., the number successfully spawned.<sup>6</sup> This function is:

$$(1) \quad N_t = N_o(t)^\rho,$$

where  $N_t$  = the number of animals remaining at time  $t$ ;  $N_o$  = initial number of animals spawned; and  $\rho$  = a parameter to be set by the user,  $0 > \rho > -1$ .

While this general functional relationship is thought to be an accurate representation of an invertebrate population's survival rate (Emmel, 1976, p. 205), its tendency to go to infinity as  $t$  goes to zero caused a problem in writing the computer code. To be consistent with the use of  $t$  as an index of the passage of time for other calculations, it was necessary to have equation (1) at  $t=0$  record the initial number of animals in the batch ( $N_o$ ). Thus for convenience of programming,  $N_t$  was defined as the number of animals remaining at the end of any week  $t$ , where  $t$  is incremented in discrete time. Thus, to reconcile the initial estimate of the number of bivalves with the continuous time function above and the other functions in the model, the following function is used:

$$(2) \quad N_t = N_o(t+1)^\rho.$$

---

<sup>5</sup> The specific forms of the mathematical relationships used to describe the algorithms were chosen primarily because they were found to provide the "best" statistical results given the data used to estimate the parameters of the functions empirically. The empirical estimates are described in subsequent sections.

<sup>6</sup> Although the cost to condition the adult population (stage 1), which spawns the initial egg population, affects the total cost of producing a batch, the number of adults is not important in determining the initial population in this model, because  $N_o$  is input by the user as a fixed parameter rather than as a function of the fecundity of the adults spawned.

An estimate of the average number of animals surviving during a period is necessary to calculate an average quantity of inputs used during  $j$ . The actual average number over the period from time  $a$  to  $b$  is calculated by:

$$(3) \quad N_{(a,b)} = \frac{N_o}{(b-a)} \int_a^b (t+1)^{\rho} dt.$$

This integral was estimated in the code with a Reiman sum using the trapezoidal approximation to the area under a curve. The method gives a satisfactory estimate of the integral and increases the generalizability of the code since it does not depend on the actual form of the integrand.<sup>7</sup>

#### Size

Both the size of an individual bivalve at the end of each period and the average size during the period are estimated. This information is useful in assessing potential market price and is also necessary for computing the amount of inputs used within a period.

Data relating the size of bivalves to their age were difficult to find in the litera-

<sup>7</sup> To estimate the definite integral  $\int_a^b f(t)dt$  by the trapezoidal rule, one must calculate

$$A = \int_a^b f(t)dt \approx \sum_{k=1}^{n-1} f(t_k) \Delta t + \frac{f(a) + f(b)}{2} \Delta t.$$

For coding purposes, it was efficient to calculate

$$A = \sum_{k=1}^n f(t_k) \Delta t + \frac{f(a) - f(b)}{2} \Delta t$$

and  $\Delta t = 1/7$ , a daily increment. If  $f$  is continuous on  $a \leq t \leq b$  and twice differentiable on  $a < t < b$ , then there is a number  $a < \epsilon < b$  such that

$$\int_a^b f(t)dt = A - \frac{(b-a)}{12} f''(\epsilon) \Delta t^2$$

There is no set procedure for finding the value of  $\epsilon$  such that this is true, but one can obtain some idea of the magnitude of the error by investigating the size of  $f''(t)$  between  $a$  and  $b$  (Thomas, 1960, pp. 207-17 and pp. 385-88).

ture as well. However, Epifanio, Logan and Turk (1976) and Epifanio (1975) report a small amount of data relating the size  $H_t$  (in mm of shell height) to age in weeks ( $t$ ) for both clams and oysters. These data suggest that the relationship is quadratic, at least for young bivalves:

$$(4) \quad H_t = \alpha t + \beta t^2,$$

where one would expect  $\alpha \geq 0$  and  $\beta < 0$ , (the parameters are chosen by the user). The only potential problem with an empirical specification of this function is that it may yield a maximum size at too early an age. This problem is not serious because the function behaves well over ages that are likely to exist in a hatchery setting.

Using this relationship, the average size during any period starting at time  $a$  and ending at  $b$  is given by:

$$(5) \quad AH_{(a,b)} = \frac{1}{(b-a)} \int_a^b (\alpha t + \beta t^2) dt.$$

This integral also is approximated in the code using the trapezoidal rule, which computes the value of the function at each day of the period.

#### Costs

Estimating the variable cost to produce a single bivalve to a certain age or size involves calculating the total variable cost ( $TVC_t$ ) for an entire batch at some time  $t$  and dividing by  $N_t$ , the number of animals surviving is given by:

$$(6) \quad AVC_t = TVC_t / N_t.$$

This average variable cost per animal is derived at the end of each period.<sup>8</sup>

As stated above, each batch is initially divided into stages ( $i=1,\dots,S$ ), which are in turn divided into periods. To facilitate cod-

<sup>8</sup> The stage associated with the period must be greater than 1 since no animals are produced in the conditioning stage.

ing, and allow for a different number of periods in each stage, periods ( $j=1, \dots, J$ ) are indexed consecutively over the entire batch. Thus total variable costs at time  $t^*$  (measured in weeks) is the sum of all variable costs ( $PTVC_j$ ) across all periods included in the first  $t^*$  weeks:

$$(7) TVC_{t^*} = \sum_{j \in t^*} PTVC_j.$$

Because the code keeps track of periods and stages by incrementing the  $j$ th loop within the  $i$ th loop, costs up to the end of a period may also be thought of as the sum of the costs in all periods included in the completed stages ( $i=1, \dots, S^*-1$ ) plus the sum of the costs for all completed periods up to time  $t^*$  included in the current stage,  $S^*$ ,

$$(8) TVC_{t^*} = \sum_{j \in (S=1, \dots, S^*-1)} PTVC_j + \sum_{j \in (S^* \cap t \leq t^*)} PTVC_j.$$

Variable costs during a period are of course determined by the quantity of the inputs used during that period and by the price of the input during the month in which the input is used. This is incorporated into the model as:

$$(9) PTVC_j = \sum_{n=1}^N \sum_{k=1}^K \sum_{r=1}^R P_{nkm_j} Q_{nkrj},$$

where  $P_{nkm_j}$  = price per unit of input  $n$ ,

type  $k$ , in the month during which  $j$  occurs;  $Q_{nkrj}$  = quantity of input  $n$ , type  $k$ , for use  $r$  used during  $j$ ;  $N$  = number of inputs  $n$ ;  $K$  = number of types  $k$  of input  $n$ ;  $R$  = number of uses of input  $n$ , type  $k$ ; and  $m_j$  = month during which  $j$  occurs.

#### Prices and Quantities

As discussed previously, prices and quantities are either computed by algorithms within the code or are input by the user, thus allowing the user to focus on specific parts in the production process. Although the model allows any of the input prices to be computed, this option is probably most useful in computing the cost, or internal price, of algae and culch, because these two inputs are often

intermediate inputs produced within the hatchery. This analysis focuses on the computation of this internal price for algae and on the use of this input as feed during the production of the bivalves.

As shown in equation (9), the model allows for a price for  $K$  different types of input  $n$  in each month of production, thus creating a 3-dimensional price array,  $P_{nkm}$ . The code requires a value for each element of the array regardless of whether the price is computed or input by the user so that a corresponding price can be found for inputs used during any month.

From the computations for the quantity of inputs used in each period, where a quantity is calculated for each type of each input for each period,  $Q_{nkj}$ , another 3-dimensional array is constructed. Determining the quantity of each type involves summing over the  $R$  different uses for the input. For expository purposes, this is included as an additional subscript on the quantity variable in equation (9),  $Q_{nkrj}$ . However, in the code, this sum of input quantities by use is calculated in the quantity subroutine, and this subscript does not appear in the quantity array used in the cost calculations.

If the quantity of input is entered by the user rather than computed by algorithms in the code (*step 1d*), a lump sum is entered as  $Q_{nki}$ , which is then divided equally among each period of that stage (*step 11*).

#### *Algae Price and Quantity*

The decision to focus this report on algae production (price) and on the quantity of algae used for feed in bivalve production is justified for three reasons.

First, the production of algae in the hatchery was the most well defined and firmly established of the hatchery's production processes. It was easy to separate the inputs used in the production of algae from those used as direct inputs to bivalve production, because some different types were used and because the two processes were physically separated in the hatchery.

Second, the algae production model closely resembles the bivalve code structure because of the similarity in the production processes. By first writing the algae price code, potential problems with the full bivalve code could be anticipated.<sup>9</sup>

Third, and most important, is the fact that feed costs have been identified as the largest single component of hatchery costs, ranging from 29% and 33% of variable costs in early studies (Im *et al.* 1976) to as high as 85% in more recent studies (Bolton 1982). Thus, from a practical point of view,


... the major bottleneck to commercialization of intensive bivalve culture systems [lies] in the ability to economically culture massive quantities of these suitable algal species. Intensive study of problems associated with such mass culture of algae is certainly warranted... (Epifanio 1975, p. 190).

These concerns become even more important as hatcheries attempt to raise animals to a larger size.

The algae system, described in Section III, can be broken into stages of relatively short, fixed duration, thus distinguishing different algae "types" by the age of the culture. Algae can be fed to the bivalves from different stages and as such provides one example of how different types of one input can be used within the code. These different types of algae are also distinguished on the basis of their different physical characteristics, such as density, and overall purity, and by the cost of production. In the subroutine where algae costs are computed, a price, which in this case equals production costs, is computed for each of the different algae types.

Figure 8 is a representation of the algorithms used to simulate algae production. As in the bivalve program, only one batch of algae is followed at a time. It is also assumed that only one algae species, or a single mixture of species, is produced in a batch and that all tanks within a stage are used. By assuming that a new batch is begun on the first

FIGURE 8. ALGAE PRICE SUBROUTINE



<sup>9</sup> In fact, the two processes are very similar inasmuch as both operate on a batch-like principle. The transfer of algae to successively larger tanks as they mature is comparable to the transfer of animals between stages. Both systems are a function of the number of animals or cells; however, the number of organisms increases over time in the algae system due to cell reproduction, although cell size remains constant.


of every month, one can derive the 3-dimensional price array required by the bivalve program,  $P_{nkmj}$ , from equation (9) (where  $n =$  algae), from the internal cost of producing each type of algae.

The cost of producing any stage of algae depends on the amount of inputs --electricity, labor and materials-- used during the stage and on the price of those inputs during the month in which the stage occurs. Because algae production in the hatchery is separate from bivalve production, there is a provision in the data input to specify different prices for those inputs used in producing algae from those used in other parts of bivalve production. The algorithms used to derive the internal costs of producing algae are discussed in detail in Myers (1985).

The algae quantity subprogram is used to compute the feed requirements for a batch of bivalves at any time  $t$  (see Figure 9). It allows for the feeding of algae from any stage of maturity to bivalves at any stage of growth, although it is likely that average production costs per unit of algae would be too high in the early stages. Exactly at what stage during the algae production process algae should start to be used for feed is an empirical question that can only be answered by the information generated from the program itself. More is said about this issue below.

To estimate the algae required for a batch of bivalves, one must know the relationship between feed consumption and some measure of a bivalve's age or size, in addition to the number of animals remaining in the batch at time  $t$ . In the literature, one finds feed requirements related directly to age (Epifanio 1975) and directly to size, as measured by shell height or dry weight (Epifanio *et al.* 1975 and Bolton 1982). To maintain the flexibility of the program, both options were included in the code. This was accommodated easily, because regardless of the relationship used --size and feed or age and feed-- the data are consistent with a function that is linear in logarithms. One might well interpret the relationship as a Cobb-Douglas production function.

FIGURE 9. ALGAE QUANTITY SUBROUTINE



In the first option contained in the code, the daily consumption of algae (10,000 cell units) at time  $t$ ,  $C_t$ , is related to shell size in mm of shell height as determined from equation (4). Thus,

$$(10) \quad \ln(H_t) = \delta + \gamma \ln C_t$$

Solving for  $C_t$  and substituting,

$$(11) \quad C_t = (e)^{-(\delta/\gamma)} (H_t)^{1/\gamma}.$$

Multiplying this expression by the number of animals surviving ( $N_t$ ) and substituting from equations (7) and (9), the total daily feed requirements (in 10,000 cell units) for a batch at time  $t$ ,  $C_t^*$  can be written entirely in terms of  $t$ :

$$(12) \quad C_t^* = [(\alpha t + \beta t^2)^{1/\gamma}] [(t+1)^\rho N_o] [e^{-\delta/\gamma}].$$

The total number of algae cells consumed (in 10,000 cell units) during a period is then given by

$$(13) \quad \bar{C}_t = e^{-\delta/\gamma} \int_a^b [\alpha t + \beta t^2]^{1/\gamma} [(t+1)^\rho] dt.$$

Again, the values of the parameters are left to the user.

The second option relates the cells consumed,  $c_t$ , directly to age in weeks, thereby eliminating the need to evaluate equation (4). The new relationship is<sup>10</sup>

$$(14) \quad \ln t = \xi + \eta \ln c_t.$$

Thus,

$$(15) \quad c_t = (e)^{-(\xi/\eta)} t^{1/\eta};$$

$$(16) \quad c_t^* = [t^{1/\eta}] [(t+1)^\rho N_o] [e^{-\xi/\eta}];$$

and

$$(17) \quad \bar{c}_t = e^{-\xi/\eta} \int_a^b [t^{1/\eta}] [(t+1)^\rho N_o] dt.$$

Both of these options create the 2-dimensional array for algae quantity.<sup>11</sup> The price determined in the algae price routines can then be referenced to compute the cost of the algae inputs used in each period of bivalve production.

<sup>10</sup> The variables  $C_t$  and  $c_t$  refer to feed consumption (in 10,000 cell units), calculated according to options 1 and 2, respectively.

<sup>11</sup> Both equations (13) and (17) are evaluated in the program using procedures similar to those for the survival and size equations above and reference equations (2) and (4) for each day of the period.

## V - EMPIRICAL TESTS OF THE SIMULATION MODEL

The purpose of this section is to describe the results from initial experimentation with the bivalve cost simulation model. The decision to apply the model to oyster seed production (*Crassostrea virginica*) was based primarily on the availability of data. As emphasized above, the empirical results reported here focus on simulating the cost, or internal price, of algae and on the quantities of algae and total feed costs of raising a batch of bivalves to various sizes. The remainder of this section begins with a discussion of the data used for the analysis. Next, the empirical results are presented for a base case, after which a variety of sensitivity analyses are conducted to see how the empirical results are affected by changes in the key parameters of the model.<sup>12</sup> Finally, these results are compared with other cost and bivalve seed price information to provide initial indications of the optimum size at which to market the seed.

### The Data

To experiment with the model, it was necessary to obtain two sets of information: a) the data required by the subprogram that calculates the cost or internal price of algae and b) data by which to estimate the parameters of the survival function, the age-size relationships and the algae feed functions [equations (1), (4), (10) and (14)]. The data for the algae price (internal cost) estimates are based in large measure on several weeks of observation at the hatchery during the summer of 1983. There were, however, a few items taken from other sources, but these were modified to be consistent with the other production processes and specific environmental conditions of the hatchery. These data are summarized in Appendix B and are described in detail by Myers (1985).

For any single type of bivalve, there were very few data in the literature on which to base the estimates of the parameters for

<sup>12</sup> Myers (1985) conducts considerable sensitivity analysis on the internal price of algae, so these results focus on changes in other parameters of the system.

equations (1), (4), (10) and (14). However, except for the survival equation, it was possible to combine data from several sources to derive consistent parameter estimates for the oyster.

Table 5 contains an empirical estimate of equation (4a), relating the size of oysters to age. The data are from Epifanio *et al.* (1976).

Within the simulation model, there are two alternatives for estimating the feed (algae) requirements for bivalves. One way is to relate feed directly to shell height as in equations (10a,b) of Table 5. To estimate equation (10a), equation (4a) was used first to predict shell height at some selected ages for which Claus and Adler (1970) report algae consumption data. These predicted shell heights are then regressed on the algae consumption figures (in log-linear form) to obtain equation (10a). Equation (10b) was estimated directly from data in Bolton (1982). A second way to estimate feed requirements accommodated in the model was to relate algae consumption directly to age. Data from Claus and Adler (1970) were used to estimate equation (14a) directly, while equation (14b) was estimated directly from information in Bolton (1982).<sup>13</sup>

The primary reason for developing these four ways of estimating feed requirements was to test the sensitivity of the model. It is somewhat surprising, but encouraging, to find that the algae production elasticities in these four equations are similar, ranging only from 0.35 to 0.42.

Data with which to estimate the survival rate parameters were among the most difficult to obtain. There were no population

<sup>13</sup> Bolton (1982) estimates oyster feed requirements through a mathematical relationship between the whole weight of oysters and the daily consumption of algae (due to Pruder *et al.* (1977)):

$$Y = 8.2 x^{-0.21}$$

where  $x$  = whole weight in grams; and  $Y$  = algal cells cleared  $\times 10^8$  /g whole weight/day. Because they report data on age and shell height for oysters of different whole weights, once this equation is used to estimate algal consumption based on weight, these predicted values can in turn be regressed on age and shell height to obtain equations (10b) and (14b).

Table 5. Estimated Equations for Use in Simulations

Equation Number <sup>a</sup>	Estimated Equation
(1a) <sup>b</sup>	$\ln N_t = 8.55 - (46.09) \cdot 0.255 \ln t$ $R^2 = 0.66$ [0.041]
(4a) <sup>c,d</sup>	$H_t = 0.737(t) - (22.14) \cdot 0.00252(t)^2$ [0.033] [-6.99] [0.0004]
(10)	
(a) <sup>f</sup>	$\ln H_t = -0.631 + (-1.65) \cdot 0.338 \ln C_t$ $R^2 = 0.81$ [0.074]
(b) <sup>e</sup>	$\ln H_t = -1.769 + (-5.85) \cdot 0.416 \ln C_t$ $R^2 = 0.98$ [0.022]
(14)	
(a) <sup>e</sup>	$\ln t = -0.331 + (-0.85) \cdot 0.345 \ln C_t$ $R^2 = 0.81$ [0.075]
(b) <sup>d</sup>	$\ln t = -2.244 + (-5.644) \cdot 0.407 \ln C_t$ $R^2 = 0.97$ [0.029]

<sup>a</sup> See text for equation numbers and variable definitions. The numbers in parentheses and brackets are t-ratios and standard errors, respectively.

<sup>b</sup> The source of the data is Eldridge, *et al.* (1979).

<sup>c</sup>  $R^2$ 's are not applicable for restricted regressions.

<sup>d</sup> The source of the data is Epifanio *et al.* (1976).

<sup>e</sup> The source of the data is Bolton (1982).

<sup>f</sup> The sources of data are Claus and Adler (1970) and Epifanio *et al.* (1976).

survival data for oysters raised in a hatchery, although Bolton (1982) makes allowance for a 20% mortality rate during the cultivation period. Because most assumptions in Bolton's experiment proved extremely optimistic, it seemed inadvisable to use this overall mortal-

ity rate as a base of comparison. Furthermore, there was no easy way to infer from this overall rate how mortality changes over time.

Since hatchery data on which to estimate survival rates of oysters or other bivalves were unavailable, the best alternative to obtain survival rates that are at least in the relevant range was to estimate equation (1) from data for hard clams planted in protective trays in a South Carolina estuary (Eldridge *et al.* 1979). The size of the clams planted was larger than bivalves raised in a hatchery, and the survival function for smaller bivalves had to be extrapolated beyond the limits of the data. Initially, one might expect this to lead to an overestimate of the survival of hatchery-sized bivalves. However, the data do reflect a less controlled environment than found in the hatchery, a factor that may partially offset the effect of size. The estimated equation, in log-linear form, is in Table 5; its parameters imply a constant mortality rate of 25.5%. When accumulated on a weekly basis, this function leads to over 60% mortality in a year's time, three times that for oysters in Bolton's (1982) prototype hatchery. This discrepancy is large, but it is unlikely that Bolton anticipated a year-long cultivation period. Equation (1a) from Table 5 implies about a 50% mortality after 18 weeks. This is closer to Bolton's initial assumptions. These two extremes are a useful range over which to test the sensitivity of the results.

#### *Empirical Results*

A computer program was designed that implements the simulation model described in detail above. This program was designed to be used by the hatchery operator to generate information about a batch of bivalves at different states of development. Input costs and requirements, and bivalve age, survival and size are reported at the end of each production period and as cumulative amounts. Most important, it gives an average cost per bivalve at the end of each period to give the user information about production timing and about the expected revenue from bivalve seed sale that might be expected.

#### Initial Simulations

A partial listing of the printout generated by the program is in Appendix B. It contains information for a batch of oysters under base run conditions (run 1). The batch is assumed to begin in April and last for 420 days, or 59 periods. Myers' (1985) input and price data were used to calculate the internal cost per unit of algae.<sup>14</sup> The data reflect economic conditions at the hatchery in 1983. The batch contains one million oysters initially; the survival rate for bivalves is given by equation (1a), of Table 5; the age-size and feed-size relationships are given by equations (4) and (10a), respectively. The bivalve feeding filtering rate is assumed to be 50%, and 12% of the diet is assumed to come from algae in circulating sea water in the juvenile stage.

The assumptions underlying the base run and 11 other simulations are given in Table 6. These runs contain numerous combinations of the functional relationships described earlier in Table 5. The first three runs assume the same mortality rates but accommodate different ways of estimating the physical feed requirements. Runs 1 and 3 assume feed is directly related to size, although in run 2, feed is directly related to age. The results of these runs are summarized in Table 7.

For the base run, the number of bivalves surviving over the duration of the batch falls from a million at spawning to an estimated 938,000 at the beginning of the larval stage. After setting, the estimated number is 716,000. If the batch were to remain in the hatchery for 420 days (including 60 days for conditioning the broodstock), the

---

<sup>14</sup> To facilitate comparisons, all other runs of the model are assumed to begin in April and run for 59 periods (420 days). The simulation was run over such a long duration primarily to test the behavior of the model since, due to bivalve size and the cost of feed, it is clear that bivalve seed would be sold or transferred to grow-out long before this last period was reached. Conditioning takes a month, and the spawning, larvae and setting stages were broken into periods of a few days each; but once the juvenile stage was reached, periods were a constant one week in length. For ease of exposition, much of the discussion in the text focuses on monthly intervals or on individual stages.

Table 6. Description of Parameters for Alternative Simulations Runs<sup>a</sup>

Simulation	Survival Equation	Age-Size Equation	Feed-Size Equation	Feed-Age Equation
Run 1 (Base Run)	$N_o = 1 \times 10^6$ $\rho = -.0255$	$\alpha = 0.737$ $\beta = -0.00252$	$\delta = -0.631$ $\gamma = 0.0338$	N.A.
Run 2	same as run 1	same as run 1	N.A.	$\xi = -0.331$ $\eta = 0.345$
Run 3	same as run 1	same as run 1	$\delta = -1.769$ $\gamma = 0.0416$	N.A.
Run 4	$N_o = 1 \times 10^6$ $\rho = -0.214$	same as run 1	same as run 1	N.A.
Run 5	$N_o = 1 \times 10^6$ $\rho = -0.296$	same as run 1	same as run 1	N.A.
Run 6	$N_o = 1 \times 10^6$ $\rho = -0.163$	same as run 1	same as run 1	N.A.
Run 7	$N_o = 1 \times 10^6$ $\rho = -0.337$	same as run 1	same as run 1	N.A.
Run 8	same as run 1	$\alpha = 0.671$ $\beta = -0.00252$	same as run 1	N.A.
Run 9	same as run 1	$\alpha = 0.671$ $\beta = -0.00332$	same as run 1	N.A.
Run 10	same as run 1	same as run 1	$\delta = -0.631$ $\gamma = 0.412$	N.A.
Run 11	same as run 1	same as run 1	$\delta = -0.631$ $\gamma = 0.486$	N.A.
Run 12	same as run 1	$\alpha = 0.671$ $\beta = -0.00332$	$\delta = -0.631$ $\gamma = 0.486$	N.A.

<sup>a</sup> See Table 5 and equations (1), (4), (10) and (15) in the text for parameter definitions and estimated relationships.

number of animals surviving would be approximately 357,000.

The estimated average size of the bivalves ranges from just under 2 mm after the setting stage to just over 33 mm at the end of the simulation. As the batch progresses through time, the additional cost of feeding larger bivalves is partially offset by declining

numbers. Because the feed cost per surviving bivalve includes the accumulated costs of feeding those that did not survive, the costs rise rather rapidly throughout the batch. At the end of period 18 (about 100 days), the feed cost per animal is still less than one cent, but if left in the hatchery through the 59th period, feed costs would average more than a dollar per surviving bivalve.

Table 7. Summary of Simulation Output for Initial Runs

Period <sup>a</sup>	Stage	Cumulative	Animals	Average	Cumulative	Cost per	Bivalve
		Days to End of Period	Sur- viving (thousands)	Size of Animal (mm)	Run 1 <sup>b</sup> (Base) (\$)	Run 2 (\$)	Run 3 (Proportion of Run 1)
6	Setting	49	716	1.98	c	0.98	8.77
10	Juvenile	77	594	4.84	c	0.97	5.72
14	Juvenile	105	534	7.61	0.003	0.98	4.29
18	Juvenile	133	495	10.30	0.009	0.99	3.59
22	Juvenile	161	467	12.91	0.022	1.01	3.16
26	Juvenile	189	446	15.44	0.045	1.04	2.86
30	Juvenile	217	429	17.89	0.082	1.06	2.63
34	Juvenile	245	414	20.26	0.137	1.09	2.46
38	Juvenile	273	402	22.55	0.214	1.12	2.31
42	Juvenile	301	391	24.76	0.316	1.16	2.20
46	Juvenile	329	382	26.88	0.449	1.19	2.10
50	Juvenile	357	373	28.93	0.616	1.23	2.01
54	Juvenile	385	366	30.90	0.822	1.27	1.94
59	Juvenile	420	357	33.24	1.144	1.32	1.86

<sup>a</sup>Period 6 is through setting (about 50 days including adult conditioning, spawning and larval stages); the remaining periods are a week in length.

<sup>b</sup>See Table 6 for assumptions about runs.

<sup>c</sup>Less than 0.001.

Data for the other runs in Table 7 suggest that these results can be quite sensitive to the assumptions made regarding the feed equations. In the base run, feed requirements are related directly to the size of the animal, whereas in run 2 it is assumed that feed requirements are determined by equation (14a) relating feed directly to age. Assuming the same survival rates, this alternative way of calculating feed leads to a slightly lower estimate of feed cost per surviving animal through period 18, but then cost rises to 1.3 times that of the base run by period 59. The feed relationships in these two runs were estimated from the same basic data; despite these differences, the results are quite consistent.

The cumulative costs from run 3 are in sharp contrast to those for runs 1 and 2. In this latter run, costs are consistently higher than in the base case, and within the same

run the differential is most extreme in the early periods. This result is somewhat surprising, given that the production elasticity for algae is higher in runs 3, 4 and 6 than in the base run. However, the intercepts on the size or age axes of the production function are lower than for the base run. Thus, even though the percentage gain in size is larger for each one-percent change in algae consumption, absolute feed requirements to raise these oysters to any size over the duration of the batch remains higher than in the base run.

The implications of this result lie in the importance of measuring feed requirements accurately in early stages of the growth process. That is, if one were to look only at a single animal, the tendency would be to conclude that feed costs would be relatively unimportant compared to costs for older animals. However, because the number of animals falls throughout the process, costs (due

to what appear to be minor cost differences in early periods) accumulate rapidly as more animals die and as total costs are averaged over fewer survivors. Thus, reliable estimates for both feed costs at different points in the growth process and the number of animals surviving are more critical in estimating costs for raising bivalves than for larger animals (e.g., cattle or other livestock) with much lower mortality rates.

#### A Digression on Algae Production

The primary focus of the analysis so far has been on the costs of feeding a batch of bivalves and the effect of survival and feeding ratios on costs. However, it is evident (Table 8) that cumulative algae consumption in runs 2 and 3 is a smaller multiple of the base run consumption than it is for cost in early periods and is a larger multiple

in later periods. The primary reason for this is that feed costs for bivalves are also a function of the implicit price or cost of algae that is raised for feed. The program uses algorithms from Myers (1985) to calculate the price of algae, which are assumed to be constant in any month but which can vary by month because of seasonal differences.

The "algae price" algorithm assumes that algae are produced in a batch process, which is divided into 6 stages covering about 6 to 7 weeks. During the first four stages, most of the algae are needed to inoculate the subsequent stage. Thus, the net production of algae available is negligible until the tertiary stage (about 6 weeks). Up to this point the algae are grown in well water and the batch is quite pure, thus making it ideal for feeding to bivalves at the larval and set stages. These tertiary algae tanks are located (in the study's hatchery) adjacent to the larval tanks, which

Table 8. Algae Consumption for Selected Simulations

Period <sup>a</sup>	Cumulative Algae Consumption			Ratio of Cost to Algae Consumption <sup>c</sup>	
	Run 1 <sup>b</sup> (Base Case) (10 bil. cells)	Run 2 (Proportion of Run 1)	Run 3	Run 2	Run 3
6	48	0.98	8.05	1.00	1.09
10	580	0.97	4.79	1.00	1.19
14	2,153	0.98	3.68	1.00	1.17
18	5,067	1.01	3.10	1.00	1.16
22	9,513	1.04	2.73	0.97	1.16
26	15,603	1.07	2.47	0.97	1.16
30	23,382	1.11	2.27	0.95	1.16
34	32,846	1.15	2.12	0.95	1.16
38	43,947	1.20	2.00	0.93	1.16
42	56,605	1.25	1.89	0.93	1.16
46	70,711	1.30	1.81	0.92	1.16
50	86,133	1.36	1.74	0.90	1.16
54	102,720	1.42	1.67	0.89	1.16
59	124,850	1.50	1.61	0.88	1.16

<sup>a</sup>See Table 7.

<sup>b</sup>See Table 7.

<sup>c</sup>Ratio of column 7 (Table 7) to column 3 (Table 8) and column 8 (Table 7) to column 4 (Table 8), respectively.

helps keep the costs of moving algae around the hatchery to a minimum. There are an estimated 9,650 litres of algae available at the tertiary stage, most of which is fed to the larvae and animals at set. Some of the tertiary algae is also used to innoculate a solar stage to which seawater, rather than well water, has been added, thus making it unusable for the sensitive larvae. After another week or two, there are an estimated 67,000 litres of algae available to feed juvenile bivalves. Because the cost per cell of algae in the solar stage is about 25% of that in the tertiary stage, it is not surprising that in the early periods feed cost is higher relative to algae consumption than it is in later periods (Table 8).

A further explanation for the behavior of costs relative to algae consumption lies in a slightly higher algae cost estimate in the summer months because of the air conditioning required to maintain cool temperatures suitable for algae production. For example, assuming the batch is started in April, about

82% of the cost of algae in the solar stage is due to labor, 5% to materials and 13% to electricity, much of which is used for cooling. Thus, as the bivalve production moves into autumn, cooling requirements are reduced and production costs fall by as much as 3% relative to a batch available in July.

#### Implications of Different Survival Rates

The accuracy of hatchery cost estimates derived by this analysis depends on the number of animals surviving at any stage. To test the sensitivity of these results, the survival parameter was changed by plus and minus one and two standard derivatives from the least squares estimate of equation (1). A decrease in the absolute value of the parameter, as in runs 4 and 6, implies a lower mortality rate, whereas an increase in its absolute value leads to higher mortality (runs 5 and 7).

The results of these sensitivity tests are summarized in Table 9. The change in

Table 9. Summary of Simulations to Test Sensitivity of Survival Parameter

Period <sup>a</sup>	<u>Number of Bivalves Surviving</u>				<u>Total Feed Cost</u>				<u>Cumulative Feed Cost Per Bivalve</u>			
	<u>Run</u>				<u>Run</u>				<u>Run</u>			
	4 <sup>b</sup>	5	6	7	4	5	6	7	4	5	6	7
	(Proportion of Run 1)				(Proportion of Run 1)				(Proportion of Run 1)			
6	1.06	0.95	1.13	0.89	1.05	0.96	1.11	0.91	0.99	1.01	0.98	1.02
10	1.09	0.92	1.21	0.85	1.08	0.93	1.18	0.87	0.99	1.01	0.97	1.02
14	1.11	0.90	1.25	0.82	1.10	0.91	1.23	0.84	0.99	1.01	0.98	1.02
18	1.12	0.89	1.29	0.80	1.11	0.90	1.26	0.82	0.99	1.01	0.98	1.02
22	1.13	0.89	1.32	0.78	1.12	0.90	1.29	0.80	0.99	1.01	0.98	1.02
26	1.14	0.88	1.34	0.77	1.13	0.89	1.31	0.79	0.99	1.01	0.98	1.02
30	1.15	0.87	1.36	0.76	1.13	0.88	1.33	0.78	0.99	1.01	0.98	1.02
34	1.15	0.87	1.38	0.75	1.13	0.88	1.34	0.77	0.99	1.01	0.98	1.02
38	1.16	0.86	1.39	0.75	1.14	0.87	1.36	0.76	0.99	1.01	0.98	1.02
42	1.16	0.86	1.40	0.74	1.15	0.87	1.37	0.76	0.99	1.01	0.98	1.02
46	1.17	0.86	1.42	0.73	1.15	0.87	1.38	0.75	0.99	1.01	0.97	1.02
50	1.17	0.85	1.43	0.73	1.15	0.86	1.39	0.75	0.99	1.01	0.97	1.02
54	1.18	0.85	1.44	0.72	1.16	0.86	1.40	0.74	0.99	1.01	0.97	1.02
59	1.18	0.85	1.45	0.72	1.16	0.86	1.41	0.74	0.99	1.01	0.97	1.03

<sup>a</sup>See Table 7 for explanation of periods.

<sup>b</sup>See Table 6 for assumptions about runs.

the number of surviving bivalves is predictable directly from the parameter changes. Because of the form of the survival function, mortality is assumed to occur at a compound rate. Thus, for increases in mortality, the proportion of bivalves surviving, relative to changes in the base run, falls as the batch moves to later stages. Total feed costs (relative to the base run) fall as well, but not quite so fast. Exactly the opposite is true when mortality rates are reduced.

Perhaps the most important result from this analysis is that despite significant differences in the number of surviving animals (from 0.72 to 1.5 times the base run), cumulative costs per bivalve are never any more than plus or minus 3% of the base run costs. The important implication of this result for hatchery owners is that relatively accurate estimates of per-unit variable costs can be made somewhat independently of good estimates of survival. This is critical in compar-

ing per-unit costs to the potential price of seed to determine when to sell.

#### Sensitivity of the Relationship of Age and Feed to Size

Although the average cost per bivalve is relatively insensitive to the assumptions made about survival rates, the results of the initial simulations suggest that this is not the case for the relationships between age and size and between feed and size. The importance of these relationships is perhaps best understood by systematically changing the parameters of equations (4) and (10a), assuming the other assumptions in run 1 are constant. The results of this experimentation are summarized in Table 10.

To begin the analysis, the coefficient on the linear term was changed by plus and minus one and two standard deviations. It was readily apparent that although the change

Table 10. Sensitivity of Simulation Results to the Age-Size and Feed-Size Relationships

Period <sup>a</sup>	Run <sup>b</sup>	Average Size of Bivalve				Cumulative Feed Cost per Bivalve <sup>c</sup>				
		Run 8	Run 9	Run 10	Run 11	Run 12	Run 8	Run 9	Run 10	Run 11
		(Proportion of Run 1)					(Proportion of Run 1)			
6	0.91	0.91	1.00	1.00	0.91	0.76	0.75	0.58	0.40	0.33
10	0.91	0.90	1.00	1.00	0.90	0.75	0.74	0.38	0.20	0.17
14	0.91	0.90	1.00	1.00	0.90	0.75	0.73	0.29	0.13	0.10
18	0.91	0.89	1.00	1.00	0.89	0.75	0.72	0.25	0.09	0.07
22	0.90	0.88	1.00	1.00	0.88	0.75	0.71	0.22	0.08	0.06
26	0.90	0.88	1.00	1.00	0.88	0.74	0.70	0.20	0.06	0.05
30	0.90	0.87	1.00	1.00	0.87	0.74	0.69	0.18	0.06	0.04
34	0.90	0.86	1.00	1.00	0.86	0.74	0.67	0.17	0.05	0.04
38	0.90	0.86	1.00	1.00	0.86	0.74	0.66	0.16	0.05	0.03
42	0.90	0.85	1.00	1.00	0.85	0.73	0.65	0.15	0.04	0.03
46	0.90	0.84	1.00	1.00	0.84	0.73	0.64	0.15	0.04	0.03
50	0.89	0.83	1.00	1.00	0.83	0.73	0.63	0.14	0.04	0.03
54	0.89	0.83	1.00	1.00	0.83	0.72	0.61	0.14	0.03	0.02
59	0.89	0.82	1.00	1.00	0.82	0.72	0.60	0.13	0.03	0.02

<sup>a</sup>See Table 7 for explanation of periods.

<sup>b</sup>See Table 6 for assumptions about runs.

<sup>c</sup>Total feed costs as a proportion of those in run 1 are the same as the feed costs per bivalve.

in feed costs was more dramatic than the change in bivalve size, both remained about proportional to the change in the parameter,  $\alpha$ . Thus, only one of these runs, where  $\alpha$  is reduced by two standard deviations, is reported (run 8). This change reduced the estimated size by about 90% of the base run size while reducing feed costs to between 72 to 76% of base run costs.

The next step in the analysis was to change the coefficient on the  $t^2$  term in equation (4). In run 9, the coefficient on the linear term was the same as in run 11, but the absolute value of the coefficient on the squared term was increased by two standard deviations. The effect of this change on both size and feed costs was minimal in the early stages that would be most relevant to a hatchery producing seed to be sold or transferred to growout facilities. At later stages, the effect of this parameter change becomes increasingly important.

The most significant results from Table 10, however, seem to be the large reductions in feed costs due to an increase in the production elasticity of algae in equation (10a). By raising the production elasticity to one standard deviation above its original level, costs fall to just under 60% of the base run (run 10) in period 6 and to 13% by the end of the simulation. Increasing the production elasticity by an additional one standard deviation leads to further, but not quite so dramatic, cost reductions (run 11).

#### Preliminary Observations about Total Costs of Production

Because data were unavailable from the hatchery on other aspects of the production of oyster seed, it was impossible to implement all parts of the simulation model and estimate costs other than feed costs. However, at least two previous studies contain estimates of the proportion of total costs accounted for by algae feed. On the basis of these estimates, it is possible to make some preliminary observations about total production costs and thus about when during the batch costs will exceed the price that a hatchery operator might receive for the seed.

As reported in Table 4, Im *et al.* (1976) estimate that algae feed costs are about one-third of total production costs; Bolton (1982) reports estimates ranging from 15 to 85% of total costs. Feed costs, as a percentage of total costs, probably increase with age. However, the nature of this relationship is not known; so to obtain some idea of what this wide range in estimates implies for the simulation results above, two kinds of sensitivity analysis were conducted. The first was to assume that feed costs are a constant fraction of total costs (first three panels of Table 11). By assuming feed costs are 33, 59 and 85% of total costs, one is able to reflect all but the low range of Bolton's estimates. It was believed that this low end of the range would apply only to very young bivalves. To reflect increasing relative feed costs, the last two panels in the table assume that costs rise linearly from 15 to 85% of feed costs from period 1 through 59 and period 1 through 30, respectively.

These sensitivity results have important implications for marketing bivalve seed in that Matthiessen and Smith (1979) argued that oyster seed (2 to 3 mm in size) could be purchased at from \$2.00 to \$5.00 per thousand. Assuming these prices, base run conditions and feed costs at 33% of total, seed would have to be sold prior to the 11th period (5.5 mm shell height) to cover costs if the price were \$2.00 per thousand, and before the 13th period (6.9 mm shell height) if the price were to be \$5.00 per thousand. This "window of opportunity" would occur much earlier (periods 5 and 6) under the most costly conditions (run 3), but only slightly later (periods 14 and 17) for the least cost scenario (run 10).<sup>15</sup>

These results are remarkably consistent, given the wide range of parameters in the age-size and size-feed relationship reflected in these runs. The "marketing win-

---

<sup>15</sup> Since this is a "batch process" and per bivalve feed costs are relatively insensitive to survival rates, it's unlikely that these marketing opportunities would change much if different survival rates were assumed. This does not mean, however, that survival rates are unimportant in terms of the overall profitability of the batch.

Table 11. Alternative Estimates of Total Cumulative Costs of Production (\$ Per Oyster)

Period <sup>a</sup>	Run 1 <sup>b</sup>	Run 2	Run 3	Run 8	Run 9	Run 10
<u>33.3%<sup>c</sup></u>						
6	d	d	0.009	d	d	d
10	0.001	0.001	0.061	0.001	0.001	0.001
14	0.008	0.008	0.275	0.006	0.006	0.002
18	0.026	0.026	0.808	0.020	0.019	0.006
22	0.065	0.066	1.845	0.049	0.046	0.014
26	0.135	0.140	3.584	0.100	0.094	0.027
30	0.247	0.262	6.247	0.183	0.169	0.045
(0.002) <sup>e</sup>	11	11	5	12	12	14
(0.005)	13	13	6	14	14	17
(0.010)	15	15	7	16	16	21
<u>59%<sup>c</sup></u>						
6	d	d	0.005	d	d	d
10	0.001	0.001	0.034	0.001	0.001	c
14	0.004	0.004	0.155	0.003	0.003	0.001
18	0.015	0.015	0.456	0.011	0.011	0.004
22	0.037	0.037	1.041	0.027	0.026	0.008
26	0.076	0.079	2.022	0.057	0.053	0.015
30	0.139	0.148	3.526	0.103	0.095	0.025
(0.002) <sup>e</sup>	12	13	6	13	13	16
(0.005)	15	15	6	16	16	20
(0.010)	17	17	8	18	18	24
<u>85%<sup>c</sup></u>						
6	d	d	0.004	d	d	d
10	0.001	0.001	0.024	d	d	d
14	0.003	0.003	0.108	0.002	0.002	0.001
18	0.010	0.010	0.317	0.008	0.007	0.003
22	0.026	0.026	0.722	0.019	0.018	0.006
26	0.045	0.055	1.404	0.039	0.037	0.010
30	0.097	0.103	2.448	0.072	0.067	0.018
(0.002) <sup>e</sup>	13	13	6	14	14	18
(0.005)	16	16	7	17	17	22
(0.010)	18	18	9	20	20	26
<u>15-85%<sup>c</sup></u>						
6	d	d	0.015	d	d	d
10	0.002	0.002	0.078	0.001	0.001	0.001
14	0.008	0.008	0.298	0.006	0.006	0.002
18	0.024	0.024	0.758	0.018	0.018	0.006
22	0.054	0.054	1.522	0.040	0.038	0.012
26	0.100	0.103	2.642	0.074	0.070	0.020
30	0.164	0.174	4.161	0.122	0.112	0.030
(0.002) <sup>e</sup>	11	11	5	11	11	14
(0.005)	13	13	5	14	14	18
(0.010)	15	15	6	16	16	21
<u>15-85%<sup>c</sup></u>						
6	d	d	0.011	d	d	d
10	0.001	0.001	0.055	0.001	0.001	d
14	0.006	0.005	0.197	0.004	0.004	0.002
18	0.016	0.015	0.480	0.012	0.011	0.004
22	0.033	0.033	0.935	0.025	0.023	0.007
26	0.060	0.062	1.584	0.044	0.041	0.012
30	0.100	0.103	2.448	0.072	0.066	0.018
(0.002) <sup>e</sup>	12	12	5	12	12	15
(0.005)	14	14	6	15	15	20
(0.010)	17	17	6	18	18	25

<sup>a</sup>See Table 7 for explanation of periods.<sup>b</sup>See Table 6 for assumptions about runs.<sup>c</sup>These are the percentages that feed costs are assumed to be of total cost. In the last two panels, costs are assumed to increase linearly from 15 to 85% of feed cost from periods 1 through 30 and periods 1 through 59, respectively.<sup>d</sup>Less than 0.0005.<sup>e</sup>Actual period during which costs reach \$0.002, \$0.005 and \$0.010, respectively, per bivalve.

dow" is extended by only 2 or 3 weeks if feed costs actually turn out to be 85% rather than 33% of total production costs or if feed costs are assumed to rise to 85% of total costs over 30 rather than 59 periods. A doubling of the price (from \$0.005 to \$0.01 would extend the marketing period by only an additional 2 to 3 weeks as well.<sup>16</sup>

It is also important to recognize that during these "break-even" marketing periods, it is estimated that the size of the seed is much larger than the 2 to 3 mm on which the prices are based. This implies that the hatchery owners may be in an advantageous bargaining position to sell seed that meets the size requirements earlier, thus reducing costs, or by asking for a higher price for larger seed, with its higher survival rates during growout.

#### *VI - SUMMARY AND CONCLUSIONS*

The bivalve aquacultural industry in the northeastern United States has been strengthened in recent years by declining and fluctuating natural harvests in the surrounding coastal region. The hatchery industry, in which the bivalves are grown only to a small seed size, has gained acceptance as an important link in the aquacultural process; the firms in the industry are potentially able to provide a consistent, reliable source of seed with which the bivalve aquaculturist begins the production process. They have also gained acceptance as a seed supplier to local governments, who artificially seed depleted natural bivalve beds with the seed purchased from hatcheries. Hatcheries as an industry are likely to play a larger role in the overall production of bivalves if they can overcome some of the technical and economic problems with which they must contend.

The techniques involved in the production of bivalve seed in hatcheries have evolved only recently, yet are technically sophisticated compared to other open-water aquacultural techniques. The production pro-

cesses differ, but each requires a precise sequence of steps under carefully-controlled and -monitored environmental conditions. Mature adult clams or oysters taken from hatchery-maintained broodstock are induced to spawn. The gametes are transferred into large, conical larvae tanks where they freely circulate in warm, algae-rich seawater. The concentrated algae used for feed are grown in the hatchery or taken from seawater. After several weeks, the bivalves undergo a metamorphosis where they change from free-swimming larvae to bottom-dwelling spat. Prior to this time, the animals are transferred to set tanks where they are induced to attach to culch -- a material purchased or produced in the hatchery generally from crushed bivalve shell. Once the animals have passed through the delicate setting period, they are transferred to the juvenile tanks where they can be exposed to large quantities of algae and seawater. The animals remain in these tanks until they reach a suitable size for transfer to open-waters.

There are a number of production options available to the hatcherist, the most apparent being the design of the production system and the source of inputs. The algae production method is the greatest cause of variations affecting cost among hatcheries. These methods include the centrifuging of algae cells from seawater, continuous culture methods, or a batch process in which production is separated into distinct stages. The type of culch used and the number of animals produced as culched or culchless (attached or non-attached to the culch) also differ among hatcheries. The hatcherist has a great deal of discretion regarding the optimum survival rate of the animals in relation to the quantity and quality of inputs used. Finally, and most central to this research, is the choice of when to sell seed.

Despite the growth of the industry and advancements in the technical processes of hatcheries, little is known about the economics of the hatchery. There have been several important studies on this aspect of hatcheries (Im, *et al.*, 1976 and by Im and Langmo, 1977). These and other studies fail to pay sufficient attention to the wide variation that exists among hatcheries and to break

<sup>16</sup> This price was included because there has undoubtedly been an increase in the price of seed between that reported by Matthiessen and Smith (1979) and 1982, the year for which the algae cost data apply.

cost down by period of the animal's life. Estimates of cost by period are essential to determine the optimal time (*viz.* size) at which to sell the seed, inasmuch as the optimal selling age is at the point where marginal cost is equal to the price that could potentially be received.

The purpose of this research is to contribute to the existing body of knowledge about the hatchery industry by examining the factors that influence production costs. The research is partly based on observations of a working hatchery located in the northeast and on current literature, with particular attention paid to the variations that exist among hatcheries. The data collected were used to develop the framework for a computer program that could estimate variable production costs. The program routine was intended to enable the individual hatcherist to determine the optimal size at which to sell the bivalve seed.

Creation of this program required the development of an algorithm that simulates the production process. The model sums costs at different stages of animal growth, which are further broken down into smaller units of time. In this way a cost can be computed at any given point in the animals' lives. The cost incurred during any given period of time is a function of the quantity of inputs used and their prices. Thus, the major component of the model was designed to determine the quantity of inputs used and the cost incurred during any given period.

There was sufficient information available from the literature and hatchery observations to simulate the overall production system. Unfortunately, records concerning the expenditures did not distinguish between hatchery and open-water field operations nor did they break costs down by stage. Consequently, data were not available to allow for all aspects of the model to be completed in detail. Data on the batch algae production process used by the hatchery were, however, more readily isolated from the rest of the hatchery because of the physical separation of the two systems in the hatchery. Thus, the focus of the empirical analysis was on the estimation of costs of algae production and is

justified since feeding costs are recognized as a large proportion of total costs.

The FORTRAN computer program that is based on the model focuses on the cost of producing algae feed, but as more details become available about other hatchery production processes it would be a simple matter to incorporate them into the existing FORTRAN code. The computer program calculates the cost, or internal price of algae to the hatchery, and then combines this information with survival and growth rates and feeding efficiencies to determine total (and per unit) feeding costs to raise a batch of bivalves to various sizes. The algorithms are used to examine the sensitivity of the algae production parameters and to the parameters of the bivalve growth, survival and feeding relationships.

The model is applied to raising Atlantic Oyster seed, using data on algae production from the hatchery and data on survival rates, feeding efficiency and growth rates available from the literature. Twelve initial simulations were made. The output from each contained a summary of input prices and requirements and costs for bivalves at different periods throughout the production process, along with the age, size and number of bivalves and the total and average costs per bivalve.

The 12 simulations reflect a range of assumptions about survival rates, growth rates and algae feeding rates. In the base run, feed requirements are related directly to animal size. The feed cost per surviving bivalve remains less than one cent up to 100 days but rises rapidly thereafter. In a second run, based on the same data to relate feed directly to age, feed cost remains lower for the first 100 days, but then rises to 1.3 times the base run.

As might be expected, the feed costs per bivalve are extremely sensitive to the parameters of the feeding relationships and to the age-size relationship. This is not particularly encouraging because there seem to be few sources of data in the literature from which to estimate these relationships. The one encouraging thing is that feed costs are

less sensitive to these parameters for very young bivalves than for older ones, and it is the young bivalve that is of most interest to a hatchery. Another important result is that for wide differences in the numbers of animals surviving, the feed cost per surviving animal (including the accumulated cost of feeding those that did not survive) varied by no more than plus or minus 3%. Having good estimates of feed costs per animal in the face of considerable uncertainty about survival rates may well facilitate planning when to sell the bivalve seed or transfer them for growout.

To explore the implications of these feed cost simulations for the marketing of bivalve seed, some rough estimates of total costs of production per bivalve were developed on the basis of others' estimates of the proportion that feed costs are of total costs. Total costs were estimated for several simulations assuming feed costs were 1) one-third, 2) nearly 60%, 3) 85% of total costs, and 4) a linear function of total costs starting at 15% and rising to 85% by the end of the production period. While the time period at which price just covered costs differed significantly over the scenarios, within a given scenario the "break-even" marketing periods varied only by a week or two in spite of diverse assumptions about the fraction feed costs are of total costs.

In conclusion, this research demonstrates that it is possible and practical to estimate algae and feeding costs for a seed producing bivalve hatchery using a computer program to simulate the production system. The most optimistic assumptions suggest that bivalve seed can be produced and marketed profitably. These conclusions must be qualified by the fact that sufficient data were not available from the hatchery under study to estimate all components of total cost. Collection of these types of data would help in further testing of the algorithm as well as contributing to an overall understanding of hatchery costs at each stage of production. Equally important from a research perspective is the need for better data on hatchery survival rates and the size-age and feeding rates. What data do exist in the literature on which to estimate these rates lead to quite different implications for feed costs, and none of the

existing data sets contains more than a few observations. Thus, in the absence of more extensive research data on which to estimate these important physical relationships, anyone attempting to use this kind of software for management decisions is well advised to use information that is specific to a particular hatchery.

### References

- Agnello, R. J. and L. P. Donnelley. "Externalities and Property Rights in the Fisheries". Land Economics, 52 (1976):518-529.
- Baab, J. S., G. L. Hamm, K. C. Hanes, A. Chu and O. A. Roels. "Shellfish Mariculture in an Artificial Upwelling System." Proceedings of the National Shellfisheries Association, 63 (1973):63-68.
- Bardach, J. E., J. Ryther and W. McLarney. Aquaculture - The Farming and Husbandry of Freshwater and Marine Organisms. New York: John Wiley and Sons Inc., 1972.
- Bolton, E. (ed.) Intensive Marine Bivalve Cultivation in a Controlled Recirculating Seawater Prototype System. DEL-SG-07-82, University of Delaware, 1982.
- Breese, W. P. and R. E. Malouf. "Hatchery Manual for the Pacific Oyster". Oregon State University Sea Grant College Program, Agriculture Exper. Station, Special Report Number 443, December, 1975.
- Castagna, M. and J. N. Kraeuter. "Mercenaria Culture Using Stone Aggregate for Predator Protection". Proceedings of the National Shellfisheries Association, 67 (1977):1-6.
- Claus, G. and C. Adler. "Optimal Conditions for Oysters Grown in Closed Environments". Proceedings of the Marine Technical Society, 1 (1970):671-679.
- Donohue, M. J., D. A. Storey and P. G. Allen. "Technical and Economic Feasibility of Alternative Aquacultural Production Systems for American Oysters and Other Bivalve Mollusks in New England". Massachusetts Agricultural Experiment Station Research Bulletin, Number 668, June 1981.
- Dressel, D. M. and D. S. Fitzgibbon. "The U.S. Molluscan Shellfish Industry". United States Dept. of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Washington, D.C., pp. 255-261, in "Drugs and Food from the Sea; Myth or Reality?", P.N. Kaul, and C.J. Sindermann (eds.), University of Oklahoma, 1978.
- Eldridge, P., A. Eversole and J. Whetstone. "Comparative Survival and Growth of Hard Clams Mercenaria Mercenaria, Planted in Trays Subtidally and Intertidally at Varying Densities in a South Carolina Estuary". Proceedings of the National Shellfisheries Association, 69 (1979):30-39.
- Emmel, T. G. Population Biology. New York: Harper and Row, Publishers, 1976.
- Epifanio, C. E. "Culture of Bivalve Mollusks in Recirculating Systems: Nutritional Requirements". Proceedings of the First International Conference on Aquaculture Nutrition, , pp. 173-194, October, 1975.
- Epifanio, C. E., C. M. Logan and C. Turk. "Culture of Six Species of Bivalves in a Recirculating Seawater System". Proceedings of the 10th European Symposium of Marine Biology, 1976.

- Freedman, M. and T. Morris. "A Dwindling Harvest Industry". Sunday Newsday, Long Island, New York, Suffolk Edition, p.3, March 6, 1983.
- Galtsoff, P. S. "The American Oyster Crassostrea Virginica Gmelin". Fishery Bulletin of the Fish and Wildlife Service, 64(1964):397-301.
- Gates, J. M., G. C. Matthiessen and C. A. Griscom. "Aquaculture in New England". Sea Grant University of Rhode Island, Marine Technical Report Series Number 18, Kinston, 1974.
- Goldstein, M. "Plan for Clams - Without Rocking the Boat". Sunday Newsday, Long Island Newspaper, Suffolk Edition, p.21, March 7, 1983
- Hanson, J.A. (ed.) "Open Sea Mariculture". Pennsylvania: Hutchinson and Ross, 1974.
- Hardin, G. "The Tragedy of the Commons". Science, 162 (1968):1243-1247.
- Henderson, S. P. "Shellfish Hatcheries, an Industry View" in Drug and Food from the Sea: Myth or Reality? P.N. Kaul and C.J. Sindermann, eds., University of Oklahoma, 1978.
- Im, K. H., R. S. Johnston and R. D. Langmo. "The Economics of Hatchery Production of Pacific Oyster Seed: A Research Progress Report". Proceedings of the National Shellfisheries Association, 66 (1976):81-94.
- Im, K. H. and R. D. Langmo. "Economic Analysis of Producing Pacific Oyster Seed in Hatcheries". Proceedings of the National Shellfisheries Association, 67 (1977):17-28.
- Kochiss, J. M. "Oystering from New York to Boston". Connecticut: Wesleyan University Press, 1974.
- Korringa, P. "Farming the Cupped Oyster of the Genus Crassostrea". New York: Elsevier Scientific Publishing Co., 1976.
- Loosanoff, V. L. and H. C. Davis. "Rearing of Bivalve Mollusks". Advances in Marine Biology 1(1963a):1-136.
- Loosanoff, V. L. and H. C. Davis. "Shellfish Hatcheries and Their Future". United States Department of the Interior, Fish and Wildlife Service, Washington, D.C., Commercial Fisheries Review, 25, 1963b.
- Mansfield, E. "Microeconomics - Theory and Application". New York: WW Norton and Co., 1979.
- Manzi, J. J., V. G. Burrell Jr. and H. Q. Clawson. "Commercialization of the Hard Clam (*Mercenaria mercenaria*), Mariculture in South Carolina". Preliminary Report, in J. W. Avault Jr. (ed.), Proceedings of the Twelfth Annual Meeting World Mariculture Society. Louisiana State University, 1982a.
- Manzi, J. J., V. G. Burrell Jr., H. Q. Clawson, F. S. Stevens and M. B. Maddox. "Hard Clam, *Mercenaria mercenaria* (Linne), Mariculture in South Carolina", draft for review only, Oct., 1982b.
- Matthiessen, G. C. "Review of Oyster Culture and the Oyster Industry in North America". Number 2528, Woods Hole Oceanographic Institute, 1970.
- Matthiessen, G. C. "The Oyster Industry of Massachusetts and the Introduction of Exotic Species" in R. Mann (ed.), Exotic Species and Mariculture, pp. 213-224, MIT Press, Cambridge, 1979.
- Matthiessen, G. C. and L. J. Smith. "Analysis of Methods for the Culture of *Crassostrea virginica* in New England". Proceedings of the World Mariculture Society, 10 (1979):609-623.
- Matthiessen, G. C. and R. C. Toner. "Possible Methods of Improving the Shellfish Industry of Martha's Vineyard, Duke's County, Massachusetts". Marine Research Foundation, Inc., Edgartown, Massachusetts, 1966.
- McHugh, J. and A. Williams. "Historical Statistics of the Fisheries of the New York Bight", New York Sea Grant Institute, Albany, New York, 1976.
- Miller, R. "A View Through the Porthole by a Commercial Fisherman" in O. Terry (ed.) Proceedings of the Symposium on Aquaculture in New York State, 1977.
- Myers, J. M. Production and Feeding Costs of a Bivalve Seed Hatchery. Unpublished M.S. thesis, June 1985.
- Pillay, T. V. R. "The State of Aquaculture 1975". FAO Technical Conference on Aquaculture, May 26 - June 2, 1976, Kyoto, Japan, pp. 1-13.
- Regional Marine Resources Council, Committee of the Nassau-Suffolk Regional Planning Board. "Guidelines for Management of Long Island Hard Clam Resources", pp. 2-13, 1974.
- Rhodes, E. W. and W. S. Landers. "Growth of Oyster Larvae *Crassostrea virginica* of Various Sizes in Different Concentrations of the chrysophyte, *Isochrysis galbana*". Proceedings of the National Shellfisheries Association, 63(1973):53-59.
- Rhodes, T. "Comments of a Bayman" in O. Terry and D. Chase (eds.) Proceedings of the Symposium on Mariculture in New York State, Southampton College, New York, 1974.
- Ritchie, T. P. "A Comprehensive Review of the Commercial Clam Industries in the United States". Delaware Sea Grant Program Publication, #DEL-SG-26-76, United States Government Printing Office, Washington, D.C., 1977.
- Smith, B. STOP Project Director, interviews, 1982-83.
- Terry, O. W. "Aquaculture," MESA New York Bight Atlas Monograph Number 17, New York Sea Grant Institute, Albany, New York, June 1977.
- Thomas, G. B., Jr. Calculus and Analytic Geometry. 3rd ed. Reading, Mass.: Addison-Wesley Publishing Co., Inc. 1960.
- Ukeles, R. "View on Bivalve Larvae Nutrition". Proceedings of the First International Conference on Aquaculture Nutrition, pp. 127-162, 1975.
- United States Department of Commerce, NOAA, NMFS. "New York Landings, Annual Summary 1973". Current Fisheries Statistics Number 6412, p.2, Washington, D.C., 1974.
- United States Department of Commerce, NOAA, NMFS. "Fishery Statistics of the United States". Washington, D.C., 1979.

**APPENDIX A**



C QUANT3 COMPUTES QUANT OF INPUT 3 USED IN J  
 C QUANT4 COMPUTES QUANT OF INPUT 4 USED IN J  
 C QUANT5 COMPUTES QUANT OF INPUT 5 USED IN J  
 C QUANT6 COMPUTES QUANT OF INPUT 6 USED IN J  
 C COSTJ SUMS AND COMPUTES COST OF INPUTS IN PERIOD, J  
 C WROUT PRINTS COSTS AND QUANTITIES  
 C WOUTN1 PRINTS ADDITIONAL INFO GENERATED ABOUT INPUT 1  
 C WOUTN2 PRINTS ADDITIONAL INFO GENERATED ABOUT INPUT 2  
 C WOUTN3 PRINTS ADDITIONAL INFO GENERATED ABOUT INPUT 3  
 C WOUTN4 PRINTS ADDITIONAL INFO GENERATED ABOUT INPUT 4  
 C WOUTN5 PRINTS ADDITIONAL INFO GENERATED ABOUT INPUT 5  
 C WOUTN6 PRINTS ADDITIONAL INFO GENERATED ABOUT INPUT 6  
 C AGENO FUNCTION RELATING AGE TO SURVIVAL  
 C AGESZ FUNCTION RELATING AGE TO SIZE  
 C BLK1 BLOCK DATA FOR GENERAL PROGRAM  
 C BLK2 BLOCK DATA FOR ORGANISM DEPENDENT VARIABLES  
 C (NOTE: OPTIONAL SUBROUTINES LISTED UNDER APPROPRIATE SUBROUTINES)

C NAME	C TYPE	C COMMON DIMENS	C DESCRIPTION OF COMMON BLOCK VARIABLES
C AGEUPJ R	C MAIN	MAXJI	AGE OF ORGANISM AT END OF PERIOD
C ANYNOR I	C MAIN	NOOFN	TO DETERMINE IF ANY N ARE TO BE COMPUTED
C BEGINI I	C MAIN	NOOFN	STAGE TO BEGIN COST CALCULATIONS.
C BEGINM I	C MAIN	NOOFN	OPTION 1, BEGIN AT STAGE(BEGINI+1)
C BEGINR I	C MAIN	NOOFN	FIRST MONTH OF PRODUCTION
C CSOOFJ R	C MAIN	MAXJI	NO. OF THE 1ST PERIOD OF EACH STAGE
C CSOUPJ R	C MAIN	MAXJI	COST/ORGANISM - AVERAGE IN STAGE
C CSNOFJ R	C MAIN	MAXJI	COST/ORGANISM UP TO END OF PERIOD, J
C CSNUPJ R	C MAIN	MAXJI	COST OF EACH INPUT, N, DURING PERIOD, J
C CSOFJ R	C MAIN	MAXJI	COST OF ALL INPUTS, N, DURING PERIOD, J
C CSUPJ R	C MAIN	MAXJI	OPTION 1: COST UP TO 1ST STAGE TO CALCULATE
C DATE C**50	C MAIN	MAXJI	COST OF ALL INPUTS, N, UP TO END OF PERIOD, J
C DAYUPJ R	C MAIN	MAXJI	CURRENT DATE
C DAYOPI R	C MAIN	NOOFI	TOTAL ELAPSED DAYS UP TO END OF PERIOD, J
C DAYSI R	C MAIN	NOOFI	TOTAL DAYS IN EACH STAGE, I
C DAYSJ R	C MAIN	NOOFI	LAST PERIOD CALCULATED COST FOR
C ENDM I	C MAIN	MAXJI	LAST MONTH OF THE PRODUCTION SEASON
C TOFJ I	C MAIN	MAXJI	STAGE THAT EACH PERIOD IS IN
C JPRI I	C MAIN	NOOFI	NUMBER OF PERIODS, J, IN STAGE, I
C MAXI I	C PARI	MAXJI	MAXIMUM NO. OF STAGES, I, POSSIBLE: PAR=6
C MAXJ I	C PARI	MAXJI	MAX NO. OF PERIODS, J, IN STAGES: PAR=10
C MAXJT I	C PARI	MAXJI	MAX NO. PERIODS: PAR = MAXJ X MAXN
C MAXM I	C PARI	MAXJI	MAX NO. OF MONTHS: PAR=12
C MAXN I	C PARI	MAXJI	MAX NO. OF INPUTS, N: PAR=6
C MAXT I	C PARI	MAXJI	MAX NO. OF TYPES, T, IN INPUT N: PAR=6
C MOFJ I	C MAIN	MAXJI	MONTH THAT J OCCURS IN
C M1 I	C MAIN	MAXJI	MONTH TO START CALCULATIONS ON
C M2 I	C MAIN	MAXJI	MONTH TO END CALCULATIONS ON
C NAMEF1 C**10	C MAIN	NAMEF1	NAME OF INPUT FILE USED
C NAMEM C**10	C BLK1	MAIN	NAME OF EACH MONTH, M
C NAMES C**25	C MAIN	NAMEF1	NAME OF SYSTEM USED
C NOOFO I	C PARI	NOOFI	NO. OF ORGANISM TYPES, O: PAR=2
C NOOFI I	C PARI	NOOFI	NO. OF STAGES, I: PAR=5
C NOOFN I	C PARI	NOOFN	NO. OF INPUTS, N: PAR=6
C NOOFT I	C BLK1	NOOFN	NO. OF TYPE, T, OF EACH INPUT, N
C NQBYM C**1	C MAIN	NOOFN	DOES QUANTITY OF INPUT N VARY BY MONTH?
C N\$BYM C**1	C MAIN	NOOFN	DOES PRICE OF INPUT, N, VARY BY MONTH?

0187 NO. OF ORGANISMS AT BEGIN  
 0188 AVG. NO. OF ORGANISMS DURING EACH PERIOD, J  
 0189 NO. OF ORGANISMS AT END OF EACH PERIOD, J  
 0190 TYPE OF ORGANISM TO DO CALCULATIONS FOR  
 0191 CAN PROGRAM COMPUTE PRICE OF INPUT, N?  
 0192 0=YES, 1=NO  
 0193 DOES THE USER WANT TO ENTER N IN I?  
 0194 CAN PROGRAM COMPUTE INPUT, N, IN STAGE, I?  
 0195 0=YES, 1=NO  
 0196 OPTIONS 1-3 THAT IS CHOSEN TO COMPUTE Q  
 0197 CAN PROGRAM DO EACH OPTION FOR COMPUTING?  
 0198 0=YES, 1=NO  
 0199 CAN USER CHANGE NO. OF TYPES, T OF INPUT, N?  
 0200 0=YES, 1=NO  
 0201 IS IT POSSIBLE TO WRITE ADDITIONAL INFO N?  
 0202 IF PAST MONTHS OF PROD., SET TO 1.  
 0203 PRICE OF TYPE, T, OF INPUT, N, IN MONTH, M.  
 0204 MAXT  
 0205 MAXT  
 0206 MAXT  
 0207 MAXT  
 0208 MAXT  
 0209 UNIT NUMBER FOR READ FILE  
 0210 UNIT NUMBER FOR READ INTERACTIVE  
 0211 AVG SIZE OF ORGANISM DURING PERIOD  
 0212 SIZE OF ORGANISM AT END OF PERIOD  
 0213 NO. OF STAGES TO DO COMPUTATIONS FOR (OP=1) 0213  
 0214 QUANTITY OF TYPE, T, OF INPUT, N, USED IN  
 0215 MONTH, M, FOR STAGE, I.  
 0216 MAXT  
 0217 MAXT  
 0218 MAXT  
 0219 MAXT  
 0220 MAXT  
 0221 MAXT  
 0222 MAXT  
 0223 MAXT  
 0224 MAXT  
 0225 MAXT  
 0226 MAXT  
 0227 MAXT  
 0228 MAXT  
 0229 MAXT  
 0230 MAXT  
 0231 MAXT  
 0232 MAXT  
 0233 MAXT  
 0234 MAXT  
 0235 MAXT  
 0236 MAXT  
 0237 MAXT  
 0238 MAXT  
 0239 MAXT  
 0240 MAXT  
 0241 MAXT  
 0242 MAXT  
 0243 MAXT  
 0244 MAXT  
 0245 MAXT  
 0246 MAXT  
 0247 MAXT  
 0248 MAXT  
 0249 MAXT  
 0250 MAXT

```

0315      C NAMENU C*5 CBLKO NOOFN UNIT NAME OF INPUT, N          0251
0316      C NAMET C*10 CBLKO NOOFN NAME OF EACH TYPE T, OF INPUT, N 0252
0317      C NOO1  R CORG MAXT NO. ORGANISM IN I=1, CONDITIONING STAGE 0253
0318      C N5AVAL R CORG MAXT AMOUNT OF EACH TYPE OF INPUT N=5 PRODUCED 0254
0319      C N5ABYM C CORG MAXT DOES THE AMOUNT OF N=5 PROD. VARY BY MONTH? 0255
0320      C N5REQ  R CORG MAXT FRACTION OF EACH TYPE OF N=5 USED / STAGE 0256
0321      C QEQQ5  I CORG MAXT READ(RI,*), UNIN(N) 0257
0322      C QIAOFJ R CORG MAXT UNIN(N) = UNIN(N) 0258
0323      C          ENDIF 0259
0324      C          CONTINUE 0260
0325      C          WF = WI 0261
0326      C          RF = RI 0262
0327      C          RERUN = .TRUE. 0263
0328      C          CALL REREAD(WI,RI,RERUN) 0264
0329      C          GOTO 100 0265
0330      C          ----- 0266
0331      C          ----- 0267
0332      C          ----- 0268
0333      C          ----- 0269
0334      C          ----- 0270
0335      C          ----- 0271
0336      C          ----- 0272
0337      C          ----- 0273
0338      C          ----- 0274
0339      C          ----- 0275
0340      C          ----- 0276
0341      C          ----- 0277
0342      C          ----- 0278
0343      C          ----- 0279
0344      C          ----- 0280
0345      C          ----- 0281
0346      C          ----- 0282
0347      C          ----- 0283
0348      C          ----- 0284
0349      C          ----- 0285
0350      C          ----- 0286
0351      C          ----- 0287
0352      C          ----- 0288
0353      C          ----- 0289
0354      C          ----- 0290
0355      C          ----- 0291
0356      C          ----- 0292
0357      C          ----- 0293
0358      C          ----- 0294
0359      C          ----- 0295
0360      C          ----- 0296
0361      C          ----- 0297
0362      C          ----- 0298
0363      C          ----- 0299
0364      C          ----- 0300
0365      C          ----- 0301
0366      C          ----- 0302
0367      C          ----- 0303
0368      C          ----- 0304
0369      C          ----- 0305
0370      C          ----- 0306
0371      C          ----- 0307
0372      C          ----- 0308
0373      C          ----- 0309
0374      C          ----- 0310
0375      C          ----- 0311
0376      C          ----- 0312
0377      C          ----- 0313
0378      C          ----- 0314

C 1. PARAMETERS
C INCLUDE ((IPAR1))
C 2. ARGUMENTS
INTEGER RED, WRT
LOGICAL RERUN
C 3. COMMON BLOCKS
INCLUDE ((ICBLK1))
INCLUDE ((ICBLKO))
INCLUDE ((ICUNIT))
INCLUDE ((ICMINT))
INCLUDE ((ICUNIT))
C 4. LOCAL VARIABLES
CHARACTER *1 QUEST, QUEST2
INTEGER UNIT, UNITN
DIMENSION UNITN(NOOFN)

C NAME DESCRIPTION OF LOCAL VARIABLES AND ARGUMENTS
C QUEST QUESTION TO CONTINUE WITH PROGRAM DESCRIPTION
C QUEST2 QUESTION TO RERUN PROGRAM WITH MINOR DATA CHANGES
C RED ARGUE. FOR UNIT NUMBERS READ
C RERUN ARGUE. TO RUN AGAIN WITH A FEW CHANGES
C WRT ARGUE. FOR UNIT NUMBERS WRITE
C UNIT UNIT NO. REQUESTED FOR OUTPUT FILES
C UNITN UNIT NO. REQUESTED FOR SPECIFIC PROD. INPUT OUTPUT FILES
C          ----- 0284
C          ----- 0285
C          ----- 0286
C          ----- 0287
C          ----- 0288
C          ----- 0289
C          ----- 0290
C          ----- 0291
C          ----- 0292
C          ----- 0293
C          ----- 0294
C          ----- 0295
C          ----- 0296
C          ----- 0297
C          ----- 0298
C          ----- 0299
C          ----- 0300
C          ----- 0301
C          ----- 0302
C          ----- 0303
C          ----- 0304
C          ----- 0305
C          ----- 0306
C          ----- 0307
C          ----- 0308
C          ----- 0309
C          ----- 0310
C          ----- 0311
C          ----- 0312
C          ----- 0313
C          ----- 0314

RERUN = .FALSE.

C C OPTION TO RERUN WITH A FEW CHANGES TO DATA
WRITE(WI,30000)
NAME
READ(RI,1000) NAME
READ(RI,1000) QUEST2
IF (QUEST .EQ. 'Y') CALL INFORM

C C READ IN DATA AND CALL BODY TO COMPUTE QUANT. AND COSTS
CALL READIN (RERUN)
C          ----- 0305
C          ----- 0306
C          ----- 0307
C          ----- 0308
C          ----- 0309
C          ----- 0310
C          ----- 0311
C          ----- 0312
C          ----- 0313
C          ----- 0314

C C WRITE OUT INFORMATION CONCERNING PROGRAM USE
SUBROUTINE INFORM
C          ----- 0301
C          ----- 0302
C          ----- 0303
C          ----- 0304
C          ----- 0305
C          ----- 0306
C          ----- 0307
C          ----- 0308
C          ----- 0309
C          ----- 0310
C          ----- 0311
C          ----- 0312
C          ----- 0313
C          ----- 0314

C 1. PARAMETERS
INCLUDE ((IPAR1))
C 2. COMMON BLOCKS
INCLUDE ((ICBLK1))
INCLUDE ((ICUNIT))
INCLUDE ((ICMAIN))
INCLUDE ((ICLINK))
C 3. LOCAL VARIABLES
DATE
READ(RI,*)
IF (WRITE(WI,40000) .EQ. 'Y') THEN
  WRITE(WI,40000)
  READ(RI,*)
ENDIF

```

```

INTEGER I, N, O
CHARACTER*1 QUEST

C NAME      DESCRIPTION OF LOCAL VARIABLES
C-----+
C I         LOOP COUNTER FOR STAGES
C N         LOOP COUNTER FOR INPUTS
C O         LOOP COUNTER FOR ORGANISM
C QUEST     TO CONTINUE
C

2   12, ' TYPES' /' THE USER CAN ',          0443
3   'CHOOSE THE NUMBER OF TYPES OF THE FOLLOWING INPUTS: ' 0444
0380 0381 0382 32000 FORMAT '(2X,A10)
0383 0384 0385 40000 FORMAT(' // THE PROGRAM IS CAPABLE OF COMPUTING THE FOLLOWING ', 0445
0386 0387 0388 41000 FORMAT(' ,1X,A,'IN THE ',A10,' STAGE')
0389 0390 0391 50000 FORMAT(' // THE PROGRAM IS CAPABLE OF COMPUTING THE FOLLOWING ', 0446
0392 0393 0394 40000 FORMAT(' // INPUT QUANTITIES: ') 0447
0395 0396 0397 51000 FORMAT(' ,1X,A10)
0398 0399 0400 60000 FORMAT(' // NOTE: ALTHOUGH THE PROGRAM CAN COMPUTE FOR ',12, 0448
0401 0402 0403 0404 0405 0406 70000 FORMAT(' // FOLLOW THESE GUIDELINES TO ENTER REQUIRED DATA: /' 0449
0407 0408 0409 0410 0411 0412 0413 0414 0415 0416 0417 0418 0419 0420 0421 0422 0423 0424 0425 0426 0427 0428 0429 0430 0431 0432 0433 0434 0435 0436 0437 0438 0439 0440 0441 0442 0443 0444 0445 0446 0447 0448 0449 0450 0451 0452 0453 0454 0455 0456 0457 0458 0459 0460 0461 0462 0463 0464 0465 0466 0467 0468 0469 0470 0471 0472 0473 0474 0475 0476 0477 0478 0479 0480 0481 0482 0483 0484 0485 0486 0487 0488 0489 0490 0491 0492 0493 0494 0495 0496 0497 0498 0499 0500 0501 0502 0503 0504 0505 0506

32
CONTINUE
WRITE(WI,31000) MAXT
DO 32 N=1,NOEN
  IF (OPTR(N) .EQ. 0) WRITE (WI,32000) NAMEN(N)
CONTINUE
WRITE(WI,30001) .EQ. 0
READ(RI,1000) QUEST
WRITE(WI,40000)
DO 40 N=1,NOEN
  DO 41 I=1,NOIFI
    IF (OPNR(I,N) .EQ. 0) WRITE(WI,41000) NAMEN(N), NAMEI(I)
41  CONTINUE
40  CONTINUE
WRITE(WI,50000)
DO 50 N=1,NOEN
  IF (OPNSOK(N) .EQ. 0) WRITE(WI,51000) NAMEN(N)
CONTINUE
WRITE(WI,60000) MAXM, NAMEM(1), NAMEM(MAXM)
WRITE(WI,30001) QUEST
READ(RI,1000) QUEST
WRITE(WI,70000)
WRITE(WI,30001) QUEST
READ(RI,1000) QUEST

1000 FORMAT(A1)
30001 FORMAT(' // CONTINUE? ')
10000 FORMAT(' // PROGRAM IS CAPABLE OF CALCULATING COSTS FOR THE ', 0443
2   2   ' UNITS OR PERIODS. /' THE MAXIMUM NUMBER OF PERIODS ', 0444
3   3   ' ALLOWED IN ALL STAGES COMBINED IS ',I2)
20000 FORMAT(' // THE PRODUCTION PROCESS IS BROKEN INTO THE FOLLOWING ', 0445
2   2   ' 2   ' FOLLOWING ',A,'TYPES: /'6(2X,A10))
25000 FORMAT(' // THESE STAGES CAN THEN BE BROKEN DOWN INTO SMALLER ', 0446
2   2   ' UNITS OR PERIODS. /' THE MAXIMUM NUMBER OF PERIODS ', 0447
3   3   ' ALLOWED IN ALL STAGES COMBINED IS ',I2)
30000 FORMAT(' // THE QUANTITY OF EACH OF THE ',I2,' INPUTS USED IN ', 0448
2   2   ' EVERY STAGE IS EITHER ',/,' COMPUTED OR ENTERED. ', 0449
3   3   ' THESE INPUTS ARE: /',6(1X,A10,' PER ',A5,' /))
31000 FORMAT(' // SOME OF THESE INPUTS MAY BE SEPARATED INTO UP TO ', 0450

C STRUCTURE: OPTION TO INPUT BY FILE OR INTERACTIVELY
C             DISPLAY AND CHOOSE OPTIONS FOR DATA INPUT
C             READ IN DATA BY CALLING:
C             READS
C             READN, N=1, NUMBER OF INPUTS
C             2ND ENTRY POINT, REREAD, TO RERUN WITH ONLY MINOR CHANGE
C             OPTION TO CHECK, CORRECT, AND PRINT DATA
C             READ
C             C 1. PARAMETERS
C             INCLUDE '(IPAR1)'
C             C 2. ARGUMENTS
C               INTEGER RED, RERED, WRT, REWRT, HREAD
C               LOGICAL RERUN, RUN2
C             C 3. COMMON BLOCKS
C               INCLUDE '(ICMAIN)'
C               INCLUDE '(ICBLKL)'
```

```

INCLUDE  (ICBLKO)
INCLUDE  (ICUNITS)
C 4. LOCAL VARIABLES
    INTEGER I, INFO, K, K2, KGOTO, KLAST, N, NOOKF, OPEN,
    2      R, UNIT, W, Z
    LOGICAL CHECK, RIPROB, RUN1
    CHARACTER * 1 QFILE, QUEST, QUEST2, QUEST3
    PARAMETER (NOOKF = 8)
    DIMENSION INFO(NOOKF)

C NAME      DESCRIPTION OF LOCAL VARIABLES AND ARGUMENTS
C ----- -----
C CHECK     USED TO DETERMINE IF LOOP HAS BEEN ENTERED
C READ      ARGUE: IOSTAT TO SHOW PROBLEM IN READING FILE (2-WAY COMM.)
C I         LOOP COUNTER FOR STAGE, I
C INFO      INFORMATION LINE NUMBER
C K         COUNTER FOR LINE NUMBER
C K2        COUNTER FOR SUBROUTINE LINE NUMBERS
C KCOTO    NO. OF THE INFO LINE TO REINPUT DATA
C KLAST    THE LAST INFO LINE, K, INPUT
C N         LOOP COUNTER FOR INPUTS
C NOOKF   NUMBER OF INPUT LINES
C OPEN     COUNT OF N. OF N TO ENTER FOR EACH I
C QFILE    WANT TO USE A DATA FILE?
C QUEST    GENERAL QUESTION FOR USER (IS WRITTEN OVER)
C QUEST2   OPTION TO STOP, REDO, OR CONTINUE WITH DATA ENTRY
C QUEST3   OPTION TO SEE OPTION INFORMATION
C R         UNIT NUMBER FOR READ
C RED      ARGUE: UNIT NUMBERS FOR READ
C RERUN    ARGU: UNIT NO. FOR 2ND ENTRY POINT - ALWAYS INTERACTIVE
C RERUN    ARGU: TO ALLOW INFO PRINTING IF NOT RUN1, FROM 2ND ENTRY PNT037
C RERUN    ARGU: UNIT NO FOR 2ND ENTRY POINT - ALWAYS INTERACTIVE
C RERWT    TO DETERMINE IF FILE PROBLEM HAS BEEN ENCOUNTERED
C RIPROB   USED TO PRINT INFO THE FIRST RUN THROUGH
C RUN1    ARGUE: IF FIRST RUN THROUGH DATA INPUT ALREADY DONE
C UNIT     UNIT NO FOR OUTPUT FILE TO GO TO
C W         UNIT NUMBER FOR WRITE
C WRT     ARGUE: UNIT NUMBERS FOR WRITE
C Z         2ND COUNTER FOR LINE NUMBERS
C ----- -----
C SET ALL LINES TO 0 IN ORDER TO READ IN THE FIRST TIME
    DO 100 K=1,NOOKF
        INFO(K) = 0
100    CONTINUE
        RUN1 = .TRUE.
        RUN2 = .FALSE.

C INITIALIZE FILE VARIABLES
    RIPROB = .FALSE.
    READ = 0

C OPTION TO TERMINATE
    WRITE(WI,20000)
    READ(RI,1000) QUEST
    IF (QUEST .EQ. 'Y') STOP 'COMPLETE SPREAD SHEET'

C READ IN PROGRAM DATA/IDENTIFIER AND OUTPUT FILE UNIT NUMBER
    WRITE(WI,50000)
    READ(RI,*) DATE
    WRITE(WI,260000)
    READ(RI,*) UNIT
    WO = UNIT

C OPTION TO INPUT BY SEQUENTIAL FILE
    WRITE(WI,300000)
    READ(RI,1000) QFILE
    IF (QFILE .EQ. 'Y') THEN
        WRITE(WI,310000)
        READ(RI,*) NAMEF1
        WRITE(WI,320000)
        W = WF
        R = RF
        WRT = WF
        RED = RF
    ELSE
        W = WI
        R = RI
        WRT = WI
        RED = RI
        NAMEF1 = 'NOFILE'
    ENDIF
    C LIST OPTIONS AVAILABLE TO READ DATA IF OPTION TO DISPLAY CHOSEN
    250   WRITE(WI,40000)
    WRITE(WI,40010)
    READ(R,1000) QUEST3
    IF (QUEST3 .EQ. 'Y') THEN
        IF (OPNOK(1) .EQ. 0) THEN
            WRITE(WI,40100)
            WRITE(WI,40150)
        ENDIF
        IF (OPNOK(2) .EQ. 0) THEN
            WRITE(WI,40200)
            WRITE(WI,40150)
        ENDIF
        IF (OPNOK(3) .EQ. 0) THEN
            WRITE(WI,40500)
        ENDIF
        READ(R,1000,ERR=600,IOSTAT=HREAD) QUEST
        C LIST INPUTS CALCULATABLE IN EACH STAGE
        0551   WRITE(WI,41000)
        DO 300 N=1,NOOFN
            WRITE(WI,41100) N, NAMEN(N)
            CHECK = .TRUE.
            DO 310 I=1,NOIFI
                IF (OPNOK(I,N) .EQ. 0) THEN
                    WRITE(WI,41200) NAMEI(I)
                ELSE
                    CHECK = .FALSE.
                ENDIF
                CONTINUE
                IF (CHECK) WRITE(WI,41300)
            CONTINUE
            WRITE(WI,40500)
        READ(R,1000,ERR=600,IOSTAT=HREAD) QUEST
        C USER CHOOSES AN OPTION
        WRITE(WI,42000)
        READ(R,*,ERR=600,IOSTAT=HREAD) OPTION
        0569   0570
    ENDIF
    C USER CHOOSES AN OPTION
    WRITE(WI,42000)
    READ(R,*,ERR=600,IOSTAT=HREAD) OPTION
    0570

```

```

0699      IF (QUEST .EQ. 'Y') CALL ROUT(W0)
0700
0701      C OFFER OPTION TO TERMINATE
0702          WRITE(WL,45200)
0703          READ(RI,1000) QUEST
0704          IF (QUEST .EQ. 'Y') STOP 'PROGRAM STOPPED IN OPTION SECTION'
0705
0706          C DISPLAY NECESSARY SUBROUTINES AND THEIR NUMBERS
0707          400 IF (RUN1) THEN
0708              WRITE(W,50000)
0709              WRITE(W,50010)
0710              K2 = 1
0711              WRITE(W,50020) K2
0712              K2 = K2+1
0713              WRITE(W,50030) K2, NAME
0714              DO 410 N=1,NOOFN
0715                  K2 = K2+1
0716                  IF (ANYOK(N) .LT. TOTALI) WRITE(W,50040) K2, NAMEN(N)
0717
0718          CONTINUE
0719          HREAD = 0
0720          RIREB = .FALSE.
0721
0722          C READ IN DATA NECESSARY TO RUN GENERAL PROGRAM
0723          C CONTROL TRANSFERRED HERE IF INPUT CORRECTIONS ARE TO BE MADE
0724          420 CONTINUE
0725          K = 1
0726          IF (INFO(K) .EQ. 0) THEN
0727              WRITE(WL,51000) K
0728              CALL READS(WRT,RED,HREAD,RUN2)
0729          IF (HREAD .NE. 0) GOTO 550
0730
0731          ENDIF
0732          K = K + 1
0733          IF (INFO(K) .EQ. 0) THEN
0734              WRITE(WL,51100) K, NAME
0735              CALL READ0(WRT,RED,HREAD,RUN2)
0736          IF (HREAD .NE. 0) GOTO 550
0737
0738          C READ IN DATA PERTAINING TO QUANTITY OF INPUT, N, ONLY IF
0739          C SOME STAGE IS TO BE COMPUTED FOR THAT INPUT
0740          DO 500 N=1,NOOFN
0741              K = K+1
0742              IF (INFO(K) .EQ. 0 .AND. ANYOK(N) .LT. TOTALI) THEN
0743                  WRITE(WL,53000) K, NAMEN(N)
0744                  IF (N .EQ. 1) CALL READ1(WRT,RED,HREAD,RUN2)
0745                  IF (N .EQ. 2) CALL READ2(WRT,RED,HREAD,RUN2)
0746                  IF (N .EQ. 3) CALL READ3(WRT,RED,HREAD,RUN2)
0747                  IF (N .EQ. 4) CALL READ4(WRT,RED,HREAD,RUN2)
0748                  IF (N .EQ. 5) CALL READ5(WRT,RED,HREAD,RUN2)
0749                  IF (N .EQ. 6) CALL READ6(WRT,RED,HREAD,RUN2)
0750                  IF (HREAD .NE. 0) GOTO 550
0751
0752          ENDIF
0753          WRITE(WL,54000)
0754          CONTINUE
0755          KLAST = K
0756
0757          C MESSAGE IF ERROR DETECTED IN READING FILE
0758          IF (QUEST .EQ. 'Y') GOTO 250
0759
0760          C CONTROL TRANSFERRED HERE IF ERROR FOUND IN READING FILE
0761          600 IF (QFILE .EQ. 'Y') THEN
0762              KLAST = K
0763
0764          C OPTION TO PRINT OPTION INFORMATION
0765          WRITE(WL,45300)
0766          READ(RI,1000) QUEST
0767          IF (NAME1 .EQ. 'NOFILE') THEN
0768              WRITE(WI,45100)
0769              WRITE(WI,45200)
0770              READ(RI,1000) QUEST
0771              IF (QUEST .EQ. 'Y') GOTO 250
0772
0773          C OPTION TO CHECK AND REINPUT OPTION INFORMATION
0774          WRITE(WI,45000)
0775          IF (QUEST .EQ. 'Y') CALL ROUT(WI)
0776          IF (NAME1 .EQ. 'NOFILE') THEN
0777              WRITE(WI,45100)
0778              READ(RI,1000) QUEST
0779              IF (QUEST .EQ. 'Y') GOTO 250
0780
0781          CONTINUE
0782          TOTALI = NOOFI - BEGINI + 1
0783
0784          C OPTION TO CHECK AND REINPUT OPTION INFORMATION
0785          WRITE(WI,45000)
0786          IF (QUEST .EQ. 'Y') CALL ROUT(WI)
0787          IF (NAME1 .EQ. 'NOFILE') THEN
0788              WRITE(WI,45100)
0789              WRITE(WI,45200)
0790              READ(RI,1000) QUEST
0791              IF (QUEST .EQ. 'Y') GOTO 250
0792
0793          CONTINUE
0794
0795          C CONTROL TRANSFERRED HERE IF ERROR FOUND IN READING FILE
0796          696 IF (QFILE .EQ. 'Y') THEN
0797              KLAST = K
0798
0799          C MESSAGE IF ERROR DETECTED IN READING FILE
0800          WRITE(WL,45300)
0801          READ(RI,1000) QUEST

```





```

1019   63000 FORMAT(' ',/,/, 'YOU NOW HAVE SEVERAL OPTIONS CONCERNING ALL ', 1020   C DISPLAY OR PRINT ALL DATA ENTERED IN SUBROUTINE READIN
1020   2      ' , THE DATA ENTERED. YOU CAN: /' 1021   C HEADING FOR OUTPUT
1021   3      ' , '2X,'ENTER S TO STOP AFTER AN OPTION TO PRINT THE DATA' / 1022   WRITE(W,10000) NAME, DATE
1022   3      ' , '2X,'ENTER R TO REDO DATA ENTIRELY INTERACTIVELY' ) 1023   IF (NAMEFL .NE. 'NFILE') WRITE(W,20000) NAMEFL
1023   1024   1025   C OPTION INFORMATION
1025   1026   C OPTION INFORMATION
1026   1027   WRITE(W,30000) OPTION
1027   1028   IF (OPTION .EQ. 1) WRITE(W,31000)
1028   1029   IF (OPTION .EQ. 2) WRITE(W,32000)
1029   1030   IF (OPTION .EQ. 3) WRITE(W,33000)
1030   1031   IF (OPTION .EQ. 1) WRITE(W,34000) NAMEI(BEGINI)
1031   1032   IF (OPTION .EQ. 1) WRITE(W,34000) NAMEI(BEGINI)
1032   1033   IF (OPTION .EQ. 1) WRITE(W,34000) NAMEI(BEGINI)
1033   1034   IF (OPTION .EQ. 1 .OR. OPTION .EQ. 2) THEN
1034   1035   WRITE(W,40000) (NAMEI(L), L=1,NOOFI)
1035   1036   DO 100 N=1,NOOFN
1036   1037   WRITE(W,41000) NAMEN(N), (OPNNW(I,N), I=1,NOOFI)
1037   1038   CONTINUE
1038   1039   ENDIF
1039   1040
1040   1041   C THE NUMBER AND NAME OF EACH INPUT SUBROUTINE
1041   1042   K = 1
1042   1043   WRITE(W,45000)
1043   1044   WRITE(W,50000) K
1044   1045   K=K+1
1045   1046   WRITE(W,51000) K, NAME
1046   1047   DO 200 N=1,NOOFN
1047   1048   K = K+1
1048   1049   K = K+1
1049   1050   IF(ANYNOK(N) .LT. TOTALI) WRITE(W,52000) K, NAMEN(N)
1050   1051   CONTINUE
1051
1052   1052
1053   1053
1054   1054   10000 FORMAT('1',A10,' COST SIMULATION MODEL: INPUTS ENTERED' /
1055   1055   2      ',49( '-' )/ '
1055   1056   3      ', 'DATE: ', A50// ', 'OPTION DATA ENTERED' ')
1056
1057   1057   20000 FORMAT(' ', 'NAME OF FILE USED: ',2X,A10)
1058   1058
1059   1059   30000 FORMAT(' ', 'OPTION CHOSEN: ',1X,11)
1060   1060
1061   1061   31000 FORMAT(' ', '15X,'COST UP TO FIRST STAGE OF INTEREST IS ENTERED' /
1062   1062   2      ', '15X,'USER CHOOSES INPUTS TO COMPUTE IN REMAINING STAGES') 1127
1063   1063
1064   1064   1120
1065   1065   32000 FORMAT(' ', '15X,'USER CHOOSES WHICH INPUTS TO COMPUTE IN EACH ' ,
1066   1066   2      ', 'STAGE')
1067   1067
1068   1068   33000 FORMAT(' ', '15X,'ALL INPUTS IN ALL STAGES ARE COMPUTED ' ,
1069   1069   2      ', 'BY THE PROGRAM')
1070   1070
1071   1071   34000 FORMAT(' ', '15X,'THE FIRST STAGE OF INTEREST IS THE' ,1X,A10,
1072   1072   2      ', 'STAGE')
1073   1073
1074   1074   1137
1075   1075   40000 FORMAT(' ', 'IF C, THE PROGRAM COMPUTES THE QUANTITY' /
1075   1075   2      ', 'IF E, THE USERS ENTERS THE QUANTITY' /
1076   1076   3      ', 'IF -, THE QUANTITY IS NEITHER ENTERED ' ,
1077   1077   4      ', 'NOR COMPUTED' /
1078   1078   5      ', 'INPUT',T15,7(A10,1X))
1079   1079
1080   1080   41000 FORMAT(' ', 'A10,T15,7(A1,10X)
1081   1081
1082   1082   45000 FORMAT(' ', ' / ', 'THE NUMBER OF EACH SECTION FOR WHICH DATA ' ,
1082
C NAME DESCRIPTION OF LOCAL VARIABLES AND ARGUMENTS
C -----
C I LOOP COUNTER FOR STAGES
C K INPUT LINE NUMBER
C N LOOP COUNTER FOR INPUTS
C W UNIT NUMBERS FOR WRITE
C WRT ARGUE: UNIT NUMBER FOR WRITE
C -----
C TRANSLATE ARGUMENTS TO LOCAL VARIABLES
W = WRT

```

2 'MUST BE ENTERED.'

```

50000 FORMAT(' ',2X,'SECTION',1X,I2,: GENERAL
      DATA')
51000 FORMAT(' ',2X,'SECTION',1X,I2,: 'A,' DATA')
52000 FORMAT(' ',2X,'SECTION',1X,I2,: '1X,A10,1X,'DATA')

      RETURN
      END

      SUBROUTINE READS(WRT,RED,HREAD,RUN2)
      C
      C ORGNSM:
      C
      C SUBROUTINE DOES NOT VARY
      C VARIABLES DO NOT CHANGE
      C **NOTE: SOME INFO(K) MAY HAVE TO BE FORCED TO 0 DEPENDING ON HOW
      C          ORGANISM DEPENDENT SUBROUTINES ARE MANAGED

      STRUCTURE: READS IN GENERAL DATA TO RUN PROGRAM
      READS IN PRICE OF EACH INPUT OR ELSE CALLS PRICE, N, TO
      COMPUTE PRICE
      OPTION TO CHECK, CORRECT, OR PRINT DATA

      C 1. PARAMETERS
      INCLUDE (IPAR1)
      C 2. ARGUMENTS
      INTEGER HREAD, RED, RED2, WRT, WRT2, WW
      LOGICAL RUN2, RUN3
      C 3. COMMON BLOCKS
      INCLUDE (ICMAIN)
      INCLUDE (ICBLK1)
      INCLUDE (ICBLKO)
      INCLUDE (ICONIT)

      C 4. LOCAL VARIABLES
      INTEGER I, INFO, K, K2, KGOTO, KLAST, M, NOOFF, R, T, W
      LOGICAL CHECK, RUN1
      CHARACTER * 1 QUEST, QUEST2
      PARAMETER (NOOFF = 54)
      DIMENSION INFO(NOOFF)

      C NAME      DESCRIPTION OF LOCAL VARIABLES AND ARGUMENTS
      C
      C CHECK      TO DETERMINE IF A LOOP HAS BEEN ENTERED
      C HREAD      ARGUE: ISOSTAT TO INDICATE PROBLEMS IN READING FILE
      C I           LOOP COUNTER FOR STAGES
      C INFO       INFORMATION LINE NUMBER
      C K           NUMBER OF THE INFO LINE
      C K2          LOOP COUNTER TO FORCE REINPUT
      C KGOTO      LINE NUMBER TO CORRECT
      C KLAST      LAST INFO LINE NUMBER TO BE INPUT
      C M           LOOP COUNTER FOR MONTH
      C N           LOOP COUNTER FOR INPUT
      C NOOFF      NUMBER OF INPUT LINES
      C QUEST      GENERAL VARIABLE FOR USER QUESTION (IS WRITTEN OVER)

      C QUEST2     OPTION TO STOP, REDO, OR CONTINUE DATA ENTRY
      C R           UNIT NUMBER FOR READ
      C RED         ARGUE: UNIT NUMBER TO READ
      C RED2        ARGUE: UNIT NO. TO READ FOR NEXT LEVEL SUBR.
      C RUN1        FIRST RUN THROUGH
      C RUN2        ARGUE: INDICATES IF ISNT FIRST RUN THROUGH
      C RUN3        ARGUE: INDICATES TO NEXT LEVEL SUBROUTINE NOT 1ST RUN THRU
      C T           LOOP COUNTER FOR TYPE OF INPUT, I
      C W           UNIT NUMBER FOR WRITE
      C WRT         ARGUE: FOR UNIT NUMBERS TO WRITE
      C WRT2        ARGUE: UNIT NO. TO WRITE TO NEXT LEVEL SUBR.
      C WW          ARGUE: UNIT NO. TO WRITE AS OUTPUT FILE
      C
      C
      C NOT FIRST RUN, IMMEDIATELY OFFER OPTION TO CHECK AND CHANGE DATA
      IF (RUN2) THEN
      DO 120 K=1,NOOFF
      INFO(K) = 1
      CONTINUE
      120
      1161
      1162
      1163
      1164
      1165
      1166
      1167
      1168
      1169
      1170
      1171
      1172
      1173
      1174
      1175
      1176
      1177
      1178
      1179
      1180
      1181
      1182
      1183
      1184
      1185
      1186
      1187
      1188
      1189
      1190
      1191
      1192
      1193
      1194
      1195
      1196
      1197
      1198
      1199
      1200
      1201
      1202
      1203
      1204
      1205
      1206
      1207
      1208
      1209
      1210
      1211
      1212
      1213
      1214
      1215
      1216
      1217
      1218
      1219
      1220
      1221
      1222
      1223
      1224
      1225
      1226
      1227
      1228
      1229
      1230
      1231
      1232
      1233
      1234
      1235
      1236
      1237
      1238
      1239
      1240
      1241
      1242
      1243
      1244
      1245
      1246
      1247
      1248
      1249
      1250
      1251
      1252
      1253
      1254
      1255
      1256
      1257
      1258
      1259
      1260
      1261
      1262
      1263
      1264
      1265
      1266
      1267
      1268
      1269
      1270
      1271
      1272
      1273
      1274
      49

```

```

READ(R,*,ERR=900, IOSTAT=HREAD) BEGINM
C   FORCE ALL INFO(K) TO 0 IN CASE OF REINPUT
DO 405 K2 = 2,NOOFK
INFO(K2) = 0
CONTINUE
RUN2 = .FALSE.
ENDIF
K = K+1
IF (INFO(K) .EQ. 0) THEN
WRITE(W,22000) K
READ(R,*,ERR=900, IOSTAT=HREAD) ENDM
FORCE ALL REMAINING INFO(K) TO 0 IN CASE OF REINPUT
DO 406 K2 = 3,NOOFK
INFO(K2) = 0
CONTINUE
RUN2 = .FALSE.
ENDIF
C DETERMINE THE LENGTH OF STAGES, I, AND THE NO. OF PERIODS,J, IN EACH I
1275      WRITE(W,51100) K, NAMEN(N)
1276      C
1277      IF (QOPNS(N) .EQ. 'Y') THEN
1278      SET RELEVANT INFO LINES TO 0 TO FORCE REINPUTTING
1279      IF (INFO(K+5*NOOFN) = 0
1280      ELSEIF (QOPNS(N) .EQ. 'N') THEN
1281      CONTINUE
1282      ENDIF
1283      500    CONTINUE
1284      1347
1285      1348
1286      C LOOP THRU EACH INPUT N, TO ENTER INFO IF OPTION TO COMPUTE $ = NO
1287      IF (RUN1) WRITE(W,30000) NOOFN
1288      DO 600 N=1,NOOFN
1289      1352
1290      1353
1291      C NUMBER OF TYPES, T, OF EACH INPUT,N
1292      K = K+1
1293      IF (INFO(K) .EQ. 0 .AND. OPTOK(N) .EQ. 0 .AND. QOPNS(N) .EQ. 'N') THEN
1294      WRITE(W,31000) K, NAMEN(N)
1295      READ(R,* ,ERR=900, IOSTAT=HREAD) NOOFT(N)
1296      C SET RELEVANT INFO LINES TO 0 TO FORCE REINPUT
1297      IF (NOOFT(N) .GT. 1) THEN
1298      INFO(K+1) = 0
1299      INFO(K+3) = 0
1300      C NEXT FORCING DOESNT WORK. MAKE COMMENT FOR NOW
1301      INFO(K+5*NOOFN) = 0
1302      C NAME OF EACH TYPE, T
1303      K = K+1
1304      IF (INFO(K) .EQ. 0 .AND. OPTOK(N) .EQ. 0 .AND. QOPNS(N) .EQ. 'N') THEN
1305      WRITE(W,53000) K, NAMEN(N)
1306      1369
1307      DO 610 T=1,NOOFT(N)
1308      WRITE(W,32000) K, T, NAMEN(N)
1309      READ(R,* ,ERR=900, IOSTAT=HREAD) NAMET(N,T)
1310      610    CONTINUE
1311      C
1312      NAMET(N,1) = 'ONE'
1313      IF (INFO(K) .EQ. 0) THEN
1314      IF (OPTION .EQ. 1) THEN
1315      WRITE(W,24000) K,
DO 420 I=BEGINI,NOOFI
WRITE(W,2100) NAMEI(I)
READ(R,* ,ERR=900, IOSTAT=HREAD) JPERI(I)
CONTINUE
ENDIF
K = K+1
IF (INFO(K) .EQ. 0) THEN
IF (OPTION .EQ. 1) THEN
WRITE(W,25000) K, NAMEI(BEGINI)
READ(R,* ,ERR=900, IOSTAT=HREAD) DAYOP1
ENDIF
K = K+1
IF (INFO(K) .EQ. 0) THEN
IF (OPTION .EQ. 1) THEN
WRITE(W,26000) K, NAMEI(BEGINI)
READ(R,* ,ERR=900, IOSTAT=HREAD) CSOP1
ENDIF
ENDIF
C DAYS AND COST UP TO THE END OF OPTION 1
1316      C DOES THE PRICE VARY BY MONTH?
1317      K = K+1
1318      IF (INFO(K) .EQ. 0 .AND. QOPNS(N) .EQ. 'N') THEN
1319      IF ((ENDM-BIGINM) .GT. 0) THEN
1320      WRITE(W,53000) K, NAMEN(N)
1321      READ(R,* ,ERR=900, IOSTAT=HREAD) NSBYM(N)
1322      C SET RELEVANT INFO LINES TO 0 TO FORCE REINPUT
1323      INFO(K+1) = 0
1324      ELSE
1325      NSBYM(N) = '-'
1326      ENDIF
1327      C PRICE OF EACH TYPE, T, IN ALL PRODUCTION MONTHS
1328      K = K+1
1329      IF (INFO(K) .EQ. 0 .AND. QOPNS(N) .EQ. 'N') THEN
1330      WRITE(W,54000) K, NAMEN(N), NAMET(N,T)
1331      DO 620 T=1,NOOFT(N)
1332      IF (NOOFT(N) .GT. 1) WRITE(W,55000) NAMET(N,T)
1333      IF (NSBYM(N) .EQ. 'Y') THEN
1334      IF (NSBYM(N) .EQ. 'Y') THEN
1335      WRITE(W,51000) K, NAMEN(N)
1336      READ(R,1000,ERR=900, IOSTAT=HREAD) QOPNS(N)
1337      ELSE
1338      QOPNS(N) = 'N'
1339      ENDIF
1340      C
1341      IF THE OPTION EXISTS TO COMPUTE INPUT PRICE, OFFER IT
1342      K = K+1
1343      IF (INFO(K) .EQ. 0) THEN
1344      IF (OPNSOR(N) .EQ. 0) THEN
1345      WRITE(W,51000) K, NAMEN(N)
1346      READ(R,1000,ERR=900, IOSTAT=HREAD) QOPNS(N)
1347      ELSE
1348      READ(R,* ,ERR=900, IOSTAT=HREAD) PRICET(N,T,M)
1349      ENDIF
1350      ENDIF
1351      ENDIF
1352      ENDIF
1353      ENDIF
1354      ENDIF
1355      ENDIF
1356      ENDIF
1357      ENDIF
1358      ENDIF
1359      ENDIF
1360      ENDIF
1361      ENDIF
1362      ENDIF
1363      ENDIF
1364      ENDIF
1365      ENDIF
1366      ENDIF
1367      ENDIF
1368      ENDIF
1369      ENDIF
1370      ENDIF
1371      ENDIF
1372      ENDIF
1373      ENDIF
1374      ENDIF
1375      ENDIF
1376      ENDIF
1377      ENDIF
1378      ENDIF
1379      ENDIF
1380      ENDIF
1381      ENDIF
1382      ENDIF
1383      ENDIF
1384      ENDIF
1385      ENDIF
1386      ENDIF
1387      ENDIF
1388      ENDIF
1389      ENDIF
1390      ENDIF
1391      ENDIF
1392      ENDIF
1393      ENDIF
1394      ENDIF
1395      ENDIF
1396      ENDIF
1397      ENDIF
1398      ENDIF
1399      ENDIF
1400      ENDIF
1401      ENDIF
1402      ENDIF

```



```

C OPTION TO PRINT THE DATA ENTERED          1555
850  WRITE(WI,80000)                         1596
     READ(RL,1000) QUEST                      1597
     IF (QUEST.EQ.'Y') THEN
       CALL ROUTS(WO)
     ENDIF

     IF(QUEST2.EQ.'S') STOP 'PROGRAM STOP IN DATA ENTRY SECTION 1'

1000  FORMAT (A1)

21000 FORMAT(' /' ,I2,'. THE NUMBER OF PERIODS IN EACH STAGE. ',I2,'. THE NUMBER OF DISTINCT TYPES OF A/ 1531
21000 FORMAT(' /' ,I2,'. SEASON BEGINS /' ,T6,'ENTER AN INTEGER: ') 1532
1533   53000 FORMAT(' /' ,I2,'. DOES THE PRICE OF',1X,A,1X,'VARY BY MONTH?') 1598
1534   54000 FORMAT(' /' ,I2,'. PRICE OF',1X,A,1X,'PER',1X,A5) 1599
1535   55000 FORMAT(' /' ,I2,'. OF TYPE',1X,A,':') 1600
1536   56000 FORMAT(' /' ,I2,'. IN',1X,A) 1601
1537   56000 FORMAT(' /' ,I2,'. OF',1X,A,':') 1602
1538   56000 FORMAT(' /' ,I2,'. IN',1X,A) 1603
1539   56000 FORMAT(' /' ,I2,'. DO YOU WANT TO SEE THE GENERAL DATA?') 1604
1540   56000 FORMAT(' /' ,I2,'. DO YOU WANT TO SEE THE GENERAL DATA?') 1605
1541   56000 FORMAT(' /' ,I2,'. DO YOU WANT TO SEE THE GENERAL DATA?') 1606
1542   64000 FORMAT(' /' ,I2,'. ENTER S TO STOP EXECUTION, AFTER AN OPTION TO', 1607
1543   64000 FORMAT(' /' ,I2,'. PRINT GENERAL DATA /', 1608
1544   64000 FORMAT(' /' ,I2,'. ENTER R TO REDO ALL DATA IN THIS SECTION ', 1609
1545   64000 FORMAT(' /' ,I2,'. INTERACTIVELY /', 1610
1546   64000 FORMAT(' /' ,I2,'. ENTER C TO CONTINUE EXECUTION WITH CORRECTIONS ', 1611
1547   64000 FORMAT(' /' ,I2,'. LINE BY LINE /', 1612
1548   64000 FORMAT(' /' ,I2,'. ENTER S, R, OR C') 1613
1549   64000 FORMAT(' /' ,I2,'. ENTER S, R, OR C') 1614
1550   70000 FORMAT(' /' ,I2,'. YOU MAY CORRECT THE GENERAL DATA A LINE AT A TIME /', 1615
1551   70000 FORMAT(' /' ,I2,'. ENTER 0 TO MAKE NO CORRECTIONS OR /', 1616
1552   70000 FORMAT(' /' ,I2,'. 2X, ENTER THE NUMBER OF THE LINE YOU WANT TO CORRECT.') 1617
1553   70000 FORMAT(' /' ,I2,'. 2X, (NOTE: IF YOU CHANGE MONTHS OF PRODUCTION, ALL ', 1618
1554   70000 FORMAT(' /' ,I2,'. DATA MUST BE REINPUT.) /', 1619
1555   70000 FORMAT(' /' ,I2,'. CHANGES TO OTHER DATA ', 1619
1556   70000 FORMAT(' /' ,I2,'. MAY FORCE REINPUT OF APPROPRIATE DATA.) /', 1620
1557   70000 FORMAT(' /' ,I2,'. ENTER AN INTEGER FROM 0 TO ,1X,12) 1621
1558   80000 FORMAT(' /' ,I2,'. DO YOU WANT A PRINTED COPY OF DATA ENTERED IN ', 1622
1559   80000 FORMAT(' /' ,I2,'. THE GENERAL SECTION?') 2 1623
1560   80000 FORMAT(' /' ,I2,'. THE GENERAL SECTION?') 2 1624
1561   900    RETURN 1625
1562   900    RETURN 1626
1563   900    RETURN 1627
1564   1564  SUBROUTINE ROUTS (WRT) 1628
1565   1565  C ----- 1629
1566   1566  C ORGNSM: 1630
1567   1567  C ----- 1631
1568   1568  C SUBROUTINE DOES NOT VARY 1632
1569   1569  C VARIABLES DO NOT CHANGE 1633
1570   1570  C STRUCTURE: DISPLAYS OR PRINTS DATA FROM SUBROUTINE READS 1634
1571   1571  C ----- 1635
1572   1572  C ORGNSM: 1636
1573   1573  C ----- 1637
1574   1574  C FOR OPTIONS 1 AND 2, ENTER THE QUANTITY OF ', 1638
1575   1575  C ----- 1639
1576   1576  C EACH INPUT USED EACH STAGE /) 1640
1577   1577  C ----- 1641
1578   1578  C STRUCTURE: DISPLAYS OR PRINTS DATA FROM SUBROUTINE READS 1642
1579   1579  C 1. PARAMETERS 1643
1580   1580  C 1. INCLUDE (IPAR1) 1644
1581   1581  C 2. ARGUMENTS 1645
1582   1582  C 3. COMMON BLOCKS 1646
1583   1583  C INCLUDE (ICBLK1) 1647
1584   1584  C INCLUDE (ICBLKO) 1648
1585   1585  C INCLUDE (ICUNIT) 1649
1586   1586  C LOCAL VARIABLES 1650
1587   1587  C 4. LOCAL VARIABLES 1651
1588   1588  C INCLUDE (ICUNIT) 1652
1589   1589  C ----- 1653
1590   1590  C INTEGER I, K, M, N, T, W 1654
1591   1591  C NAME DESCRIPTION OF LOCAL VARIABLES AND ARGUMENTS 1655
1592   1592  C ----- 1656
1593   1593  C ----- 1657
1594   1594  C ----- 1658
      51100 FORMAT(' /' ,I2,'. NO STAGES REQUIRE QUANTITIES OF THIS INPUT /') 1659
      50000 FORMAT(' /' ,I2,'. FOR SOME INPUTS YOU CAN CHOOSE TO ', 1660
      50000 FORMAT(' /' ,I2,'. LET THE PROGRAM COMPUTE INPUT PRICE OR /', 1661
      50000 FORMAT(' /' ,I2,'. YOU CAN CHOOSE TO INPUT THE PRICE/UNIT YOURSELF.') 1662
      50100 FORMAT(' /' ,I2,'. PRICE OF',1X,A,':') 1663
      51000 FORMAT(' /' ,I2,'. DO YOU WANT THE PROGRAM TO COMPUTE THE PRICE', 1664
      51000 FORMAT(' /' ,I2,'. OF ',A,'?') 1665
      52000 FORMAT(' /' ,I2,'. DATA FOR COMPUTING PRICE OF ',A) 1666

```

```

C K INPUT LINE NUMBER
C M LOOP COUNTER FOR MONTHS
C N LOOP COUNTER FOR INPUTS
C T LOOP COUNTER FOR TYPES OF INPUT
C W UNIT NUMBER FOR WRITE
C WRT ARGUMENT FOR UNIT NUMBER TO WRITE
C

C TRANSLATE ARGUMENTS TO LOCAL VARIABLES
W = WRT

C GENERAL INFORMATION
WRITE(W,10000) NAME, DATE
K = 1
WRITE(W,11000) K, NAMEN(BEGINM), BEGINM
K = K+1
WRITE(W,12000) K, NAMEN(ENDM), ENDM
WRITE(W,12500) I, I-BEGINI, NOOFI
K = K+1
WRITE(W,13000) K, (DAYSI(I), I=BEGINI, NOOFI)
K = K+1
WRITE(W,14000) K, (JPERI(I), I=BEGINI, NOOFI)
IF (OPTION .EQ. 1) WRITE(W,15000) K, BEGINI, DAYOP1
K = K+1
IF (OPTION .EQ. 1) WRITE(W,16000) K, BEGINI, CSOP1
IF (OPTION .EQ. 1) WRITE(W,16000) K, BEGINI, CSOP1

C CALCULATE OR INPUT PRICE?
WRITE(W,40000)
DO 500 N=1,NOOFN
WRITE(W,41000) NAMEN(N)
K = K+1
WRITE(W,42000) K, QOPNS(N)
500 CONTINUE

C LOOP THRU EACH N TO WRITE INFO WHEN PRICE OPTION WAS NOT CHOSEN
DO 600 N=1,NOOFN NAMEN(N)
WRITE(W,41000) NAMEN(N)
NUMBER AND NAME OF TYPES, T
K = K+1
IF (QOPNS(N) .EQ. 'N') THEN
  IF (OPTOK(N) .EQ. 'N') THEN
    IF (OPTOK(N) .EQ. 0) .AND. (NOOF(N) .GT. 1)
      WRITE(W,32000) K, (T, NAMEN(N,T), T=1,NOOF(N))
  ENDIF
ENDIF
K = K+1
IF (QOPNS(N) .EQ. 'N') THEN
  IF (OPTOK(N) .EQ. 0) .AND. (NOOF(N) .GT. 1)
    WRITE(W,31000) K, NOOF(N)
  ENDIF
ENDIF
DOES PRICE VARY BY MONTH?
K = K+1
IF (QOPNS(N) .EQ. 'N' .AND. (ENDM-BEGINM) .GT. 0) THEN
  IF (OPTOK(N) .EQ. 44000) K, NSBYM(N)
  ENDIF
ENDIF
PRICE OF EACH TYPE, T, IN EACH INPUT MONTH
K = K+1
IF (QOPNS(N) .EQ. 'N') THEN
  WRITE(W,45000) K, NAMEN(N), NAMEM(M), (NAMEM(M), M=BEGINM, ENDM)
ELSE
  2
  31000 FORMAT(' ',I2,'. NUMBER OF TYPES : ',I2)
  1722
  1785
  1786
  1787
  1788
  1789
  1790
  1791
  1792
  1793
  1794
  1795
  1796
  1797
  1798
  1799
  1800
  1801
  1802
  1803
  1804
  1805
  1806
  1807
  1808
  1809
  1810
  1811
  1812
  1813
  1814
  1815
  1816
  1817
  1818
  1819
  1820
  1821
  1822
  1823
  1824
  1825
  1826
  1827
  1828
  1829
  1830
  1831
  1832
  1833
  1834
  1835
  1836
  1837
  1838
  1839
  1840
  1841
  1842
  1843
  1844
  1845
  1846
  1847
  1848
  1849
  1850
  1851
  1852
  1853
  1854
  1855
  1856
  1857
  1858
  1859
  1860
  1861
  1862
  1863
  1864
  1865
  1866
  1867
  1868
  1869
  1870
  1871
  1872
  1873
  1874
  1875
  1876
  1877
  1878
  1879
  1880
  1881
  1882
  1883
  1884
  1885
  1886
  1887
  1888
  1889
  1890
  1891
  1892
  1893
  1894
  1895
  1896
  1897
  1898
  1899
  1900
  1901
  1902
  1903
  1904
  1905
  1906
  1907
  1908
  1909
  1910
  1911
  1912
  1913
  1914
  1915
  1916
  1917
  1918
  1919
  1920
  1921
  1922
  1923
  1924
  1925
  1926
  1927
  1928
  1929
  1930
  1931
  1932
  1933
  1934
  1935
  1936
  1937
  1938
  1939
  1940
  1941
  1942
  1943
  1944
  1945
  1946
  1947
  1948
  1949
  1950
  1951
  1952
  1953
  1954
  1955
  1956
  1957
  1958
  1959
  1960
  1961
  1962
  1963
  1964
  1965
  1966
  1967
  1968
  1969
  1970
  1971
  1972
  1973
  1974
  1975
  1976
  1977
  1978
  1979
  1980
  1981
  1982
  1983
  1984
  1985
  1986
  1987
  1988
  1989
  1990
  1991
  1992
  1993
  1994
  1995
  1996
  1997
  1998
  1999
  2000
  2001
  2002
  2003
  2004
  2005
  2006
  2007
  2008
  2009
  2010
  2011
  2012
  2013
  2014
  2015
  2016
  2017
  2018
  2019
  2020
  2021
  2022
  2023
  2024
  2025
  2026
  2027
  2028
  2029
  2030
  2031
  2032
  2033
  2034
  2035
  2036
  2037
  2038
  2039
  2040
  2041
  2042
  2043
  2044
  2045
  2046
  2047
  2048
  2049
  2050
  2051
  2052
  2053
  2054
  2055
  2056
  2057
  2058
  2059
  2060
  2061
  2062
  2063
  2064
  2065
  2066
  2067
  2068
  2069
  2070
  2071
  2072
  2073
  2074
  2075
  2076
  2077
  2078
  2079
  2080
  2081
  2082
  2083
  2084
  2085
  2086
  2087
  2088
  2089
  2090
  2091
  2092
  2093
  2094
  2095
  2096
  2097
  2098
  2099
  2100
  2101
  2102
  2103
  2104
  2105
  2106
  2107
  2108
  2109
  2110
  2111
  2112
  2113
  2114
  2115
  2116
  2117
  2118
  2119
  2120
  2121
  2122
  2123
  2124
  2125
  2126
  2127
  2128
  2129
  2130
  2131
  2132
  2133
  2134
  2135
  2136
  2137
  2138
  2139
  2140
  2141
  2142
  2143
  2144
  2145
  2146
  2147
  2148
  2149
  2150
  2151
  2152
  2153
  2154
  2155
  2156
  2157
  2158
  2159
  2160
  2161
  2162
  2163
  2164
  2165
  2166
  2167
  2168
  2169
  2170
  2171
  2172
  2173
  2174
  2175
  2176
  2177
  2178
  2179
  2180
  2181
  2182
  2183
  2184
  2185
  2186
  2187
  2188
  2189
  2190
  2191
  2192
  2193
  2194
  2195
  2196
  2197
  2198
  2199
  2200
  2201
  2202
  2203
  2204
  2205
  2206
  2207
  2208
  2209
  2210
  2211
  2212
  2213
  2214
  2215
  2216
  2217
  2218
  2219
  2220
  2221
  2222
  2223
  2224
  2225
  2226
  2227
  2228
  2229
  2230
  2231
  2232
  2233
  2234
  2235
  2236
  2237
  2238
  2239
  2240
  2241
  2242
  2243
  2244
  2245
  2246
  2247
  2248
  2249
  2250
  2251
  2252
  2253
  2254
  2255
  2256
  2257
  2258
  2259
  2260
  2261
  2262
  2263
  2264
  2265
  2266
  2267
  2268
  2269
  2270
  2271
  2272
  2273
  2274
  2275
  2276
  2277
  2278
  2279
  2280
  2281
  2282
  2283
  2284
  2285
  2286
  2287
  2288
  2289
  2290
  2291
  2292
  2293
  2294
  2295
  2296
  2297
  2298
  2299
  2300
  2301
  2302
  2303
  2304
  2305
  2306
  2307
  2308
  2309
  2310
  2311
  2312
  2313
  2314
  2315
  2316
  2317
  2318
  2319
  2320
  2321
  2322
  2323
  2324
  2325
  2326
  2327
  2328
  2329
  2330
  2331
  2332
  2333
  2334
  2335
  2336
  2337
  2338
  2339
  2340
  2341
  2342
  2343
  2344
  2345
  2346
  2347
  2348
  2349
  2350
  2351
  2352
  2353
  2354
  2355
  2356
  2357
  2358
  2359
  2360
  2361
  2362
  2363
  2364
  2365
  2366
  2367
  2368
  2369
  2370
  2371
  2372
  2373
  2374
  2375
  2376
  2377
  2378
  2379
  2380
  2381
  2382
  2383
  2384
  2385
  2386
  2387
  2388
  2389
  2390
  2391
  2392
  2393
  2394
  2395
  2396
  2397
  2398
  2399
  2400
  2401
  2402
  2403
  2404
  2405
  2406
  2407
  2408
  2409
  2410
  2411
  2412
  2413
  2414
  2415
  2416
  2417
  2418
  2419
  2420
  2421
  2422
  2423
  2424
  2425
  2426
  2427
  2428
  2429
  2430
  2431
  2432
  2433
  2434
  2435
  2436
  2437
  2438
  2439
  2440
  2441
  2442
  2443
  2444
  2445
  2446
  2447
  2448
  2449
  2450
  2451
  2452
  2453
  2454
  2455
  2456
  2457
  2458
  2459
  2460
  2461
  2462
  2463
  2464
  2465
  2466
  2467
  2468
  2469
  2470
  2471
  2472
  2473
  2474
  2475
  2476
  2477
  2478
  2479
  2480
  2481
  2482
  2483
  2484
  2485
  2486
  2487
  2488
  2489
  2490
  2491
  2492
  2493
  2494
  2495
  2496
  2497
  2498
  2499
  2500
  2501
  2502
  2503
  2504
  2505
  2506
  2507
  2508
  2509
  2510
  2511
  2512
  2513
  2514
  2515
  2516
  2517
  2518
  2519
  2520
  2521
  2522
  2523
  2524
  2525
  2526
  2527
  2528
  2529
  2530
  2531
  2532
  2533
  2534
  2535
  2536
  2537
  2538
  2539
  2540
  2541
  2542
  2543
  2544
  2545
  2546
  2547
  2548
  2549
  2550
  2551
  2552
  2553
  2554
  2555
  2556
  2557
  2558
  2559
  2560
  2561
  2562
  2563
  2564
  2565
  2566
  2567
  2568
  2569
  2570
  2571
  2572
  2573
  2574
  2575
  2576
  2577
  2578
  2579
  2580
  2581
  2582
  2583
  2584
  2585
  2586
  2587
  2588
  2589
  2590
  2591
  2592
  2593
  2594
  2595
  2596
  2597
  2598
  2599
  2600
  2601
  2602
  2603
  2604
  2605
  2606
  2607
  2608
  2609
  2610
  2611
  2612
  2613
  2614
  2615
  2616
  2617
  2618
  2619
  2620
  2621
  2622
  2623
  2624
  2625
  2626
  2627
  2628
  2629
  2630
  2631
  2632
  2633
  2634
  2635
  2636
  2637
  2638
  2639
  2640
  2641
  2642
  2643
  2644
  2645
  2646
  2647
  2648
  2649
  2650
  2651
  2652
  2653
  2654
  2655
  2656
  2657
  2658
  2659
  2660
  2661
  2662
  2663
  2664
  2665
  2666
  2667
  2668
  2669
  2670
  2671
  2672
  2673
  2674
  2675
  2676
  2677
  2678
  2679
  2680
  2681
  2682
  2683
  2684
  2685
  2686
  2687
  2688
  2689
  2690
  2691
  2692
  2693
  2694
  2695
  2696
  2697
  2698
  2699
  2700
  2701
  2702
  2703
  2704
  2705
  2706
  2707
  2708
  2709
  2710
  2711
  2712
  2713
  2714
  2715
  2716
  2717
  2718
  2719
  2720
  2721
  2722
  2723
  2724
  2725
  2726
  2727
  2728
  2729
  2730
  2731
  2732
  2733
  2734
  2735
  2736
  2737
  2738
  2739
  2740
  2741
  2742
  2743
  2744
  2745
  2746
  2747
  2748
  2749
  2750
  2751
  2752
  2753
  2754
  2755
  2756
  2757
  2758
  2759
  2760
  2761
  2762
  2763
  2764
  2765
  2766
  2767
  2768
  2769
  2770
  2771
  2772
  2773
  2774
  2775
  2776
  2777
  2778
  2779
  2780
  2781
  2782
  2783
  2784
  2785
  2786
  2787
  2788
  2789
  2790
  2791
  2792
  2793
  2794
  2795
  2796
  2797
  2798
  2799
  2800
  2801
  2802
  2803
  2804
  2805
  2806
  2807
  2808
  2809
  2810
  2811
  2812
  2813
  2814
  2815
  2816
  2817
  2818
  2819
  2820
  2821
  2822
  2823
  2824
  2825
  2826
  2827
  2828
  2829
  2830
  2831
  2832
  2833
  2834
  2835
  2836
  2837
  2838
  2839
  2840
  2841
  2842
  2843
  2844
  2845
  2846
  2847
  2848
  2849
  2850
  2851
  2852
  2853
  2854
  2855
  2856
  2857
  2858
  2859
  2860
  2861
  2862
  2863
  2864
  2865
  2866
  2867
  2868
  2869
  2870
  2871
  2872
  2873
  2874
  2875
  2876
  2877
  2878
  2879
  2880
  2881
  2882
  2883
  2884
  2885
  2886
  2887
  2888
  2889
  2890
  2891
  2892
  2893
  2894
  2895
  2896
  2897
  2898
  2899
  2900
  2901
  2902
  2903
  2904
  2905
  2906
  2907
  2908
  2909
  2910
  2911
  2912
  2913
  2914
  2915
  2916
  2917
  2918
  2919
  2920
  2921
  2922
  2923
  2924
  2925
  2926
  2927
  2928
  2929
  2930
  2931
  2932
  2933
  2934
  2935
  2936
  2937
  2938
  2939
  2940
  2941
  2942
  2943
  2944
  2945
  2946
  2947
  2948
  2949
  2950
  2951
  2952
  2953
  2954
  2955
  2956
  2957
  2958
  2959
  2960
  2961
  2962
  2963
  2964
  2965
  2966
  2967
  2968
  2969
  2970
  2971
  2972
  2973
  2974
  2975
  2976
  2977
  2978
  2979
  2980
  2981
  2982
  2983
  2984
  2985
  2986
  2987
  2988
  2989
  2990
  2991
  2992
  2993
  2994
  2995
  2996
  2997
  2998
  2999
  3000
  3001
  3002
  3003
  3004
  3005
  3006
  3007
  3008
  3009
  3010
  3011
  3012
  3013
  3014
  3015
  3016
  3017
  3018
  3019
  3020
  3021
  3022
  3023
  3024
  3025
  3026
  3027
  3028
  3029
  3030
  3031
  3032
  3033
  3034
  3035
  3036
  3037
  3038
  3039
  3040
  3041
  3042
  3043
  3044
  3045
  3046
  3047
  3048
  3049
  3050
  3051
  3052
  3053
  3054
  3055
  3056
  3057
  3058
  3059
  3060
  3061
  3062
  3063
  3064
  3065
  3066
  3067
  3068
  3069
  3070
  3071
  3072
  3073
  3074
  3075
  3076
  3077
  3078
  3079
  3080
  3081
  3082
  3083
  3084
  3085
  3086
  3087
  3088
  3089
  3090
  3091
  3092
  3093
  3094
  3095
  3096
  3097
  3098
  3099
  3100
  3101
  3102
  3103
  3104
  3105
  3106
  3107
  3108
  3109
  3110
  3111
  3112
  3113
  3114
  3115
  3116
  3117
  3118
  3119
  3120
  3121
  3122
  3123
  3124
  3125
  3126
  3127
  3128
  3129
  3130
  3131
  3132
  3133
  3134
  3135
  3136
  3137
  3138
  3139
  3140
  3141
  3142
  3143
  3144
  3145
  3146
  3147
  3148
  3149
  3150
  3151
  3152
  3153
  3154
  3155
  3156
  3157
  3158
  3159
  3160
  3161
  3162
  3163
  3164
  3165
  3166
  3167
  3168
  3169
  3170
  3171
  3172
  3173
  3174
  3175
  3176
  3177
  3178
  3179
  3180
  3181
  3182
  3183
  3184
  3185
  3186
  3187
  3188
  3189
  3190
  3191
  3192
  3193
  3194
  3195
  3196
  3197
  3198
  3199
  3200
  3201
  3202
  3203
  3204
  3205
  3206
  3207
  3208
  3209
  3210
  3211
  3212
  3213
  3214
  3215
  3216
  3217
  3218
  3219
  3220
  3221
  3222
  3223
  3224
  3225
  3226
  3227
  3228
  3229
  3230
  3231
  3232
  3233
  3234
  3235
  3236
  3237
  3238
  3239
  3240
  3241
  3242
  3243
  3244
  3245
  3246
  3247
  3248
  3249
  3250
  3251
  3252
  3253
  3254
  3255
  3256
  3257
  3258
  3259
  3260
  3261
  3262
  3263
  3264
  3265
  3266
  3267
  3268
  3269
  3270
  3271
  3272
  3273
  3274
  3275
  3276
  3277
  3278
  3279
  3280
  3281
  3282
  3283
  3284
  3285
  3286
  3287
  3288
  3289
  3290
  3291
  3292
  3293
  3294
  3295
  329
```

```

1787      C 1. PARAMETERS
1788      INCLUDE (IPAR1)
1789      C 2. ARGUMENTS
1790      INTEGER HREAD, RED, RED2, WRT, WRT2, WW
1791      LOGICAL RUN2
1792
1793      C 3. COMMON BLOCKS
1794      INCLUDE (ICMAIN)
1795      INCLUDE (ICBLK1)
1796      INCLUDE (ICBLK2)
1797      INCLUDE (ICUNIT)
1798      C 4. LOCAL VARIABLES
1799      INTEGER O, INFO, K, KGOTO, KLAST, NOOFK, R, W
1800      LOGICAL RUN1
1801      CHARACTER * 1 QUEST, QUEST2
1802      PARAMETER (NOOFK = 8)
1803      DIMENSION INFO(NOOFK)
1804
1805      C NAME      DESCRIPTION OF LOCAL VARIABLES AND ARGUMENTS
1806
1807      C -----
1808      C LOOP COUNTER FOR ORGANISM
1809      C READ   ARGUE: ISOSTAT TO INDICATE PROBLEMS IN READING FILE
1810      C INFO   INFORMATION LINE NUMBER
1811      C K     NUMBER OF THE INFO LINE
1812      C KGOTO  LINE NUMBER TO CORRECT
1813      C KLAST  LAST INFO LINE NUMBER TO BE INPUT
1814      C NOOFK NUMBER OF INPUT LINES
1815      C QUEST  GENERAL VARIABLE FOR USER QUESTION (IS WRITTEN OVER)
1816      C QUEST2 OPTION TO STOP, REDO, OR CONTINUE DATA ENTRY
1817      C R     UNIT NUMBER FOR READ
1818      C RED   ARGUE: UNIT NUMBERS TO READ
1819      C RED2  ARGUE: UNIT NO. TO READ FOR NEXT LEVEL SUBR.
1820      C RUN1 FIRST RUN THROUGH
1821      C RUN2 ARGUE: INDICATES IF 1ST FIRST RUN THROUGH
1822      C W     UNIT NUMBER FOR WRITE
1823      C WRT   ARGUE: FOR UNIT NUMBERS TO WRITE
1824      C WRT2  ARGUE: UNIT NO. TO WRITE FOR NEXT LEVEL SUBR.
1825      C WM    ARGUE: UNIT NO. TO WRITE TO OUTPUT FILE
1826
1827
1828
1829      C NOT FIRST RUN, IMMEDIATELY OFFER OPTION TO CHECK AND CHANGE DATA
1830      IF (RUN2) THEN
1831          DO 120 K=1,NOOFK
1832              INFO(K) = 1
1833              CONTINUE
1834          WRT2 = WI
1835          RED2 = RI
1836          W = WI
1837          R = RI
1838          WRITE(WI,60000) NAME
1839          RUN1 = .FALSE.
1840          READ(RI,1000) QUEST
1841          IF (QUEST .EQ. 'Y') CALL ROUTO(WI)
1842
1843          WRITE(WI,70000) NAME, NOOFK
1844          READ(RI,*), KGOTO
1845          IF (KGOTO .LT. 0 .OR. KGOTO .GT. NOOFK) THEN
1846              GOTO 100
1847          ELSEIF (KGOTO .GT. 0) THEN
1848              INFO (KGOTO) = 0
1849          ELSEIF (KGOTO .EQ. 0) THEN
1850              GOTO 850
1851
1852      C ORGSM:
1853      C -----
1854      C SUBROUTINE READ0 WITH ORGANISM
1855      C VARIABLES THAT MUST BE RETURNED:
1856      C THE TYPE OF THE ORGANISM : NAME(O)
1857      C THE NO. OF ORGANISMS AT BEGINNING : ONOBEG
1858      C THE TYPE OF SYSTEM USED : NAMES
1859      C CURRENT ORGANISM: BIVALVES
1860
1861      C STRUCTURE: READS IN THE ABOVE MENTIONED DATA
1862      C READS IN OTHER DATA SPECIFIC TO ORGANISM
1863      C OPTION TO CHECK, CORRECT, OR PRINT DATA
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914

```



```

1000 FORMAT (A1)
20000 FORMAT(' ',I2,'. CHOOSE ONE OF THE FOLLOWING ',A,' TYPES:')
20015 FORMAT(' ',I2,'. NAME OF THE SYSTEM USED. / ',4X,'UP TO 50 ',
2          2,'CHARACTERS IN QUOTES')
20100 FORMAT(' ',T6,I1,' = ',A)
20020 FORMAT(' ',I2,'. NUMBER OF DAYS IN THE ',A,' STAGE.')
20025 FORMAT(' ',I2,'. NUMBER OF ',A,' IN THE ',A,' STAGE')
20030 FORMAT(' ',I2,'. NUMBER OF ',A,' THAT A BATCH IS BEGUN WITH IN ',
2          2,'THE ',A,' STAGE')
40000 FORMAT(' /' THE FOLLOWING EQUATION IS USED TO EQUATE AGE IN ',
2          2,'WEEKS TO NUMBER SURVIVING: / ',4X,A)
41000 FORMAT(' ',I2,'. CHOOSE SURVIVAL PARAMETER A (SUGGESTED: ',
2          2,G14.7,')')
50000 FORMAT(' /' THE FOLLOWING EQUATION IS USED TO EQUATE AGE IN ',
2          2,'WEEKS TO SIZE IN MM: / ',4X,A)
51000 FORMAT(' ',I2,'. CHOOSE SIZE PARAMETER A (SUGGESTED: ',
2          2,G14.7,')')
52000 FORMAT(' ',I2,'. CHOOSE SIZE PARAMETER B (SUGGESTED: ',
2          2,G14.7,')')
60000 FORMAT(' /' 'DO YOU WANT TO CHECK DATA FOR ',A/)
64000 FORMAT(' ',ENTER S TO STOP EXECUTION, AFTER AN OPTION TO',
2          2,'/ ',PRINT DATA FOR ',A/
3          3,'ENTER R TO REDO ALL DATA IN THIS SECTION ',
3          3,'INTERACTIVELY '
4          4,'ENTER C TO CONTINUE EXECUTION WITH CORRECTIONS ',
5          5,'LINE BY LINE'
6          6,'ENTER S, R, OR C /')
70000 FORMAT(' ',YOU MAY CORRECT DATA LINE AT A TIME FOR ',A/
2          2,'2X,'ENTER 0 TO MAKE NO CORRECTIONS OR '/
3          3,'2X, ENTER THE NUMBER OF THE LINE YOU WANT TO CORRECT. /'
4          4,'ENTER AN INTEGER FROM 0 TO ',IX,I2)
80000 FORMAT (' /' DO YOU WANT A PRINTED COPY OF DATA ENTERED FOR ',A,
2          2,'SECTION?')
900      RETURN
        END
C -----SUBROUTINE ROUTE(WRT)
C -----ORGNSM:
C -----C
10043   C SUBROUTINE VARIES WITH ORGANISM
10044   C VARIABLES CHANGE WITH ORGANISM, BUT SOME MUST BE RETURNED
10045   C VARIABLES REQUIRED ARE THOSE REQUIRED IN READ0
10046   C CURRENT ORGANISM: BIVALVES
10047
10048   C STRUCTURE: DISPLAYS OR PRINTS DATA FROM SUBROUTINE READ0
10049
10050   C 1. PARAMETERS
10051     INCLUDE (CPAR1)
10052     C 2. ARGUMENTS
10053       INTEGER WRT
10054     C 3. COMMON BLOCKS
10055       INCLUDE (ICMAIN)
10056         INCLUDE (ICBLK1)
10057           INCLUDE (ICBLK0)
10058             INCLUDE (ICUNIT)
10059               INCLUDE (ICORG)
10060             C 4. LOCAL VARIABLES
10061               C W
10062             C NAME DESCRIPTION OF LOCAL VARIABLES AND ARGUMENTS
10063             C -----
10064               C K INPUT LINE NUMBER
10065               C W UNIT NUMBER FOR WRITE
10066               C WRT ARGUE: UNIT NUMBER FOR WRITE
10067             C -----
10068             C -----
10069
10070   C TRANSLATE ARGUMENTS TO LOCAL VARIABLES
10071   C W = WRT
10072
10073   C GENERAL INFORMATION
10074     WRITE(W,10000) NAME, DATE, NAME
10075       K = 1
10076         WRITE(W,10500) K, NAME, OTYPE, NAMEO(OTYPE)
10077           K = K+1
10078             WRITE(W,10515) K, NAMES
10079               K = K+1
10080                 WRITE(W,10520) K, NAMEI(1), DAYS1
10081                   K = K+1
10082                     WRITE(W,10525) K, NAMEI(1), NOOI1
10083                       K = K+1
10084                         WRITE(W,30000) K, NAMEO(OTYPE), ONOBEG
10085                           WRITE(W,41000) K, EQNOA
10086                           C PARAMETERS FOR SIZE/NUMBER EQUATION
10087                           K = K+1
10088                           WRITE(W,40000) EQNO(1)
10089                           C -----
10090                           WRITE(W,41000) K, EQNOA
10091
10092   C PARAMETERS FOR AGE/SIZE EQUATION
10093     WRITE(W,50000) EQSE(1)
10094       K = K+1
10095         WRITE(W,51000) K, EQSZA
10096           K = K+1
10097             WRITE(W,52000) K, EQSzb
10098             C -----
10099             C -----
10100             C COST SIMULATION MODEL: INPUT ENTERED (CONT.) /
10101               2
10102               ,A,' DATA /'
10103
10500 FORMAT(' ',I2,'. TYPE OF ',A,T46,I2,' = ',A10)
10515 FORMAT(' /' 'I2,'. NAME OF SYSTEM',T46,A)
10516
10517
10518
10519
10520
10521
10522
10523
10524
10525
10526
10527
10528
10529
10530
10531
10532
10533
10534
10535
10536
10537
10538
10539
10540
10541
10542
10543
10544
10545
10546
10547
10548
10549
10550
10551
10552
10553
10554
10555
10556
10557
10558
10559
10560
10561
10562
10563
10564
10565
10566
10567
10568
10569
10570

```

10520 FORMAT(' /' ,I2,' . NUMBER DAYS IN STAGE ',A,T46,G10.5) 2171 C STRUCTURE: READS IN DATA NECESSARY TO COMPUTE QUANTITY OF INPUT 5  
 10525 FORMAT(' /' ,I2,' . NUMBER BIVALES IN STAGE ',A,T46,G10.5) 2172 C OPTIONS TO CHECK, PRINT, CORRECT DATA  
 30000 FORMAT(' /' ,I2,' . NUMBER OF ',A,' BATCH BEGUN WITH: ',T46,G12.7) 2173 C 1. PARAMETERS  
 40000 FORMAT(' /' ,I2,' . EQUATION TO DETERMINE SURVIVAL: /' ,4X,A) 2174 C INCLUDE (IPAR1)  
 41000 FORMAT(' /' ,I2,' . PARAMETER A = ',G14.7) 2175 C 2. ARGUMENTS  
 50000 FORMAT(' /' ,I2,' . EQUATION TO EQUATE AGE TO SIZE: /' ,4X,A) 2176 C INTEGER HREAD, RED, RED2, WRT, WRT2  
 51000 FORMAT(' /' ,I2,' . PARAMETER A = ',G14.7) 2177 C LOGICAL RUN2  
 52000 FORMAT(' /' ,I2,' . PARAMETER B = ',G14.7/) 2178 C 3. COMMON BLOCKS  
 RETURN 2179 C INCLUDE (ICMAIN)  
 END 2180 C INCLUDE (ICBLK1)  
 2181 C INCLUDE (ICBLKO)  
 2182 C INCLUDE (ICUNIT)  
 2183 C INCLUDE (ICORG)  
 2184 C 4. LOCAL VARIABLES  
 2185 C INTEGER EQ, I, INFO, K, KGOTO, KLAST, M, NOOKF, R, T, W  
 2186 C LOGICAL RUN1  
 2187 C CHARACTER \* 1 QUEST, QUEST2  
 2188 C PARAMETER (NOOKF=9)  
 2189 C DIMENSION INFO(NOOKF)  
 2190 C NAME DESCRIPTION OF LOCAL VARIABLES AND ARGUMENTS  
 2191 C-----  
 2192 C-----  
 2193 C-----  
 2194 C EQ LOOP COUNTER FOR EQUATIONS  
 2195 C HREAD ARGUE: IOSTAT TO INDICATE PROBLEMS READING FILE  
 2196 C I LOOP COUNTER FOR INPUTS  
 2197 C INFO INDICATOR OF INPUT LINE  
 2198 C K INPUT LINE NUMBER  
 2199 C KGOTO INPUT LINE NO. TO GOTO TO CORRECT  
 2200 C KLAST LAST INPUT LINE INPUT  
 2201 C M LOOP COUNTER FOR MONTHS  
 2202 C NOOKF NUMBER OF INPUT LINES  
 2203 C QUEST GENERAL INTERACTIVE USER QUESTION (IS OVERWRITTEN)  
 2204 C QUEST2 OPTION TO STOP, REDO, OR CONTINUE  
 2205 C R UNIT NUMBER FOR READ  
 2206 C RED ARGUE: UNIT NUMBER FOR READ  
 2207 C RED2 ARGUE: UNIT NUMBER FOR READ FOR NEXT LEVEL SUBR.  
 2208 C RUN1 IS IT THE FIRST RUN?  
 2209 C RUN2 ARGUE: IS IT GREATER THAN THE FIRST RUN  
 2210 C T LOOP COUNTER FOR TYPES  
 2211 C W UNIT NUMBER FOR WRITE  
 2212 C WRT ARGUE: UNIT NUMBER FOR WRITE  
 2213 C WRT2 ARGUE: UNIT NUMBER TO WRITE TO NEXT LEVEL SUBROUTINE  
 2214 C-----  
 2215 C-----  
 2216 C-----  
 2217 C IF NOT 1ST RUN, IMMEDIATELY OFFER TO DISPLAY AND CHANGE DATA  
 2218 C IF (RUN2) THEN  
 2219 C DO 120 K=1,NOOKF  
 2220 C INFO(K) = 1  
 2221 C-----  
 120 CONTINUE  
 2222 C-----  
 2223 C-----  
 2224 C-----  
 2225 C-----  
 2226 C-----  
 2227 C-----  
 2228 C-----  
 2229 C-----  
 2230 C-----  
 2231 C-----  
 2232 C-----  
 2233 C-----  
 2234 C-----  
 C ORGNSM:  
 C-----  
 C SUBROUTINE VARIES WITH ORGANISM  
 C-----  
 C VARIABLES CHANGE WITH ORGANISM  
 C-----  
 C VARIABLES NOT RETURNED TO ORGNSM  
 C-----  
 C VARIABLES NECESSARY ONLY FOR QUANT5, IF OPTION TO COMPUTE QUANTITY=Y  
 C-----  
 C CURRENT ORGANISM: BIVALES

```

ELSEIF (KGOTO .GT. 0) THEN
  INFO(KGOTO) = 0
ELSEIF (KGOTO .EQ. 0) THEN
  GOTO 850
ENDIF

C INITIALIZE VARIABLES TO 0 TO READ IN DATA FIRST TIME
C ASSIGN ARGUE UNIT NO. TO LOCAL UNIT NUMBERS
DO 110 K=1,NOOFK
  INFO(K) = 0
CONTINUE
RUN1 = .TRUE.
W = WRT
R = RED
WRT2 = WRT
RED2 = RED
ENDIF

C READ IN ALL DATA
400  READ = 0
C CHOOSE ALGORITHM AND PARAMETERS TO COMPUTE QUANTITY OF ALGAE
K = 1
IF (INFO(K) .EQ. 0) THEN
  WRITE(W,10000) K
  READ(R,* ,ERR=900, IOSTAT=HREAD)
  INFO(K+1) = 0
  INFO(K+2) = 0
ENDIF
K = K+1
IF (INFO(K) .EQ. 0) THEN
  WRITE(W,10100) (EQ, EQQ5(EQ), EQ=1,2)
  READ(R,* ,ERR=900, IOSTAT=HREAD)
  INFO(K+1) = 0
  INFO(K+2) = 0
ENDIF
K = K+1
IF (INFO(K) .EQ. 0) THEN
  WRITE(W,11000) K, EQQ5AF(QEQQ5)
  READ(R,* ,ERR=900, IOSTAT=HREAD)
  EQQ5A(QEQQ5)
ENDIF
K = K+1
IF (INFO(K) .EQ. 0) THEN
  WRITE(W,11100) K, EQQ5BF(QEQQ5)
  READ(R,* ,ERR=900, IOSTAT=HREAD)
  EQQ5B(QEQQ5)
ENDIF

C FACTORS AFFECTING QUANTITY REQUIRED
C FEEDING FILTERING RATE
K = K+1
IF (INFO(K) .EQ. 0) THEN
  DO 200 I=BEGIN,NOOFI
    IF (OPNNEW(I,5) .EQ. 'C') THEN
      WRITE(W,20100) I, NAMEI(I)
      READ(R,* ,ERR=900, IOSTAT=HREAD)
      FILTER(I)
    ENDIF
  CONTINUE
200  Continue
ENDIF

C FRACTION OF EACH TYPE OF ALGAE (IF < 1) USED IN EACH STAGE
K = K+1
IF (INFO(K) .EQ. 0) THEN
  IF (NOOFT(5) .GT. 1) THEN
    WRITE(W,21000) K
    DO 210 I=BEGINI,NOOFI
      IF (OPNNEW(I,5) .EQ. 'C') THEN
        NSREQ(I,I,T)
        READ(R,* ,ERR=900, IOSTAT=HREAD)
        NSREQ(I,T)
      CONTINUE
    ENDIF
210  Continue
ENDIF

C AVAILABILITY OF ALGAE BY MONTH FROM SEAWATER, PRODUCED AND PURCHASED
2299  WRITE(W,20100) I, NAMEI(I)
  DO 211 T=1,NOOFT(5)
    IF (NOOFT(5) .GT. 1) WRITE(W,20200) NAMEI(5,T)
    READ(R,* ,ERR=900, IOSTAT=HREAD)
    NSREQ(I,T)
  CONTINUE
2300  Continue
2301  DO 220 I=BEGINI,NOOFT
    IF (OPNNEW(I,5) .EQ. 'C') THEN
      NSREQ(I,1,1) = 1.
    ENDIF
  CONTINUE
2302  Continue
2303  211  Continue
2304  ENDIF
2305  210  Continue
2306  ELSE
    DO 220 I=BEGINI,NOOFT
      IF (OPNNEW(I,5) .EQ. 'C') THEN
        NSREQ(I,1,1) = 1.
      ENDIF
    CONTINUE
2307  ENDIF
2308  220  Continue
2309  ENDIF
2310  2311  Continue
2311  ENDIF
2312  ENDIF
2313  ENDIF
2314  C AVAILABILITY OF ALGAE BY MONTH FROM SEAWATER, PRODUCED AND PURCHASED
2315  READ(R,10000,ERR=900, IOSTAT=HREAD)
  SEABYM
2316  K = K+1
  IF (INFO(K) .EQ. 0) THEN
    IF ('ENDM-BEGINM) .GT. 0) THEN
      WRITE(W,30000) K
      READ(R,10000,ERR=900, IOSTAT=HREAD)
      SEABYM
    ELSE
      SEABYM = '-'
    ENDIF
  INFO(K+1) = 0
  ENDIF
2317  2318  IF ('ENDM-BEGINM) .GT. 0) THEN
    WRITE(W,30100) K
    DO 300 I=BEGINI,NOOFI
      IF (OPNNEW(I,5) .EQ. 'C') THEN
        WRITE(W,20100) I, NAMEI(I)
        IF (SEABYM .EQ. 'Y') THEN
          WRITE(W,20200) NAMEI(I)
        ELSE
          WRITE(W,30100) I, NAMEI(I)
        ENDIF
      CONTINUE
    ENDIF
  ELSE
    WRITE(W,30100) K
    DO 301 M=BEGINM,ENDM
      IF ('ENDM-BEGINM) .GT. 0) THEN
        WRITE(W,30200) NAMEM(M)
        READ(R,* ,ERR=900, IOSTAT=HREAD)
        SEACEL(I,M)
      ELSE
        READ(R,* ,ERR=900, IOSTAT=HREAD)
        SEACEL(I,1)
      ENDIF
    ENDIF
  ENDIF
2319  2320  READ(R,10000,ERR=900, IOSTAT=HREAD)
  SEABYM
2321  2322  SEABYM = '-'
  ENDIF
2323  2324  ENDIF
2324  2325  ENDIF
2325  K = K+1
  IF (INFO(K) .EQ. 0) THEN
    WRITE(W,30100) K
    DO 300 I=BEGINI,NOOFI
      IF (OPNNEW(I,5) .EQ. 'C') THEN
        WRITE(W,20100) I, NAMEI(I)
        IF (SEABYM .EQ. 'Y') THEN
          WRITE(W,20200) NAMEI(I)
        ELSE
          WRITE(W,30100) I, NAMEI(I)
        ENDIF
      CONTINUE
    ENDIF
  ELSE
    WRITE(W,30100) K
    DO 301 M=BEGINM,ENDM
      IF ('ENDM-BEGINM) .GT. 0) THEN
        WRITE(W,30200) NAMEM(M)
        READ(R,* ,ERR=900, IOSTAT=HREAD)
        SEACEL(I,M)
      ELSE
        READ(R,* ,ERR=900, IOSTAT=HREAD)
        SEACEL(I,1)
      ENDIF
    ENDIF
  ENDIF
2326  2327  READ(R,* ,ERR=900, IOSTAT=HREAD)
  SEACEL(I,1)
2328  2329  SEACEL(I,M) = SEACEL(I,1)
  CONTINUE
2329  2330  READ(R,* ,ERR=900, IOSTAT=HREAD)
  SEACEL(I,M)
2330  2331  WRITE(W,20100) I, NAMEI(I)
  CONTINUE
2331  2332  READ(R,* ,ERR=900, IOSTAT=HREAD)
  SEACEL(I,1)
2332  2333  DO 301 M=BEGINM,ENDM
    SEACEL(I,M) = SEACEL(I,1)
  CONTINUE
2333  2334  READ(R,* ,ERR=900, IOSTAT=HREAD)
  SEACEL(I,1)
2334  2335  READ(R,* ,ERR=900, IOSTAT=HREAD)
  SEACEL(I,M)
2335  301  READ(R,10000,ERR=900, IOSTAT=HREAD)
  NSABYM
2336  300  Continue
2337  2338  READ(R,* ,ERR=900, IOSTAT=HREAD)
  SEACEL(I,1)
2338  2339  SEACEL(I,M) = SEACEL(I,1)
  CONTINUE
2339  2340  READ(R,* ,ERR=900, IOSTAT=HREAD)
  SEACEL(I,1)
2340  302  READ(R,* ,ERR=900, IOSTAT=HREAD)
  SEACEL(I,M)
2341  2342  READ(R,* ,ERR=900, IOSTAT=HREAD)
  SEACEL(I,M)
2342  2343  READ(R,* ,ERR=900, IOSTAT=HREAD)
  SEACEL(I,M)
2343  2344  READ(R,* ,ERR=900, IOSTAT=HREAD)
  SEACEL(I,M)
2344  2345  READ(R,* ,ERR=900, IOSTAT=HREAD)
  SEACEL(I,M)
2345  2346  READ(R,* ,ERR=900, IOSTAT=HREAD)
  SEACEL(I,M)
2346  2347  C QUANTITY OF ALGAE PRODUCED IF PRICE OPTION NOT CHOSEN
  C IF PRICE OPTION CHOSEN, THEN PRICE SUBR. SHOULD DETERMINE QUANT.
2347  2348  K = K+1
  IF (INFO(K) .EQ. 0) THEN
    IF ('QOPEN$5) .EQ. 'N') THEN
      IF ('ENDM-BEGINM) .GT. 0) THEN
        WRITE(W,31000) K
        READ(R,10000,ERR=900, IOSTAT=HREAD)
        NSABYM
      ELSE
        NSABYM = '-'
      ENDIF
    ENDIF
  ENDIF
2348  2349  IF (INFO(K) .EQ. 0) THEN
    IF ('QOPEN$5) .EQ. 'N') THEN
      IF ('ENDM-BEGINM) .GT. 0) THEN
        WRITE(W,31000) K
        READ(R,10000,ERR=900, IOSTAT=HREAD)
        NSABYM
      ELSE
        NSABYM = '-'
      ENDIF
    ENDIF
  ENDIF
2349  2350  IF (INFO(K) .EQ. 0) THEN
    IF ('QOPEN$5) .EQ. 'N') THEN
      IF ('ENDM-BEGINM) .GT. 0) THEN
        WRITE(W,31000) K
        READ(R,10000,ERR=900, IOSTAT=HREAD)
        NSABYM
      ELSE
        NSABYM = '-'
      ENDIF
    ENDIF
  ENDIF
2350  2351  IF (INFO(K) .EQ. 0) THEN
    IF ('QOPEN$5) .EQ. 'N') THEN
      IF ('ENDM-BEGINM) .GT. 0) THEN
        WRITE(W,31000) K
        READ(R,10000,ERR=900, IOSTAT=HREAD)
        NSABYM
      ELSE
        NSABYM = '-'
      ENDIF
    ENDIF
  ENDIF
2351  2352  IF (INFO(K) .EQ. 0) THEN
    IF ('QOPEN$5) .EQ. 'N') THEN
      IF ('ENDM-BEGINM) .GT. 0) THEN
        WRITE(W,31000) K
        READ(R,10000,ERR=900, IOSTAT=HREAD)
        NSABYM
      ELSE
        NSABYM = '-'
      ENDIF
    ENDIF
  ENDIF
2352  2353  IF (INFO(K) .EQ. 0) THEN
    IF ('QOPEN$5) .EQ. 'N') THEN
      IF ('ENDM-BEGINM) .GT. 0) THEN
        WRITE(W,31000) K
        READ(R,10000,ERR=900, IOSTAT=HREAD)
        NSABYM
      ELSE
        NSABYM = '-'
      ENDIF
    ENDIF
  ENDIF
2353  2354  IF (INFO(K) .EQ. 0) THEN
    IF ('QOPEN$5) .EQ. 'N') THEN
      IF ('ENDM-BEGINM) .GT. 0) THEN
        WRITE(W,31000) K
        READ(R,10000,ERR=900, IOSTAT=HREAD)
        NSABYM
      ELSE
        NSABYM = '-'
      ENDIF
    ENDIF
  ENDIF
2354  2355  IF (INFO(K) .EQ. 0) THEN
    IF ('QOPEN$5) .EQ. 'N') THEN
      IF ('ENDM-BEGINM) .GT. 0) THEN
        WRITE(W,31000) K
        READ(R,10000,ERR=900, IOSTAT=HREAD)
        NSABYM
      ELSE
        NSABYM = '-'
      ENDIF
    ENDIF
  ENDIF
2355  2356  IF (INFO(K) .EQ. 0) THEN
    IF ('QOPEN$5) .EQ. 'N') THEN
      IF ('ENDM-BEGINM) .GT. 0) THEN
        WRITE(W,31000) K
        READ(R,10000,ERR=900, IOSTAT=HREAD)
        NSABYM
      ELSE
        NSABYM = '-'
      ENDIF
    ENDIF
  ENDIF
2356  2357  IF (INFO(K) .EQ. 0) THEN
    IF ('QOPEN$5) .EQ. 'N') THEN
      IF ('ENDM-BEGINM) .GT. 0) THEN
        WRITE(W,31000) K
        READ(R,10000,ERR=900, IOSTAT=HREAD)
        NSABYM
      ELSE
        NSABYM = '-'
      ENDIF
    ENDIF
  ENDIF
2357  2358  IF (INFO(K) .EQ. 0) THEN
    IF ('QOPEN$5) .EQ. 'N') THEN
      IF ('ENDM-BEGINM) .GT. 0) THEN
        WRITE(W,31000) K
        READ(R,10000,ERR=900, IOSTAT=HREAD)
        NSABYM
      ELSE
        NSABYM = '-'
      ENDIF
    ENDIF
  ENDIF
2358  2359  IF (INFO(K) .EQ. 0) THEN
    IF ('QOPEN$5) .EQ. 'N') THEN
      IF ('ENDM-BEGINM) .GT. 0) THEN
        WRITE(W,31000) K
        READ(R,10000,ERR=900, IOSTAT=HREAD)
        NSABYM
      ELSE
        NSABYM = '-'
      ENDIF
    ENDIF
  ENDIF
2359  2360  IF (INFO(K) .EQ. 0) THEN
    IF ('QOPEN$5) .EQ. 'N') THEN
      IF ('ENDM-BEGINM) .GT. 0) THEN
        WRITE(W,31000) K
        READ(R,10000,ERR=900, IOSTAT=HREAD)
        NSABYM
      ELSE
        NSABYM = '-'
      ENDIF
    ENDIF
  ENDIF
2360  2361  IF (INFO(K) .EQ. 0) THEN
    IF ('QOPEN$5) .EQ. 'N') THEN
      IF ('ENDM-BEGINM) .GT. 0) THEN
        WRITE(W,31000) K
        READ(R,10000,ERR=900, IOSTAT=HREAD)
        NSABYM
      ELSE
        NSABYM = '-'
      ENDIF
    ENDIF
  ENDIF
2361  2362  IF (INFO(K) .EQ. 0) THEN
    IF ('QOPEN$5) .EQ. 'N') THEN
      IF ('ENDM-BEGINM) .GT. 0) THEN
        WRITE(W,31000) K
        READ(R,10000,ERR=900, IOSTAT=HREAD)
        NSABYM
      ELSE
        NSABYM = '-'
      ENDIF
    ENDIF
  ENDIF
2362

```

```

IF (QDFNS(5) .EQ. 'N') THEN
  WRITE(W,31100) K
  DO 311 T=1,NOOFT(5)
    WRITE(W,20200) NAMET(5,T)
    IF ('NABYM' EQ. 'Y') THEN
      DO 312 M=BEGNM,ENDM
        WRITE(W,30200) NAMEM(M)
        READ(R,* ,ERR=900,IOSTAT=HREAD) N5AVAL(T,M)
        CONTINUE
        READ(R,* ,ERR=900,IOSTAT=HREAD) N5AVAL(T,1)
      ELSE
        READ(R,* ,ERR=900,IOSTAT=HREAD) N5AVAL(T,1)
        CONTINUE
        READ(R,* ,ERR=900,IOSTAT=HREAD) N5AVAL(T,1)
        CONTINUE
        ENDIF
      ENDIF
    ENDIF
  ENDIF
  KLAST = K
  WRT2 = WI
  RED2 = RI
  W = WI
  R = RI
  C OPTION TO CHECK DATA
  WRITE(WI,60000) NAMEN(5)
  READ(RI,1000) QUEST
  IF (QUEST .EQ. 'Y') THEN
    CALL ROUTNS(WI)
  ENDIF
  C OPTION TO STOP OR REDO
  IF (RUN1) THEN
    WRITE(WI,64000) NAMEN(5)
    READ(RI,1000) QUEST2
    IF (QUEST2 .EQ. 'S') THEN
      GOTO 850
    ELSEIF (QUEST2 .EQ. 'R') THEN
      GOTO 400
    ELSEIF (QUEST .EQ. 'C') THEN
      CONTINUE
    ENDIF
    RUN1 = .FALSE.
  ENDIF
  C REASSIGN 1 TO INPUT LINES TO STOP FROM REENTERING DATA
  DO 800 K=1,NOOK
    INFO(K) = 1
    CONTINUE
  C OPTION TO CORRECT DATA, LINE BY LINE, BY CHOOSING A LINE
  810  WRITE(WI,70000) NAMEN(5), NOOK
    READ(RI,* ) KGOTO
    IF (KGOTO .LT. 0 .OR. KGOTO .GT. NOOK) THEN
      GOTO 810
    ELSEIF (KGOTO .GT. 0) THEN
      INFO(KGOTO) = 0
      GOTO 400
    ENDIF
    C OPTION TO PRINT THE DATA ENTERED
    800 FORMAT(' ',7X,A)
    2479 30200 FORMAT(' ',7X,A)
    2480 31000 FORMAT(' ',12,'. DOES THE AVAILABILITY OF ALGAE PRODUCED ',2
    2481 31000 FORMAT(' ',12,'. TOTAL AMOUNT OF EACH TYPE OF ALGAE AVAILABLE ')
    2482 31000 FORMAT(' ',12,'. CHANGE BY MONTH? ')
    2483 2484 31100 FORMAT(' ',12,'. TOTAL AMOUNT OF EACH TYPE OF ALGAE AVAILABLE ')
    2485 2486 60000 FORMAT(' ',12,'. DO YOU WANT TO CHECK THE DATA ENTERED FOR ',A)
    2487 2488 64000 FORMAT(' ',12,'. ENTER S TO STOP EXECUTION AFTER AN OPTION TO ',2
    2489 2490 3 2 3 , 'PRINT DATA FOR ',A/
    3 , 'ENTER R TO REDO ALL ALGAE DATA INTERACTIVELY / '
  850  WRITE(WI,80000) NAMEN(5)
  2427  READ(RI,1000) QUEST
  2428  IF (QUEST .EQ. 'Y') THEN
  2429    CALL ROUTNS(WO)
  2430  ENDIF
  2431  IF (QUEST2.EQ.'S') STOP 'PROGRAM STOP IN DATA ENTRY INPUT 5'
  2432  C OPTION TO PRINT OUT ADDITIONAL INFORMATION ABOUT INPUT 5
  2433  IF (OUTQN(5).EQ.0) THEN
  2434    WRITE(WI,90000)
  2435    READ(RI,1000) OUTQN(5)
  2436    IF (OUTQN(5).EQ.0) THEN
  2437      WRITE(WI,10000)
  2438      IF (OUTQN(5).EQ.'Y') THEN
  2439        WRITE(WI,91000)
  2440        READ(RI,* ) WO
  2441      ENDIF
  2442  ENDIF
  2443  C 1000 FORMAT(' ',12,'. TO COMPUTE ALGAE QUANTITY, CHOOSE ONE OF ',2
  2444  10000 FORMAT(' ',12,'. THE FOLLOWING EQUATIONS / ',4X,' THAT BEST ',2
  2445  10000 FORMAT(' ',12,'. RELATES AGE IN WEEKS TO 10,000 CELLS CONSUMED.',2
  2446  10000 FORMAT(' ',12,'. 4X, THE EQUATION WILL BE SOLVED FOR THE NUMBER ',2
  2447  10000 FORMAT(' ',12,'. OF CELLS CONSUMED PER DAY. ')
  2448  10000 FORMAT(' ',12,'. CHOOSE ALGAE PARAMETER A (SUGGESTED: ',2
  2449  10000 FORMAT(' ',12,'. G14.7, ')
  2450  10000 FORMAT(' ',12,'. CHOOSE ALGAE PARAMETER B (SUGGESTED: ',2
  2451  10000 FORMAT(' ',12,'. G14.7, ')
  2452  10000 FORMAT(' ',12,'. WHAT IS THE FEEDING FILTERING RATE OF THE ',2
  2453  10000 FORMAT(' ',12,'. ORGANISM AT EACH STAGE ',2
  2454  10000 FORMAT(' ',12,'. ,4X, FOR WHICH ALGAE QUANTITY IS TO BE COMPUTED ')
  2455  20000 FORMAT(' ',12,'. IN STAGE ',12,', ',A)
  2456  20000 FORMAT(' ',12,'. IN STAGE ',12,', ',A)
  2457  20000 FORMAT(' ',12,'. IN STAGE ',12,', ',A)
  2458  20100 FORMAT(' ',12,'. WHAT IS THE FRACTION OF EACH TYPE OF ALGAE ',2
  2459  20100 FORMAT(' ',12,'. REQUIRED IN EACH STAGE (TOTAL 100%): ')
  2460  20200 FORMAT(' ',12,'. OF TYPE ',A)
  2461  20200 FORMAT(' ',12,'. OF TYPE ',A)
  2462  20200 FORMAT(' ',12,'. DOES THE AVAILABILITY OF ALGAE FROM ',2
  2463  20200 FORMAT(' ',12,'. SEAWATER CHANGE BY MONTH? ')
  2464  20200 FORMAT(' ',12,'. FRACTION OF DIET BIVALVES RECEIVE FROM ',2
  2465  20200 FORMAT(' ',12,'. SEAWATER: ')
  2466  20200 FORMAT(' ',12,'. ')
  2467  20200 FORMAT(' ',12,'. ')
  2468  20200 FORMAT(' ',12,'. ')
  2469  20200 FORMAT(' ',12,'. ')
  2470  20200 FORMAT(' ',12,'. ')
  2471  20200 FORMAT(' ',12,'. ')
  2472  30000 FORMAT(' ',12,'. ')
  2473  30000 FORMAT(' ',12,'. ')
  2474  30000 FORMAT(' ',12,'. ')
  2475  30100 FORMAT(' ',12,'. ')
  2476  30100 FORMAT(' ',12,'. ')
  2477  30100 FORMAT(' ',12,'. ')
  2478  30200 FORMAT(' ',12,'. ')
  2479  30200 FORMAT(' ',12,'. ')
  2480  30200 FORMAT(' ',12,'. ')
  2481  30200 FORMAT(' ',12,'. ')
  2482  30200 FORMAT(' ',12,'. ')
  2483  30200 FORMAT(' ',12,'. ')
  2484  30200 FORMAT(' ',12,'. ')
  2485  30200 FORMAT(' ',12,'. ')
  2486  30200 FORMAT(' ',12,'. ')
  2487  30200 FORMAT(' ',12,'. ')
  2488  30200 FORMAT(' ',12,'. ')
  2489  30200 FORMAT(' ',12,'. ')
  2490  30200 FORMAT(' ',12,'. ')

```

```

4      ' ' ENTER C TO CONTINUE WITH OPTION TO CORRECT '
5      ' ' LINE BY LINE'/
6      ' ' ENTER S, R, OR C'
70000 FORMAT (' /' Y YOU MAY NOW CORRECT DATA A LINE AT A TIME FOR ',A/
     2      ',2X,'ENTER 0 TO MAKE NO CORRECTIONS, OR /
     3      ',2X,'ENTER THE NO. OF THE LINE YOU WANT TO CORRECT'/
     4      ',ENTER AN INTEGER FROM 0 TO ,1X,I2)
80000 FORMAT (' /' DO YOU WANT A PRINTED COPY OF DATA ENTERED FOR ',A,
     2      ',? Y OR N')
90000 FORMAT(' /' DO YOU WANT TO PRINT ADDITIONAL ',
     2      ' INFORMATION ABOUT ALGAE? Y OR N')
91000 FORMAT(' /' TO WHAT OUTPUT FILE UNIT DO YOU WANT THIS TO GO TO?',',
     2      '- INTEGER 10 TO 99,'/ (MAIN OUTPUT FILE HAS ',
     3      'A UNIT = ',I2,')')
900   RETURN
END

SUBROUTINE READN6(WRT,RED,HREAD,RUN2)
RETURN
END

SUBROUTINE READN(WRT,RED,HREAD,RUN2)
RETURN
END

SUBROUTINE ROUTN1(WRT)
RETURN
END

SUBROUTINE ROUTN2(WRT)
RETURN
END

SUBROUTINE ROUTN3(WRT)
RETURN
END

SUBROUTINE ROUTN4(WRT)
RETURN
END

SUBROUTINE ROUTN5(WRT)

```

60

---

```

2555      C -----
2556      C
2557      C ORGNSM;
2558      C -----
2559      C SUBROUTINE VARIES WITH ORGANISM
2560      C VARIABLES CHANGE, BUT NOT RETURNED TO ORGNSM
2561      C VARIABLES REQUIRED ARE THOSE TO PRINT OUT VARIABLES ENTERED
2562      C IN READN5, WHICH ARE SPECIFIC TO THE ORGANISM
2563      C CURRENT ORGANISM: BIVALVES
2564      C STRUCTURE: DISPLAYS OR PRINTS OUT DATA FOR SUBROUTINE READN5
2565      C
2566      C 1. PARAMETERS
2567      C     INCLUDE(IPAR1)
2568      C 2. ARGUMENTS
2569      C     INTEGER WRT
2570      C     COMMON BLOCKS
2571      C     INCLUDE(UCBLK1)
2572      C     INCLUDE(ICMAIN)
2573      C     INCLUDE(ICORG)
2574      C     LOCAL VARIABLES
2575      C     INTEGER I, K, M, T, W
2576      C
2577      C 3. COMMON BLOCKS
2578      C     NAME    DESCRIPTION OF LOCAL VARIABLES AND ARGUMENTS
2579      C     -----
2580      C     -----
2581      C     I        LOOP COUNTER FOR STAGE
2582      C     K        INPUT LINE NUMBER
2583      C     M        LOOP COUNTER FOR MONTH
2584      C     T        LOOP COUNTER FOR TYPES
2585      C     W        UNIT NUMBER FOR WRITE
2586      C     WRT      ARGUMENT FOR UNIT NUMBER TO WRITE
2587      C
2588      C
2589      C TRANSLATE ARGUMENTS TO LOCAL VARIABLES
2590      C     W = WRT
2591      C
2592      C
2593      C WRITE OUT HEADING
2594      C     WRITE(W,5000) NAME, DATE, NAMEN(5)
2595      C
2596      C ALGORITHM TO COMPUTE QUANTITY
2597      C     K = 1
2598      C     WRITE(W,10000) K, QEQQ5, EQQ5(QEQQ5)
2599      C     K = K+1
2600      C     WRITE(W,11000) K, EQQ5A(QEQQ5)
2601      C     K = K+1
2602      C     WRITE(W,11100) K, EQQ5B(QEQQ5)
2603      C
2604      C     WRITE(W,20000) (I, I=BEGINI,NOOFI)
2605      C
2606      C QUANTITY REQUIRED
2607      C FEEDING FILTERING RATE
2608      C
2609      C     WRITE(W,21000) K, (FILTER(I), I=BEGINI,NOOFI)
2610      C     K = K + 1
2611      C     IF (NOOFT(5) .GT. 1) THEN
2612      C       WRITE(W,21100) K
2613      C       DO 211 T=1,NOOFT(5)
2614      C         IF (NOOFT(5) .GT. 1) WRITE(W,21200) NAMET(5,T)
2615      C
2616      C     WRITE(W,21300) (NSREQ(I,T), I=BEGINI,NOOFI)
2617      C     CONTINUE(W,21300)
2618      C
END
```

61

```

C AVAILABILITY
K = K+1
  IF ((ENDM-BEGINM) .GT. 0) WRITE(W,300000) K, SEABYM
K = K+1
  IF ((ENDM-BEGINM) .LT. 0) WRITE(W,30100) K
  DO 300 M=BIGNM,ENDM
    WRITE(W,30200) NAME(M), (SEACEL(I,M), I=BEGINI,NOIFI)
  CONTINUE
300
  K = K+1
  IF (QOPNS(5).EQ. 'N' .AND. (ENDM-BEGINM) .GT. 0) THEN
    WRITE(W,31000) K, NSABYM
  ENDIF
  K = K+1
  IF (QOPNS(5) .EQ. 'N') THEN
    WRITE(W,31100) K, (NAME(M), M=BEGINM,ENDM)
  END
  DO 310 T=1,NOOPT(5)
    IF (NOOPT(5) .GT. 1) WRITE(W,21200) NAMET(5,T)
    WRITE(W,31300) (NSAVAL(T,M), M=BEGINM,ENDM)
  CONTINUE
310
  ENDIF

5000 FORMAT('1',A,' COST SIMULATION MODEL: INPUT ENTERED (CONT.) /',A50//',A,IX,'DATA' /)
10000 FORMAT(' ',I2,' AGE/ALGAE EQUATION CHOSEN: ',I1/' ',4X,A)
11000 FORMAT(' ',I2,' PARAMETER A = ',G14.7)
11100 FORMAT(' ',I2,' PARAMETER B = ',G14.7)
20000 FORMAT(' ',IN THE FOLLOWING, WHERE QUANTITY=0, QUANTITY IS ',I1/' ',I2,' ENTERD//',6X,6(I2,' STAGE',2X)/)
21000 FORMAT(' ',I2,' FILTERING RATE//',6X,6(G9.2,1X))
21100 FORMAT(' ',I2,' FRACTION OF EACH ALGAE TYPE REQUIRED ',I1/' ',I2,' PER STAGE: ')
21200 FORMAT(' ',4X,' TYPE ',A)
21300 FORMAT(' ',6X,6(G9.2,1X))
30000 FORMAT(' ',I2,' FRACTION FROM SEA VARIES BY MONTH: ',A1)
30100 FORMAT(' ',I2,' FRACTION OF DIET FROM SEA ALGAE IN EACH ',I1/' ',I2,' STAGE, EACH MONTH: ')
30200 FORMAT(' ',6X,A//',6X,6(G9.2,1X)/)
31000 FORMAT(' ',I2,' AVAILABILITY OF ALGAE VARIES BY MONTH: ',A)
31100 FORMAT(' ',I2,' AVAILABILITY EACH TYPE BY MONTH: ')
31200 FORMAT(' ',6X,A//',6X,6(A10,1X)//',6X,6(A10,1X))
31300 FORMAT(' ',6X,6(G9.2,1X)//',6X,6(G9.2,1X))

SUBROUTINE ROUTN6 (WRT)
  RETURN
END

SUBROUTINE PRICE1 (WRT,RED,HREAD,RUN2,RUN3)
  RETURN
END

SUBROUTINE PRICE2 (WRT,RED,HREAD,RUN2,RUN3)
  RETURN
END

SUBROUTINE PRICE3 (WRT,RED,HREAD,RUN2,RUN3)
  RETURN
END

SUBROUTINE PRICE4 (WRT,RED,HREAD,RUN2,RUN3)
  RETURN
END

SUBROUTINE PRICE5 (WRT,RED,HREAD,RUN2,RUN3)
  -----
C ORGNSM
  -----
C SUBROUTINE VARIES WITH ORGANISM
C VARIABLES CHANGE WITH ORGANISM
C VARIABLES NEED BE RETURNED (ONLY IF PRICE OPTION CHOSEN):
C NUMBER OF TYPES - NOOPT(5)
C NAME OF EACH TYPE - NAMET(5,T)
C PRICE PER LITRE - PRICET(5,I,M)
C AMOUNT AVAILABLE- N5AVAI(T,M),
C CURRENT ORGANISM: BIVALVE
C STRUCTURE: MAIN SUBROUTINE, PRICES, AND 8 LESSER SUBROUTINES
C MAIN SUBROUTINE DOES THE FOLLOWING:
C -CALLS SUBROUTINE RINNS, ROUT$5
C -LOOPS THROUGH PRODUCTION MONTHS
C   LOOPS THROUGH PRODUCTION STAGES
C     CALCULATES AVAILABLE ALGAE QUANTITY
C     UPDATES CURRENT PRODUCTION MONTH NJ
C     CALLS SUBROUTINE ELEC TO FIND KWHR
C     CALLS SUBROUTINE FUEL TO FIND LITRES OF OIL
C
2747      2683
2748      2684
2749      2685
2750      2686
2751      2687
2752      2688
2753      2689
2754      2690
2755      2691
2756      2692
2757      2693
2758      2694
2759      2695
2760      2696
2761      2697
2762      2698
2763      2699
2764      2700
2765      2701
2766      2702
2767      2703
2768      2704
2769      2705
2770      2706
2771      2707
2772      2708
2773      2709
2774      2710
2775      2711
2776      2712
2777      2713
2778      2714
2779      2715
2780      2716
2781      2717
2782      2718
2783      2719
2784      2720
2785      2721
2786      2722
2787      2723
2788      2724
2789      2725
2790      2726
2791      2727
2792      2728
2793      2729
2794      2730
2795      2731
2796      2732
2797      2733
2798      2734
2799      2735
2800      2736
2801      2737
2802      2738
2803      2739
2804      2740
2805      2741
2806      2742
2807      2743
2808      2744
2809      2745
2810      2746

```

```

C CALLS SUBROUTINE LABOR TO FIND HOURS OF LABOR          2811
C CALLS SUBROUTINE GOODS TO FIND MATERIAL UNITS        2812
C CALLS SUBROUTINE COST TO FIND COST OF STAGE         2813
C UPDATES STAGE TO S+1                                 2814
C CALLS SUBROUTINE WRT$5 -PRINT BY MONTH BATCH BEGUN   2815
C           UPDATE2S MONTH TO M+1
C           -ALLOWS PROGRAM TO BE RERUN WITH A FEW DATA CHANGES
C
C ???????????????????????????????????????????????????
C PRICE OF ALGAE SUBROUTINES - GENERAL INFORMATION    2816
C
C NAME      ASSOCIATED FILES USED IN PRICES
C ICN5$      FOR INCLUSION OF COMMON BLOCKS USED IN PRICES
C
C NAME      DESCRIPTION OF ASSOC. SUBROUTINES, SPECIFIC TO ORGANISM
C
C RIN5$      READS IN ALL DATA ASSOCIATED WITH PRICE OF INPUT 5
C ROUT$5     READS OUT ALL DATA READ IN IN RIN $5
C ELEC       CALCULATES AMOUNT OF ELECTRICITY USED
C FUEL       CALCULATES AMOUNT OF FUEL OIL USED
C LABOR      CALCULATES QUANTITY OF LABOR USED
C GOODS      CALCULATES QUANTITY OF MATERIALS USED
C COST       COMPUTES COST OF INPUTS USED FOR ALGAE PRODUCTION
C WRT$5     WRITES OUT PRICE OF ALGAE COMPUTATIONS
C
C NAME      COMMON DIM      DESCRIPTION OF COMMON BLOCK VARIABLE
C
C ALFOU R   CN5$      MAXT     AVERAGE LITRES USEABLE BY ANIMALS
C ALLAB R   CN5$      MAXT     TOTAL LABOR PER STAGE, LESS INNOCLATION
C BEGPRO I   CN5$      MAXT     NUMBER OF MONTH IN WHICH PRODUCTION BEGINS
C BBLIF R   CN5$      MAXT     AVERAGE LIFE TIME OF A LIGHTBULB
C BLBRM R   CN5$      MAXT     NUMBER OF LIGHTBULBS IN ROOM OF STAGE
C CALIN R   CN5$      MAXT     INPUT COST, I, IN STAGE, LESS INNOCLATION
C CINP R    CN5$      4       INPUT COST, I, PER UNIT IN STAGE, LESS INOC2851
C COMEFF R   CN5$      MAXT     FRACTION EFFICIENCY OF AIR COMPRESSOR
C CPERM R   CN5$      MAXT     AVERAGE DENSITY OF MATURE CULTURE PER STAGE2854
C DATE2 C*25 CN5$      MAXT     DATE OF ALGAE PRICE DATA
C DAYSTG R  CN5$      MAXT     DAYS REQUIRED TO COMPLETE A STAGE
C DISSTG R  CN5$      MAXT     AVERAGE UNAVOIDABLE DISCARD PER STAGE
C ENDPRO I   CN5$      MAXT     MONTH IN WHICH PRODUCTION ENDS
C HPCOM R   CN5$      MAXT     AIR COMPRESSOR INPUT POWER RATING
C HPMP R    CN5$      MAXT     PUMP INPUT POWER RATING
C HRLABS R  CN5$      6       LABOR DEVOTED TO PRODUCTION OF A STAGE
C
C HRLABW R  CN5$      6       TYPE L LABOR USE PER WEEK
C HRSAIR R  CN5$      1.2    TIME A/C IS ON PER DAY IN STAGE ROOM
C
C HRSAUT R  CN5$      MAXT    TIME AUTOCLAVE IS ON PER STAGE
C HRSBLB R  CN5$      MAXT    TIME LIGHTS ARE ON IN STAGE ROOM
C HRSCom R  CN5$      MAXT    TIME COMPRESSOR IS ON PER DAY
C H20INK R  CN5$      MAXT    WATER ADDED AT BEGINNING OF EACH STAGE
C INOTNK R  CN5$      MAXT    TANK. VOLUME FOR EACH TANK EACH STAGE
C LFODD R   CN5$      MAXT    ALGAE AVAILABLE FOR FOOD
C LITWHR R  CN5$      MAXT    ELECTRICAL ENERGY FOR LIGHTING/STAGE
C MAXLAB I   CN5$      MAXT    MAXIMUM NUMBER OF LABOR ALLOWED. PAR=6
C MONAVA R  CN5$      MAXT    MONTH THAT EACH STAGE IS AVAILABLE TO USE
C
C NAME      MAXT     NAME OF EACH ALGAE STAGE
C NUMAIR R  CN5$      MAXT    NO. OF AIRCOND. IN ROOM OR EACH STAGE
C NUMGOD I   CN5$      MAXT    NO. OF DIFFERENT GOOD TYPES, G
C NUMLAB I   CN5$      MAXT    NO. OF DIFFERENT LABOR TYPES, I
C NUMSTG I   CN5$      MAXT    NO. OF STAGES TO PRODUCE A SINGLE BATCH
C NUMTnk R   CN5$      MAXT    NUMBER OF TANKS PER STAGE
C NUTLTr R  CN5$      MAXT    NUMBER OF TANKS FED BY COMPRESSOR
C NUTRT$5 C*1 CN5$      MAXT    NUTRIENT SOLUTION ADDED PER 1 OF H2O/STG
C PGOOD R   CN5$      MAXT    PRICE OF GOOD, TYPE G
C PKWHR R   CN5$      MAXT    PRICE PER KWHR OF ELECTRICITY
C PLABOR R  CN5$      6       PRICE PER HOUR OF TYPE L LABOR
C PMPEFF R  CN5$      MAXT    FRACTION EFFICIENCY RATING OF WATER PUMP
C PMPGPM R  CN5$      MAXT    OUTPUT CAPACITY RATING OF WATER PUMP
C POIL OIL   CN5$      MAXT    PRICE PER LITRE OF FUEL OIL
C QWRT$5 C*1 CN5$      MAXT    WRITE OUT OUTPUT FOR PRICES CALCULATIONS?
C STGRM R   CN5$      MAXT    NUMBER OF STAGES IN THE ROOM
C TALLAB R  CN5$      MAXT    TOTAL TYPE L LABOR PER STAGE, PLUS INNOC.
C TALLAL R  CN5$      MAXT    TOTAL TYPE L LABOR PER STAGE/LTR, + INNOC
C TCALIN R  CN5$      MAXT    COST OF ALL INPUT, I, PER STAGE, + INNOC.
C TCCELL R  CN5$      MAXT    TOTAL CELLS OF ALGAE PRODUCED
C TCINP R   CN5$      4       COST OF INPUT, I, PER STAGE, PLUS INNOC.
C
C TCCELL R  CN5$      MAXT    TOTAL COST PER CELL OF ALGAE PRODUCED
C TCINPL R  CN5$      MAXT    COST OF INPUT, I, PER STAGE, + INNOCULATION2898
C TICLTY R   CN5$      MAXT    COST OF TYPE L LABOR PER STAGE, + INNOC.
C TCLCTR R  CN5$      MAXT    COST OF ALL INPUT, I, PER LITRE PER STAGE
C TGOTY R   CN5$      MAXT    UNITS OF TYPE G GOOD USED PER STAGE
C TKWHR R   CN5$      MAXT    ELECTRICITY USED PER STAGE, LESS INNOCULA.
C TLABTY R  CN5$      MAXT    TYPE L LABOR USED PER STAGE, LESS INNOCULA
C
C TLITRE R  CN5$      MAXT    ALGAE PRODUCED IN EACH STAGE
C TOIL R    CN5$      MAXT    OIL CONSUMPTION PER STAGE, LESS INNOCULATIO2908
C TTGOTY R  CN5$      MAXT    TYPE G GOOD USED PER STAGE, PLUS INNOC.
C
C TTKEWL R  CN5$      MAXT    ELG. PER LITRE PER STAGE, PLUS INNOCULATION2910
C TTKHMR R  CN5$      MAXT    ELEC USED PER STAGE, PLUS INNOC.
C TTLYR R   CN5$      MAXT    LABOR, TYPE LE, PER STAGE, PLUS INNOC.
C
C TTOL R    CN5$      MAXT    OIL CONSUMED PER STAGE, PLUS INNOCULATION
C TTOLL R   CN5$      MAXT    OIL CONSUMED PER LITRE OF ALGAE
C TUBTNK R  CN5$      MAXT    COMPRESSOR TUBES PER TANK PER STAGE
C WATAIR R  CN5$      MAXT    AIR COMPRESSOR INPUT RATING
C WATAUT R  CN5$      MAXT    AUTOCLAVE INPUT WATT RATING
C WATBUL R  CN5$      MAXT    LIGHTBULB INPUT RATING
C
C ???????????????????????????????????????????????????
C
C 1. COMMON BLOCK VARIABLES
C INCLUDE (IPAR1)
C INCLUDE (ICN5$)
C INCLUDE (ICUNIT)
C INCLUDE (ICBLK1)
C INCLUDE (ICORG)
C INCLUDE (ICBLKO)
C INCLUDE (ICMAIN)
C 2. ARGUMENT VARIABLES
C
C      INTEGER STAGE, TIME,
C      WRT1, WRT2, RED, RED2, HREAD,
C      LASSTG
C      LOGICAL RUN2, RUN3
C      CHARACTER*9 MONBEG
C

```

```

C 3. LOCAL VARIABLES
  INTEGER ID, ITDAY, M, MSTART, MSTOP, S
  REAL RTDAY, RTDAY
  CHARACTER*1 ANSWR3, QUEST

C NAME  DESCRIPTION OF LOCAL VARIABLES AND ARGUMENTS
C ----- -----
C ANSWR3 OPTION TO REDO WITH A FEW CHANGES
C HREAD ARGUE: IOSTAT TO HELP INDICATE PROBLEMS IN READING FILE
C ID ADDITION TO MONTH TO UPDATE FOR CURRENT STAGE
C ITDAY INTEGER TO MONTH TO UPDATE FOR CURRENT STAGE
C LASSTG ARGUE: LAST STAGE OF ALGAE CALCULATED IN THE BATCH
C MONBEG ARGUE: MONTH IN WHICH STAGE=1 OF ALGAE IS BEGUN
C MSTART FIRST MONTH OF ALGAE PRODUCTION TO LOOP THROUGH
C MSTOP LAST MONTH OF ALGAE PRODUCTION TO LOOP THROUGH
C RED ARGUE: TRANSLATE UNIT READ FROM ORGANISM TO NEXT LEVEL SUBROUTINE
C RED2 ARGUE: TRANSLATE UNIT READ TO NEXT LEVEL SUBROUTINE
C RTDAY CONVENT TOTAL DAYS ELAISSED TO REAL
C RUN2 ARGUE: INDICATES THAT ONE RUN THRU ALGAE PRICE ALREADY DONE
C RUN3 ARGUE: INDICATES THAT 1 RUN THRU FROM ABOVE LEVEL SUBR. DONE
C S LOOP COUNTER FOR STAGE
C STAGE ARGUE: CURRENT STAGE OF ALGAE THAT CALCULATIONS ARE FOR
C TODAY TOTAL ELAPSED DAYS
C TIME ARGUE: CURRENT MONTH THAT CURRENT STAGE IS IN
C QUEST OPTION TO SEE LAST OUTPUT ON SCREEN
C WRT ARGUE: TRANSLATE UNIT WRITE FROM ORGANISM TO NEXT LEVEL SUBR.
C WRT2 ARGUE: UNIT TO WRITE TO NEXT LEVEL SUBROUTINE
C ----- -----
C TRANSLATE ARGUMENTS
  WRT2 = WRT
  RED2 = RED

C CALL SUBROUTINE TO INPUT ALL DATA
  CALL RIN$$(WRT2,RED2,RUN2,RUN3)

C OFFER A SECOND INPUT ENTRY POINT TO ENABLE MINOR INPUT CHANGES
  ANSWR3 = 'X'
  10 IF (ANSWR3 .EQ. 'Y') CALL INPUT2(WRT2,RED2,RUN2,RUN3)

C OFFER AN OPTION TO PRINT OUTPUT FOR ALGAE PRICE
  WRITE(WI,400)
  READ(RI,1000) QWRT$5
  400 FORMAT(' ','DO YOU WANT A PRINTED COPY OF THE ALGAE PRICE ',2
  ,',COMPUTATIONS? Y OR N')
  2985 C
  2986 C
  2987 C
  2988 C
  2989 C
  2990 C
  2991 C
  2992 C
  2993 C
  2994 C
  2995 C
  2996 C
  2997 C
  2998 C
  2999 C
  3000 C
  3001 C
  3002 C
  3003 C
  3004 C
  3005 C
  3006 C
  3007 C
  3008 C
  3009 C
  3010 C
  3011 C
  3012 C
  3013 C
  3014 C
  3015 C
  3016 C
  3017 C
  3018 C
  3019 C
  3020 C
  3021 C
  3022 C
  3023 C
  3024 C
  3025 C
  3026 C
  3027 C
  3028 C
  3029 C
  3030 C
  3031 C
  3032 C
  3033 C
  3034 C
  3035 C
  3036 C
  3037 C
  3038 C
  3039 C
  3040 C
  3041 C
  3042 C
  3043 C
  3044 C
  3045 C
  3046 C
  3047 C
  3048 C
  3049 C
  3050 C
  3051 C
  3052 C
  3053 C
  3054 C
  3055 C
  3056 C
  3057 C
  3058 C
  3059 C
  3060 C
  3061 C
  3062 C
  3063 C
  3064 C
  3065 C
  3066 C

```

```

  DAY = 0.
  DO 30 S = 1, NUMSTG
    ACCUMULATE DAYS TO DETERMINE WHAT MONTH CURRENT STAGE IS
    BEING PRODUCED IN (FIRST CONVERT TO AN INTEGER)
    STAGE = S
    TDAY = DAYSTG(S) + TODAY
    RTDAY = TODAY + 0.9999999
    ID = (RTDAY - 1) / 30
    TIME = M + ID
    IF ((ENDPRO-BEGPRO), EQ. 11 .AND. TIME.GT.12) TIME = TIME - 12
    STOP CALCULATIONS IF STAGE CONTINUES INTO NON-PRODUCTION MONTH
    IF (TIME .GT. ENDPRO) THEN
      MONATA(S) = NOT AVAIL.
      TLITRE(S) = 0.
      LFODD(S) = 0.
      TCELL(S) = 0.
      IF (S.GT. 1) THEN
        LFODD(S-1) = TLITRE(S-1)
        TCELL(S-1) = CPERM(S-1) * 1000. * TLITRE(S-1)
      ENDIF
      ELSE
        MONTH THAT ALGAE IS AVAILABLE AND LAST STAGE OF PRODUCTION
        MONAV(A,S) = NAMEN(TIME)
        LASSTG = S
      ENDIF
    QUANTITIES PRODUCED AND AVAILABLE - CHECK FOR LAST STAGE
    TLITRE(S) = (H2OINTK(S)+INQINTK(S)) * NUMTNK(S) - DISSTG(S)
    TCELL(S) = CPERM(S) * 1000. * TLITRE(S)
    IF (S .EQ. NUMSIG) THEN
      LFODD(S) = TLITRE(S)
    ELSE
      LFODD(S) = TLITRE(S) - INQTNK(S+1) * NUMTNK(S+1)
    ENDIF
    CALCULATE INPUT QUANTITIES - BY CALLING SUBROUTINES
    CAL ELEC(STAGE,TIME)
    CAL FUEL(STAGE,TIME)
    CAL LABOR(STAGE,TIME)
    CAL GOODS(STAGE,TIME)
    CALCULATE INPUT COSTS
    CAL COST(STAGE,TIME)
    TRANSLATE PRICES VARIABLES INTO ORGNSM VARIABLS
    PRICET(5,S,TIME) = TCCEL(S)
    NSAVAL(S,TIME) = TCELL(S)
    ENDIF
    CONTINUE ('QWRT$5 .EQ. 'Y') CALL WRT$5(MONBEG,LASSTG,W0)
    IF ((QWRT$5 .EQ. 'Y') CALL WRT$5(MONBEG,LASSTG,W0)
      20 CONTINUE
    C NAME OF THE MONTH IN WHICH PRODUCTION BEGINS
    C ALLOW PRINTING OF AT LEAST ONE STAGE
    MONBEG = NAMEM(N)
    LASSTG = NUMSTG
    MONTH = M
    Loop THROUGH EACH STAGE TO CALCULATE COSTS
  
```



2000 FORMAT(' /', 'ENTER DATA CONCERNING ALGAE PRODUCTION') 3195  
 2001 FORMAT(' ', '1. DATE OF COMPUTATIONS (TO 25 CHARACTERS IN QUOTES)') 3196  
 2002 FORMAT(' ', '2. PRICE PER KWHR OF ELECTRICITY') 3197  
 2003 FORMAT(' ', '2. PRICE PER LITRE OF FUEL OIL') 3198  
 2004 FORMAT(' ', '3. PRICE PER LITRE NUTRIENT SOLUTION') 3199  
 2005 FORMAT(' ', '4. PRICE PER LIGHTBULB') 3200  
 2006 FORMAT(' ', '5. NUMBER OF LABOR TYPES - USE AN INTEGER') 3201  
 2007 FORMAT(' ', '6. PRICE PER HOUR OF EACH LABOR TYPE') 3202  
 2008 FORMAT(' ', '2X, PRICE OF TYPE', 11) 3203  
 C LINE NUMBER 7-19  
 08 IF (INFO( 8) .EQ. 0) THEN  
   WRITE(W,208)  
   READ(R,\* ,ERR= 8) WATAIR  
 ENDIF  
 09 IF (INFO( 9) .EQ. 0) THEN  
   WRITE(W,209)  
   READ(R,\* ,ERR= 9) WATBLB  
 ENDIF  
 10 IF (INFO(10) .EQ. 0) THEN  
   WRITE(W,210)  
   READ(R,\* ,ERR=10) BLBLIF  
 ENDIF  
 11 IF (INFO(11) .EQ. 0) THEN  
   WRITE(W,211)  
   READ(R,\* ,ERR=11) WATAUT  
 ENDIF  
 12 IF (INFO(12) .EQ. 0) THEN  
   WRITE(W,212)  
   READ(R,\* ,ERR=12) HPCOM  
 ENDIF  
 13 IF (INFO(13) .EQ. 0) THEN  
   WRITE(W,213)  
   READ(R,\* ,ERR=13) HRSCom  
 ENDIF  
 14 IF (INFO(14) .EQ. 0) THEN  
   WRITE(W,214)  
   READ(R,\* ,ERR=14) NUMTUB  
 ENDIF  
 15 IF (INFO(15) .EQ. 0) THEN  
   WRITE(W,215)  
   READ(R,\* ,ERR=15) COMEFF  
 ENDIF  
 16 IF (INFO(16) .EQ. 0) THEN  
   WRITE(W,216)  
   READ(R,\* ,ERR=16) PMPEFF  
 ENDIF  
 17 IF (INFO(17) .EQ. 0) THEN  
   WRITE(W,217)  
   READ(R,\* ,ERR=17) HPPMP  
 ENDIF  
 18 IF (INFO(18) .EQ. 0) THEN  
   WRITE(W,218)  
   READ(R,\* ,ERR=18) HPLABW(L)  
 ENDIF  
 19 IF (INFO(19) .EQ. 0) THEN  
   WRITE(W,2219)  
   DO 81 L=1,NUMLAB  
     WRITE(W,219) L  
     READ(R,\* ,ERR=19) HRLABW(L)  
   CONTINUE  
 ENDIF  
 208 FORMAT (' ', '8. WATT RATING OF THE AIR CONDITIONER')  
 209 FORMAT (' ', '9. AVERAGE WATT RATING OF LIGHTBULBS') 3259  
 210 FORMAT (' ', '10. AVERAGE LIFETIME OF LIGHTBULB-IN HOURS') 3260  
 211 FORMAT (' ', '11. WATT RATING OF AUTOCLAVE OR STERILIZER') 3261  
 212 FORMAT (' ', '12. HORSEPOWER RATING OF AIR COMPRESSOR') 3262  
 213 FORMAT (' ', '13. HOURS PER DAY THAT COMPRESSOR OPERATES') 3263  
 214 FORMAT (' ', '14. TOTAL NUMBER OF AIR COMPRESSOR TUBES') 3264  
 215 FORMAT (' ', '15. EFFICIENCY RATING OF AIR COMPRESSOR') 3265  
 216 FORMAT (' ', '16. EFFICIENCY RATING OF WATER PUMP') 3266  
 217 FORMAT (' ', '17. GALLONS/MINUTE RATING OF WATER PUMP') 3267  
 218 FORMAT (' ', '18. HORSEPOWER RATING OF WATER PUMP') 3268  
 219 FORMAT (' ', '19. HOURS PER WEEK OF EACH LABOR TYPE') 3269  
 3205  
 3206  
 3207  
 3208  
 3209  
 3210  
 3211  
 3212  
 3213  
 3214  
 3215  
 3216  
 3217  
 3218  
 3219  
 3220  
 3221  
 3222  
 3223  
 3224  
 3225  
 3226  
 3227  
 3228  
 3229  
 3230  
 3231  
 3232  
 3233  
 3234  
 3235  
 3236  
 3237  
 3238  
 3239  
 3240  
 3241  
 3242  
 3243  
 3244  
 3245  
 3246  
 3247  
 3248  
 3249  
 3250  
 3251  
 3252  
 3253  
 3254  
 3255  
 3256  
 3257  
 3258  
 C LINES 20-35  
 20 DO 4000 K=20,35  
 4000 CONTINUE  
 IF (INFO(20) .EQ. 0) THEN  
   WRITE(W,220) MAXT  
   READ(R,\* ,ERR=20) NUMSTG  
 ENDIF  
 IF (RETYP1 .EQ. 0) THEN  
   WRITE(W,2222)  
   WRITE(W,2223) (S, S=1,NUMSTG)  
 ENDIF  
 IF (INFO(21) .EQ. 0) THEN  
   WRITE(W,221)  
   READ(R,\* ,ERR=21) (NUTLIR(S), S=1,NUMSTG)  
 ENDIF  
 IF (INFO(22) .EQ. 0) THEN  
   WRITE(W,222)  
   READ(R,\* ,ERR=22) (STGRM (S), S=1,NUMSTG)  
 ENDIF  
 IF (INFO(23) .EQ. 0) THEN  
   WRITE(W,223)  
   READ(R,\* ,ERR=23) (H2OTNK (S), S=1,NUMSTG)  
 ENDIF  
 IF (INFO(24) .EQ. 0) THEN  
   WRITE(W,224)  
   READ(R,\* ,ERR=24) (DAYSTG (S), S=1,NUMSTG)  
 ENDIF  
 IF (INFO(25) .EQ. 0) THEN  
   WRITE(W,225)  
   READ(R,\* ,ERR=25) (OPERM (S), S=1,NUMSTG)  
 ENDIF  
 IF (INFO(26) .EQ. 0) THEN  
   WRITE(W,226)  
   READ(R,\* ,ERR=26) (INQTNK (S), S=1,NUMSTG)  
 ENDIF  
 IF (INFO(27) .EQ. 0) THEN  
   WRITE(W,227)  
   READ(R,\* ,ERR=27) (DISSTG (S), S=1,NUMSTG)  
 ENDIF  
 IF (INFO(28) .EQ. 0) THEN  
   WRITE(W,228)  
   READ(R,\* ,ERR=28) (H2OTNK (S), S=1,NUMSTG)  
 ENDIF  
 IF (INFO(29) .EQ. 0) THEN  
   WRITE(W,229)  
   READ(R,\* ,ERR=29) (TUBTNK (S), S=1,NUMSTG)  
 ENDIF  
 IF (INFO(30) .EQ. 0) THEN  
   WRITE(W,230)  
   READ(R,\* ,ERR=30) (BLBRM (S), S=1,NUMSTG)  
 ENDIF

```

3387      READ(R,* ,ERR=38) (HRSAIR(M,S) , S=1,NUMSTG)
3388      CONTINUE
3389
3390      ENDIF
3391
3392      IF (INFO(31) .EQ. 0) THEN
3393          WRITE(W,231)
3394          READ(R,* ,ERR=31) (HRSBLB(S) , S=1,NUMSTG)
3395
3396      ENDIF
3397      IF (INFO(32) .EQ. 0) THEN
3398          WRITE(W,232)
3399          READ(R,* ,ERR=32) (HRSAUT(S) , S=1,NUMSTG)
3400
3401      ENDIF
3402      IF (INFO(33) .EQ. 0) THEN
3403          WRITE(W,233)
3404          READ(R,* ,ERR=33) (NUMATR(S) , S=1,NUMSTG)
3405
3406      ENDIF
3407      IF (INFO(34) .EQ. 0) THEN
3408          WRITE(W,2234)
3409          DO 88 L=1,NUMLAB
3410              WRITE(W,234) L,
3411              READ(R,* ,ERR=34) (HRLABS(L,S) , S=1,NUMSTG)
3412
3413      CONTINUE
3414
3415      IF (INFO(35) .EQ. 0) THEN
3416          WRITE(W,235)
3417          READ(R,* ,ERR=35) (NAMALG(S) , S=1,NUMSTG)
3418
3419      ENDIF
3420
3421      FORMAT('   ',1X,6(I2,'STAGE',2X) / ' ',1X,6(I2,'STAGE',2X))
3422      FORMAT('   ',21. LITRES NUTRIENT SOL./LITRE H2O')
3423      FORMAT('   ',22. # OF TANKS')
3424      FORMAT('   ',23. # OF STAGES IN ROOM')
3425      FORMAT('   ',24. DAYS TO MATURITY')
3426      FORMAT('   ',25. CELLS/MILLIRE')
3427      FORMAT('   ',26. L. H2O ADDED/TANK')
3428      FORMAT('   ',27. L. INNOVOLA/TANK')
3429      FORMAT('   ',28. L. DISCARD)
3430      FORMAT('   ',29. # TUBES/TANK)
3431      FORMAT('   ',30. # LITEBULB/ROOM')
3432      FORMAT('   ',31. HRS LITES ON/DAY/RM')
3433      FORMAT('   ',32. HRS AUTOCLAVE )
3434      FORMAT('   ',33. # AC.S./ROOM')
3435      FORMAT('   ',34. HRS LABOR/STAGE')
3436      FORMAT('   ',35. NAME OF EACH ALGAE STAGE' / ' ,5X,
3437      2       UP TO 10 CHARACTERS IN QUOTES, )
3438
3439      C LINES 35-41
3440      DO 4001 K=36,41
3441          IF (INFO(K) .EQ. 0) RETYP2=0
3442
3443      4001  CONTINUE
3444          IF (INFO(36) .EQ. 0) THEN
3445              WRITE(W,236)
3446              READ(R,* ,ERR=36) BEGPRO
3447
3448      37      IF (INFO(37) .EQ. 0) THEN
3449              WRITE(W,237)
3450              READ(R,* ,ERR=37) ENDPRO
3451
3452          IF (INFO(38) .EQ. 0) WRITE(W,2237) (S, S=1,NUMSTG)
3453
3454          IF (INFO(38) .EQ. 0) THEN
3455              WRITE(W,2238)
3456              DO 82 M=BEGPRO,ENDPRO
3457                  WRITE(W,238) M,NAME(M)
3458
3459      38      C LINES 35-41
3460      IF (INFO(36) .EQ. 0) THEN
3461          WRITE(W,236)
3462          READ(R,* ,ERR=36) BEGPRO
3463
3464          IF (INFO(38) .EQ. 0) THEN
3465              WRITE(W,2238)
3466              DO 82 M=BEGPRO,ENDPRO
3467                  WRITE(W,238) M,NAME(M)
3468
3469      39      C LINES 35-41
3470      IF (INFO(36) .EQ. 0) THEN
3471          WRITE(W,236)
3472          READ(R,* ,ERR=36) BEGPRO
3473
3474          IF (INFO(38) .EQ. 0) THEN
3475              WRITE(W,2238)
3476              DO 82 M=BEGPRO,ENDPRO
3477                  WRITE(W,238) M,NAME(M)
3478
3479      40      C LINES 35-41
3480      IF (INFO(36) .EQ. 0) THEN
3481          WRITE(W,236)
3482          READ(R,* ,ERR=36) BEGPRO
3483
3484          IF (INFO(38) .EQ. 0) THEN
3485              WRITE(W,2238)
3486              DO 82 M=BEGPRO,ENDPRO
3487                  WRITE(W,238) M,NAME(M)
3488
3489
3490      41      C LINES 35-41
3491      IF (INFO(36) .EQ. 0) THEN
3492          WRITE(W,236)
3493          READ(R,* ,ERR=36) BEGPRO
3494
3495          IF (INFO(38) .EQ. 0) THEN
3496              WRITE(W,2238)
3497              DO 82 M=BEGPRO,ENDPRO
3498                  WRITE(W,238) M,NAME(M)
3499
3500
3501      42      C LINES 35-41
3502      IF (INFO(36) .EQ. 0) THEN
3503          WRITE(W,236)
3504          READ(R,* ,ERR=36) BEGPRO
3505
3506          IF (INFO(38) .EQ. 0) THEN
3507              WRITE(W,2238)
3508              DO 82 M=BEGPRO,ENDPRO
3509                  WRITE(W,238) M,NAME(M)
3510
3511
3512      43      C LINES 35-41
3513      IF (INFO(36) .EQ. 0) THEN
3514          WRITE(W,236)
3515          READ(R,* ,ERR=36) BEGPRO
3516
3517          IF (INFO(38) .EQ. 0) THEN
3518              WRITE(W,2238)
3519              DO 82 M=BEGPRO,ENDPRO
3520                  WRITE(W,238) M,NAME(M)
3521
3522
3523      44      C LINES 35-41
3524      IF (INFO(36) .EQ. 0) THEN
3525          WRITE(W,236)
3526          READ(R,* ,ERR=36) BEGPRO
3527
3528          IF (INFO(38) .EQ. 0) THEN
3529              WRITE(W,2238)
3530              DO 82 M=BEGPRO,ENDPRO
3531                  WRITE(W,238) M,NAME(M)
3532
3533
3534      45      C LINES 35-41
3535      IF (INFO(36) .EQ. 0) THEN
3536          WRITE(W,236)
3537          READ(R,* ,ERR=36) BEGPRO
3538
3539          IF (INFO(38) .EQ. 0) THEN
3540              WRITE(W,2238)
3541              DO 82 M=BEGPRO,ENDPRO
3542                  WRITE(W,238) M,NAME(M)
3543
3544
3545      46      C LINES 35-41
3546      IF (INFO(36) .EQ. 0) THEN
3547          WRITE(W,236)
3548          READ(R,* ,ERR=36) BEGPRO
3549
3550          IF (INFO(38) .EQ. 0) THEN
3551              WRITE(W,2238)
3552              DO 82 M=BEGPRO,ENDPRO
3553                  WRITE(W,238) M,NAME(M)
3554
3555
3556      47      C LINES 35-41
3557      IF (INFO(36) .EQ. 0) THEN
3558          WRITE(W,236)
3559          READ(R,* ,ERR=36) BEGPRO
3560
3561          IF (INFO(38) .EQ. 0) THEN
3562              WRITE(W,2238)
3563              DO 82 M=BEGPRO,ENDPRO
3564                  WRITE(W,238) M,NAME(M)
3565
3566
3567      48      C LINES 35-41
3568      IF (INFO(36) .EQ. 0) THEN
3569          WRITE(W,236)
3570          READ(R,* ,ERR=36) BEGPRO
3571
3572          IF (INFO(38) .EQ. 0) THEN
3573              WRITE(W,2238)
3574              DO 82 M=BEGPRO,ENDPRO
3575                  WRITE(W,238) M,NAME(M)
3576
3577
3578      49      C LINES 35-41
3579      IF (INFO(36) .EQ. 0) THEN
3580          WRITE(W,236)
3581          READ(R,* ,ERR=36) BEGPRO
3582
3583          IF (INFO(38) .EQ. 0) THEN
3584              WRITE(W,2238)
3585              DO 82 M=BEGPRO,ENDPRO
3586                  WRITE(W,238) M,NAME(M)
3587
3588
3589      50      C LINES 35-41
3590      IF (INFO(36) .EQ. 0) THEN
3591          WRITE(W,236)
3592          READ(R,* ,ERR=36) BEGPRO
3593
3594          IF (INFO(38) .EQ. 0) THEN
3595              WRITE(W,2238)
3596              DO 82 M=BEGPRO,ENDPRO
3597                  WRITE(W,238) M,NAME(M)
3598
3599
3600      51      C LINES 35-41
3601      IF (INFO(36) .EQ. 0) THEN
3602          WRITE(W,236)
3603          READ(R,* ,ERR=36) BEGPRO
3604
3605          IF (INFO(38) .EQ. 0) THEN
3606              WRITE(W,2238)
3607              DO 82 M=BEGPRO,ENDPRO
3608                  WRITE(W,238) M,NAME(M)
3609
3610
3611      52      C LINES 35-41
3612      IF (INFO(36) .EQ. 0) THEN
3613          WRITE(W,236)
3614          READ(R,* ,ERR=36) BEGPRO
3615
3616          IF (INFO(38) .EQ. 0) THEN
3617              WRITE(W,2238)
3618              DO 82 M=BEGPRO,ENDPRO
3619                  WRITE(W,238) M,NAME(M)
3620
3621
3622      53      C LINES 35-41
3623      IF (INFO(36) .EQ. 0) THEN
3624          WRITE(W,236)
3625          READ(R,* ,ERR=36) BEGPRO
3626
3627          IF (INFO(38) .EQ. 0) THEN
3628              WRITE(W,2238)
3629              DO 82 M=BEGPRO,ENDPRO
3630                  WRITE(W,238) M,NAME(M)
3631
3632
3633      54      C LINES 35-41
3634      IF (INFO(36) .EQ. 0) THEN
3635          WRITE(W,236)
3636          READ(R,* ,ERR=36) BEGPRO
3637
3638          IF (INFO(38) .EQ. 0) THEN
3639              WRITE(W,2238)
3640              DO 82 M=BEGPRO,ENDPRO
3641                  WRITE(W,238) M,NAME(M)
3642
3643
3644      55      C LINES 35-41
3645      IF (INFO(36) .EQ. 0) THEN
3646          WRITE(W,236)
3647          READ(R,* ,ERR=36) BEGPRO
3648
3649          IF (INFO(38) .EQ. 0) THEN
3650              WRITE(W,2238)
3651              DO 82 M=BEGPRO,ENDPRO
3652                  WRITE(W,238) M,NAME(M)
3653
3654
3655      56      C LINES 35-41
3656      IF (INFO(36) .EQ. 0) THEN
3657          WRITE(W,236)
3658          READ(R,* ,ERR=36) BEGPRO
3659
3660          IF (INFO(38) .EQ. 0) THEN
3661              WRITE(W,2238)
3662              DO 82 M=BEGPRO,ENDPRO
3663                  WRITE(W,238) M,NAME(M)
3664
3665
3666      57      C LINES 35-41
3667      IF (INFO(36) .EQ. 0) THEN
3668          WRITE(W,236)
3669          READ(R,* ,ERR=36) BEGPRO
3670
3671          IF (INFO(38) .EQ. 0) THEN
3672              WRITE(W,2238)
3673              DO 82 M=BEGPRO,ENDPRO
3674                  WRITE(W,238) M,NAME(M)
3675
3676
3677      58      C LINES 35-41
3678      IF (INFO(36) .EQ. 0) THEN
3679          WRITE(W,236)
3680          READ(R,* ,ERR=36) BEGPRO
3681
3682          IF (INFO(38) .EQ. 0) THEN
3683              WRITE(W,2238)
3684              DO 82 M=BEGPRO,ENDPRO
3685                  WRITE(W,238) M,NAME(M)
3686
3687
3688      59      C LINES 35-41
3689      IF (INFO(36) .EQ. 0) THEN
3690          WRITE(W,236)
3691          READ(R,* ,ERR=36) BEGPRO
3692
3693          IF (INFO(38) .EQ. 0) THEN
3694              WRITE(W,2238)
3695              DO 82 M=BEGPRO,ENDPRO
3696                  WRITE(W,238) M,NAME(M)
3697
3698
3699      60      C LINES 35-41
3700      IF (INFO(36) .EQ. 0) THEN
3701          WRITE(W,236)
3702          READ(R,* ,ERR=36) BEGPRO
3703
3704          IF (INFO(38) .EQ. 0) THEN
3705              WRITE(W,2238)
3706              DO 82 M=BEGPRO,ENDPRO
3707                  WRITE(W,238) M,NAME(M)
3708
3709
3710      61      C LINES 35-41
3711      IF (INFO(36) .EQ. 0) THEN
3712          WRITE(W,236)
3713          READ(R,* ,ERR=36) BEGPRO
3714
3715          IF (INFO(38) .EQ. 0) THEN
3716              WRITE(W,2238)
3717              DO 82 M=BEGPRO,ENDPRO
3718                  WRITE(W,238) M,NAME(M)
3719
3720
3721      62      C LINES 35-41
3722      IF (INFO(36) .EQ. 0) THEN
3723          WRITE(W,236)
3724          READ(R,* ,ERR=36) BEGPRO
3725
3726          IF (INFO(38) .EQ. 0) THEN
3727              WRITE(W,2238)
3728              DO 82 M=BEGPRO,ENDPRO
3729                  WRITE(W,238) M,NAME(M)
3730
3731
3732      63      C LINES 35-41
3733      IF (INFO(36) .EQ. 0) THEN
3734          WRITE(W,236)
3735          READ(R,* ,ERR=36) BEGPRO
3736
3737          IF (INFO(38) .EQ. 0) THEN
3738              WRITE(W,2238)
3739              DO 82 M=BEGPRO,ENDPRO
3740                  WRITE(W,238) M,NAME(M)
3741
3742
3743      64      C LINES 35-41
3744      IF (INFO(36) .EQ. 0) THEN
3745          WRITE(W,236)
3746          READ(R,* ,ERR=36) BEGPRO
3747
3748          IF (INFO(38) .EQ. 0) THEN
3749              WRITE(W,2238)
3750              DO 82 M=BEGPRO,ENDPRO
3751                  WRITE(W,238) M,NAME(M)
3752
3753
3754      65      C LINES 35-41
3755      IF (INFO(36) .EQ. 0) THEN
3756          WRITE(W,236)
3757          READ(R,* ,ERR=36) BEGPRO
3758
3759          IF (INFO(38) .EQ. 0) THEN
3760              WRITE(W,2238)
3761              DO 82 M=BEGPRO,ENDPRO
3762                  WRITE(W,238) M,NAME(M)
3763
3764
3765      66      C LINES 35-41
3766      IF (INFO(36) .EQ. 0) THEN
3767          WRITE(W,236)
3768          READ(R,* ,ERR=36) BEGPRO
3769
3770          IF (INFO(38) .EQ. 0) THEN
3771              WRITE(W,2238)
3772              DO 82 M=BEGPRO,ENDPRO
3773                  WRITE(W,238) M,NAME(M)
3774
3775
3776      67      C LINES 35-41
3777      IF (INFO(36) .EQ. 0) THEN
3778          WRITE(W,236)
3779          READ(R,* ,ERR=36) BEGPRO
3780
3781          IF (INFO(38) .EQ. 0) THEN
3782              WRITE(W,2238)
3783              DO 82 M=BEGPRO,ENDPRO
3784                  WRITE(W,238) M,NAME(M)
3785
3786
3787      68      C LINES 35-41
3788      IF (INFO(36) .EQ. 0) THEN
3789          WRITE(W,236)
3790          READ(R,* ,ERR=36) BEGPRO
3791
3792          IF (INFO(38) .EQ. 0) THEN
3793              WRITE(W,2238)
3794              DO 82 M=BEGPRO,ENDPRO
3795                  WRITE(W,238) M,NAME(M)
3796
3797
3798      69      C LINES 35-41
3799      IF (INFO(36) .EQ. 0) THEN
3800          WRITE(W,236)
3801          READ(R,* ,ERR=36) BEGPRO
3802
3803          IF (INFO(38) .EQ. 0) THEN
3804              WRITE(W,2238)
3805              DO 82 M=BEGPRO,ENDPRO
3806                  WRITE(W,238) M,NAME(M)
3807
3808
3809      70      C LINES 35-41
3810      IF (INFO(36) .EQ. 0) THEN
3811          WRITE(W,236)
3812          READ(R,* ,ERR=36) BEGPRO
3813
3814          IF (INFO(38) .EQ. 0) THEN
3815              WRITE(W,2238)
3816              DO 82 M=BEGPRO,ENDPRO
3817                  WRITE(W,238) M,NAME(M)
3818
3819
3820      71      C LINES 35-41
3821      IF (INFO(36) .EQ. 0) THEN
3822          WRITE(W,236)
3823          READ(R,* ,ERR=36) BEGPRO
3824
3825          IF (INFO(38) .EQ. 0) THEN
3826              WRITE(W,2238)
3827              DO 82 M=BEGPRO,ENDPRO
3828                  WRITE(W,238) M,NAME(M)
3829
3830
3831      72      C LINES 35-41
3832      IF (INFO(36) .EQ. 0) THEN
3833          WRITE(W,236)
3834          READ(R,* ,ERR=36) BEGPRO
3835
3836          IF (INFO(38) .EQ. 0) THEN
3837              WRITE(W,2238)
3838              DO 82 M=BEGPRO,ENDPRO
3839                  WRITE(W,238) M,NAME(M)
3840
3841
3842      73      C LINES 35-41
3843      IF (INFO(36) .EQ. 0) THEN
3844          WRITE(W,236)
3845          READ(R,* ,ERR=36) BEGPRO
3846
3847          IF (INFO(38) .EQ. 0) THEN
3848              WRITE(W,2238)
3849              DO 82 M=BEGPRO,ENDPRO
3850                  WRITE(W,238) M,NAME(M)
3851
3852
3853      74      C LINES 35-41
3854      IF (INFO(36) .EQ. 0) THEN
3855          WRITE(W,236)
3856          READ(R,* ,ERR=36) BEGPRO
3857
3858          IF (INFO(38) .EQ. 0) THEN
3859              WRITE(W,2238)
3860              DO 82 M=BEGPRO,ENDPRO
3861                  WRITE(W,238) M,NAME(M)
3862
3863
3864      75      C LINES 35-41
3865      IF (INFO(36) .EQ. 0) THEN
3866          WRITE(W,236)
3867          READ(R,* ,ERR=36) BEGPRO
3868
3869          IF (INFO(38) .EQ. 0) THEN
3870              WRITE(W,2238)
3871              DO 82 M=BEGPRO,ENDPRO
3872                  WRITE(W,238) M,NAME(M)
3873
3874
3875      76      C LINES 35-41
3876      IF (INFO(36) .EQ. 0) THEN
3877          WRITE(W,236)
3878          READ(R,* ,ERR=36) BEGPRO
3879
3880          IF (INFO(38) .EQ. 0) THEN
3881              WRITE(W,2238)
3882              DO 82 M=BEGPRO,ENDPRO
3883                  WRITE(W,238) M,NAME(M)
3884
3885
3886      77      C LINES 35-41
3887      IF (INFO(36) .EQ. 0) THEN
3888          WRITE(W,236)
3889          READ(R,* ,ERR=36) BEGPRO
3890
3891          IF (INFO(38) .EQ. 0) THEN
3892              WRITE(W,2238)
3893              DO 82 M=BEGPRO,ENDPRO
3894                  WRITE(W,238) M,NAME(M)
3895
3896
3897      78      C LINES 35-41
3898      IF (INFO(36) .EQ. 0) THEN
3899          WRITE(W,236)
3900          READ(R,* ,ERR=36) BEGPRO
3901
3902          IF (INFO(38) .EQ. 0) THEN
3903              WRITE(W,2238)
3904              DO 82 M=BEGPRO,ENDPRO
3905                  WRITE(W,238) M,NAME(M)
3906
3907
3908      79      C LINES 35-41
3909      IF (INFO(36) .EQ. 0) THEN
3910          WRITE(W,236)
3911          READ(R,* ,ERR=36) BEGPRO
3912
3913          IF (INFO(38) .EQ. 0) THEN
3914              WRITE(W,2238)
3915              DO 82 M=BEGPRO,ENDPRO
3916                  WRITE(W,238) M,NAME(M)
3917
3918
3919      80      C LINES 35-41
3920      IF (INFO(36) .EQ. 0) THEN
3921          WRITE(W,236)
3922          READ(R,* ,ERR=36) BEGPRO
3923
3924          IF (INFO(38) .EQ. 0) THEN
3925              WRITE(W,2238)
3926              DO 82 M=BEGPRO,ENDPRO
3927                  WRITE(W,238) M,NAME(M)
3928
3929
3930      81      C LINES 35-41
3931      IF (INFO(36) .EQ. 0) THEN
3932          WRITE(W,236)
3933          READ(R,* ,ERR=36) BEGPRO
3934
3935          IF (INFO(38) .EQ. 0) THEN
3936              WRITE(W,2238)
3937              DO 82 M=BEGPRO,ENDPRO
3938                  WRITE(W,238) M,NAME(M)
3939
3940
3941      82      C LINES 35-41
3942      IF (INFO(36) .EQ. 0) THEN
3943          WRITE(W,236)
3944          READ(R,* ,ERR=36) BEGPRO
3945
3946          IF (INFO(38) .EQ. 0) THEN
3947              WRITE(W,2238)
3948              DO 82 M=BEGPRO,ENDPRO
3949                  WRITE(W,238) M,NAME(M)
3950
3951
3952      83      C LINES 35-41
3953      IF (INFO(36) .EQ. 0) THEN
3954          WRITE(W,236)
3955          READ(R,* ,ERR=36) BEGPRO
3956
3957          IF (INFO(38) .EQ. 0) THEN
3958              WRITE(W,2238)
3959              DO 82 M=BEGPRO,ENDPRO
3960                  WRITE(W,238) M,NAME(M)
3961
3962
3963      84      C LINES 35-41
3964      IF (INFO(36) .EQ. 0) THEN
3965          WRITE(W,236)
3966          READ(R,* ,ERR=36) BEGPRO
3967
3968          IF (INFO(38) .EQ. 0) THEN
3969              WRITE(W,2238)
3970              DO 82 M=BEGPRO,ENDPRO
3971                  WRITE(W,238) M,NAME(M)
3972
3973
3974      85      C LINES 35-41
3975      IF (INFO(36) .EQ. 0) THEN
3976          WRITE(W,236)
3977          READ(R,* ,ERR=36) BEGPRO
3978
3979          IF (INFO(38) .EQ. 0) THEN
3980              WRITE(W,2238)
3981              DO 82 M=BEGPRO,ENDPRO
3982                  WRITE(W,238) M,NAME(M)
3983
3984
3985      86      C LINES 35-41
3986      IF (INFO(36) .EQ. 0) THEN
3987          WRITE(W,236)
3988          READ(R,* ,ERR=36) BEGPRO
3989
3990          IF (INFO(38) .EQ. 0) THEN
3991              WRITE(W,2238)
3992              DO 82 M=BEGPRO,ENDPRO
3993                  WRITE(W,238) M,NAME(M)
3994
3995
3996      87      C LINES 35-41
3997      IF (INFO(36) .EQ. 0) THEN
3998          WRITE(W,236)
3999          READ(R,* ,ERR=36) BEGPRO
4000
4001          IF (INFO(38) .EQ. 0) THEN
4002              WRITE(W,2238)
4003              DO 82 M=BEGPRO,ENDPRO
4004                  WRITE(W,238) M,NAME(M)
4005
4006
4007      88      C LINES 35-41
4008      IF (INFO(36) .EQ. 0) THEN
4009          WRITE(W,236)
4010          READ(R,* ,ERR=36) BEGPRO
4011
4012          IF (INFO(38) .EQ. 0) THEN
4013              WRITE(W,2238)
4014              DO 82 M=BEGPRO,ENDPRO
4015                  WRITE(W,238) M,NAME(M)
4016
4017
4018      89      C LINES 35-41
4019      IF (INFO(36) .EQ. 0) THEN
4020          WRITE(W,236)
4021          READ(R,* ,ERR=36) BEGPRO
4022
4023          IF (INFO(38) .EQ. 0) THEN
4024              WRITE(W,2238)
4025              DO 82 M=BEGPRO,ENDPRO
4026                  WRITE(W,238) M,NAME(M)
4027
4028
4029      90      C LINES 35-41
4030      IF (INFO(36) .EQ. 0) THEN
4031          WRITE(W,236)
4032          READ(R,* ,ERR=36) BEGPRO
4033
4034          IF (INFO(38) .EQ. 0) THEN
4035              WRITE(W,2238)
4036              DO 82 M=BEGPRO,ENDPRO
4037                  WRITE(W,238) M,NAME(M)
4038
4039
4040      91      C LINES 35-41
4041      IF (INFO(36) .EQ. 0) THEN
4042          WRITE(W,236)
4043          READ(R,* ,ERR=36) BEGPRO
4044
4045          IF (INFO(38) .EQ. 0) THEN
4046              WRITE(W,2238)
4047              DO 82 M=BEGPRO,ENDPRO
4048                  WRITE(W,238) M,NAME(M)
4049
4050
4051      92      C LINES 35-41
4052      IF (INFO(36) .EQ. 0) THEN
4053          WRITE(W,236)
4054          READ(R,* ,ERR=36) BEGPRO
4055
4056          IF (INFO(38) .EQ. 0) THEN
4057              WRITE(W,2238)
4058              DO 82 M=BEGPRO,ENDPRO
4059                  WRITE(W,238) M,NAME(M)
4060
4061
4062      93      C LINES 35-41
4063      IF (INFO(36) .EQ. 0) THEN
4064          WRITE(W,236)
4065          READ(R,* ,ERR=36) BEGPRO
4066
4067          IF (INFO(38) .EQ. 0) THEN
4068              WRITE(W,2238)
4069              DO 82 M=BEGPRO,ENDPRO
4070                  WRITE(W,238) M,NAME(M)
4071
4072
4073      94      C LINES 35-41
4074      IF (INFO(36) .EQ. 0) THEN
4075          WRITE(W,236)
4076          READ(R,* ,ERR=36) BEGPRO
4077
4078          IF (INFO(38) .EQ. 0) THEN
4079              WRITE(W,2238)
4080              DO 82 M=BEGPRO,ENDPRO
4081                  WRITE(W,238) M,NAME(M)
4082
4083
4084      95      C LINES 35-41
4085      IF (INFO(36) .EQ. 0) THEN
4086          WRITE(W,236)
4087          READ(R,* ,ERR=36) BEGPRO
4088
4089          IF (INFO(38) .EQ. 0) THEN
4090              WRITE(W,2238)
4091              DO 82 M=BEGPRO,ENDPRO
4092                  WRITE(W,238) M,NAME(M)
4093
4094
4095      96      C LINES 35-41
4096      IF (INFO(36) .EQ. 0) THEN
4097          WRITE(W,236)
4098          READ(R,* ,ERR=36) BEGPRO
4099
4100          IF (INFO(38) .EQ. 0) THEN
4101              WRITE(W,2238)
4102              DO 82 M=BEGPRO,ENDPRO
4103                  WRITE(W,238) M,NAME(M)
4104
4105
4106      97      C LINES 35-41
4107      IF (INFO(36) .EQ. 0) THEN
4108          WRITE(W,236)
4109          READ(R,* ,ERR=36) BEGPRO
4110
4111          IF (INFO(38) .EQ. 0) THEN
4112              WRITE(W,2238)
4113              DO 82 M=BEGPRO,ENDPRO
4114                  WRITE(W,238) M,NAME(M)
4115
4116
4117      98      C LINES 35-41
4118      IF (INFO(36) .EQ. 0) THEN
4119          WRITE(W,236)
4120          READ(R,* ,ERR=36) BEGPRO
4121
4122          IF (INFO(38) .EQ. 0) THEN
4123              WRITE(W,2238)
4124              DO 82 M=BEGPRO,ENDPRO
4125                  WRITE(W,238) M,NAME(M)
4126
4127
4128      99      C LINES 35-41
4129      IF (INFO(36) .EQ. 0) THEN
4130          WRITE(W,236)
4131          READ(R,* ,ERR=36) BEGPRO
4132
4133          IF (INFO(38) .EQ. 0) THEN
4134              WRITE(W,2238)
4135              DO 82 M=BEGPRO,ENDPRO
4136                  WRITE(W,238) M,NAME(M)
4137
4138
4139      100     C LINES 35-41
4140      IF (INFO(36) .EQ. 0) THEN
4141          WRITE(W,236)
4142          READ(R,* ,ERR=36) BEGPRO
4143
4144          IF (INFO(38) .EQ. 0) THEN
4145              WRITE(W,2238)
4146              DO 82 M=BEGPRO,ENDPRO
4147                  WRITE(W,238) M,NAME(M)
4148
4149
4150      101     C LINES 35-41
4151      IF (INFO(36) .EQ. 0) THEN
4152          WRITE(W,236)
4153          READ(R,* ,ERR=36) BEGPRO
4154
4155          IF (INFO(38) .EQ. 0) THEN
4156              WRITE(W,2238)
4157              DO 82 M=BEGPRO,ENDPRO
4158                  WRITE(W,238) M,NAME(M)
4159
4160
4161      102     C LINES 35-41
4162      IF (INFO(36) .EQ. 0) THEN
4163          WRITE(W,236)
4164          READ(R,* ,ERR=36) BEGPRO
4165
4166          IF (INFO(38) .EQ. 0) THEN
4167              WRITE(W,2238)
4168              DO 82 M=BEGPRO,ENDPRO
4169                  WRITE(W,238) M,NAME(M)
4170
4171
4172      103     C LINES 35-41
4173      IF (INFO(36) .EQ. 0) THEN
4174          WRITE(W,236)
4175          READ(R,* ,ERR=36) BEGPRO
4176
4177          IF (INFO(38) .EQ. 0) THEN
4178              WRITE(W,2238)
4179              DO 82 M=BEGPRO,ENDPRO
4180                  WRITE(W,238) M,NAME(M)
4181
4182
4183      104     C LINES 35-41
4184      IF (INFO(36) .EQ. 0) THEN
4185          WRITE(W,236)
4186          READ(R,* ,ERR=36) BEGPRO
4187
4188          IF (INFO(38) .EQ. 0) THEN
4189              WRITE(W,2238)
4190              DO 82 M=BEGPRO,ENDPRO
4191                  WRITE(W,238) M,NAME(M)
4192
4193
4194      105     C LINES 35-41
4195      IF (INFO(36) .EQ. 0) THEN
4196          WRITE(W,236)
4197          READ(R,* ,ERR=36) BEGPRO
4198
4199          IF (INFO(38) .EQ. 0) THEN
4200              WRITE(W,2238)
4201              DO 82 M=BEGPRO,ENDPRO
4202                  WRITE(W,238) M,NAME(M)
4203
4204
4205      106     C LINES 35-41
4206      IF (INFO(36) .EQ. 0) THEN
4207          WRITE(W,236)
4208          READ(R,* ,ERR=36) BEGPRO
4209
4210          IF (INFO(38) .EQ. 0) THEN
4211              WRITE(W,2238)
4212              DO 82 M=BEGPRO,ENDPRO
4213                  WRITE(W,238) M,NAME(M)
4214
4215
4216      107     C LINES 35-41
4217      IF (INFO(36) .EQ. 0) THEN
4218          WRITE(W,236)
4219          READ(R,* ,ERR=36) BEGPRO
4220
4221          IF (INFO(38) .EQ. 0) THEN
4222              WRITE(W,2238)
4223              DO 82 M=BEGPRO,ENDPRO
4224                  WRITE(W,238) M,NAME(M)
4225
4226
4227      108     C LINES 35-41
4228      IF (INFO(36) .EQ. 0) THEN
4229          WRITE(W,236)
4230          READ(R,* ,ERR=36) BEGPRO
4231
4232          IF (INFO(38) .EQ. 0) THEN
4233              WRITE(W,2238)
4234              DO 82 M=BEGPRO,ENDPRO
4235                  WRITE(W,238) M,NAME(M)
4236
4237
4238      109     C LINES 35-41
4239      IF (INFO(36) .EQ. 0) THEN
4240          WRITE(W,236)
4241          READ(R,* ,ERR=36) BEGPRO
4242
4243          IF (INFO(38) .EQ. 0) THEN
4244              WRITE(W,2238)
4245              DO 82 M=BEGPRO,ENDPRO
4246                  WRITE(W,238) M,NAME(M)
4247
4248
4249      110     C LINES 35-41
4250      IF (INFO(36) .EQ. 0) THEN
4251          WRITE(W,236)
4252          READ(R,* ,ERR=36) BEGPRO
4253
4254          IF (INFO(38) .EQ. 0) THEN
4255              WRITE(W,2238)
4256              DO 82 M=BEGPRO,ENDPRO
4257                  WRITE(W,238) M,NAME(M)
4258
4259
4260      111     C LINES 35-41
4261      IF (INFO(36) .EQ. 0) THEN
4262          WRITE(W,236)
4263          READ(R,* ,ERR=36) BEGPRO
4264
4265          IF (INFO(38) .EQ. 0) THEN
4266              WRITE(W,2238)
4267              DO 82 M=BEGPRO,ENDPRO
4268                  WRITE(W,238) M,NAME(M)
4269
4270
4271      112     C LINES 35-41
4272      IF (INFO(36) .EQ. 0) THEN
42
```

```

3515      PRINT2 = 2
3516      ENDIF
3517      IF (COPYA .EQ. 'Y' .AND. COPYB .NE. 'Y') THEN
3518          PRINT1 = 1
3519          PRINT2 = 2
3520      ENDIF
3521      IF (COPYA .NE. 'Y' .AND. COPYB .NE. 'Y') GOTO 98
3522
3523      SUBROUTINE ROUT$5
3524
3525      C ORGNSM:
3526      C -----
3527      C SUBROUTINE VARIES WITH ORGANISM
3528      C VARIABLES CHANGE WITH ORGANISM
3529      C VARIABLES NOT RETURNED TO ORGNSM, ONLY TO PRICES
3530      C CURRENT ORGANISM: BIVALVES
3531
3532      C STRUCTURE: READS OUT DATA RINGS IN SUBROUTINE INPUT,
3533      WHICH IS USED TO CALCULATE PRICES5 , ALGAE
3534
3535      C 1. COMMON BLOCKS
3536          INCLUDE '(IPAR1)'
3537          INCLUDE '(ICN5$)'
3538          INCLUDE '(ICOUNT)'
3539          INCLUDE '(ICBLK1)'
3540          INCLUDE '(ICBLK0)'
3541          INCLUDE '(ICMAIN)'
3542
3543          LOCAL VARIABLES
3544          CHARACTER*1,COPYA,COPYB
3545          INTEGER PRINT1,PRINT2,G,L,P,M,S,W
3546          NAME DESCRIPTION OF LOCAL VARIABLES AND ARGUMENTS
3547          COPYA OPTION TO PRINT INPUT DATA
3548          COPYB OPTION TO WRITE INPUT DATA
3549          G LOOP COUNTER FOR GOODS
3550          L LOOP COUNTER FOR GOODS
3551          M LOOP COUNTER FOR MONTHS
3552          P LOOP COUNTER FOR DIFFERENT PRINTING OPTIONS
3553          PRINT1 PRINTING INITIAL VALUE (1 OR 2)
3554          PRINT2 PRINTING FINAL VALUE (1 OR 2)
3555          S LOOP COUNTER STAGE
3556          W UNIT NUMBER TO WRITE
3557
3558          C TRANSLATE UNIT NUMBERS TO UNIT NUMBERS USED IN ORGNSM
3559          W = W0
3560
3561          C OPTION TO PRINT INPUT
3562          WRITE(WI,3000)
3563          3000 FORMAT(' / ', 'DO YOU WANT TO PRINT THE ALGAE PRICE DATA ',
3564              2,'ENTERED?')
3565          READ(RI,1000) COPYA
3566          1000 FORMAT( A1)
3567          WRITE(WI,3001)
3568          3001 FORMAT(' / ', 'DO YOU WANT TO SEE THE ALGAE DATA ON THE SCREEN? ',
3569              2,'Y OR N')
3570          READ(RI,1000) COPYB
3571          IF (COPYA .EQ. 'Y' .AND. COPYB .NE. 'Y') THEN
3572              PRINT1 = 1
3573              PRINT2 = 1
3574              IF (COPYB .EQ. 'Y' .AND. COPYA .NE. 'Y') THEN
3575                  PRINT1 = 1
3576                  PRINT2 = 1
3577              ENDIF
3578          ENDIF
3579
3580          PRINT2 = PRINT1,PRINT2
3581          IF (P .EQ. 2) V = WI
3582          WRITE(W,3300),NAME,DATE
3583          WRITE(W,3301),DATE2
3584          WRITE(W,3302)
3585          WRITE(W,3002) PWHR
3586          WRITE(W,3003) POIL
3587          WRITE(W,304) (PGOOD(G), G=1,NUMGOD)
3588          WRITE(W,305) NUMLAB
3589          DO 83 L=1,NUMLAB
3590              WRITE(W,306) L, PLABOR(L)
3591          CONTINUE
3592
3593          WRITE(W,3307)
3594          WRITE(W,308) WATAIR
3595          WRITE(W,309) WATBLB
3596          WRITE(W,310) BLBLIF
3597          WRITE(W,311) WATAUT
3598          WRITE(W,312) HPCOM
3599          WRITE(W,313) HRSOM
3600          WRITE(W,314) NUMTUB
3601          WRITE(W,315) COMEFF
3602          WRITE(W,316) PMPEFT
3603          WRITE(W,317) PMPGFM
3604          WRITE(W,318) HPPMP
3605          WRITE(W,3319)
3606          DO 84 L=1,NUMLAB
3607              WRITE(W,319) L, HRLABW(L)
3608          CONTINUE
3609
3610          WRITE(W,3320)
3611          WRITE(W,320) NUMSTG
3612          WRITE(W,3321)(S, S=1,NUMSTG)
3613          WRITE(W,321)(NULTR(S), S=1,NUMSTG)
3614          WRITE(W,322)(NUMINK(S), S=1,NUMSTG)
3615          WRITE(W,323)(STGRM(S),
3616          WRITE(W,324)(DAVSTG(S), S=1,NUMSTG)
3617          WRITE(W,325)(CPERM(S), S=1,NUMSTG)
3618          WRITE(W,326)(H2OINK(S),
3619          WRITE(W,327)(INCNTK(S), S=1,NUMSTG)
3620          WRITE(W,328)(DISSTG(S), S=1,NUMSTG)
3621          WRITE(W,329)(TUBINK(S), S=1,NUMSTG)
3622          WRITE(W,330)(BLBRM(S), S=1,NUMSTG)
3623          WRITE(W,331)(HRSBLB(S), S=1,NUMSTG)
3624          WRITE(W,332)(HRSAUT(S), S=1,NUMSTG)
3625          WRITE(W,333)(NDNAIR(S), S=1,NUMSTG)
3626
3627          DO 89 L=1,NUMLAB
3628              WRITE(W,334)L,(HRLABS(L,S), S=1,NUMSTG)
3629          CONTINUE
3630          WRITE(W,335) (NAMALG(S), S=1,NUMSTG)
3631
3632          WRITE(W,3336) BEGPRO, NAMEM(BEGPRO)
3633          WRITE(W,3337) ENDPRO, NAMEM(ENDPRO)
3634
3635

```





```

C      INTEGER STAGE, TIME
C      3. LOCAL VARIABLES
C         INTEGER L, MJ, S
C         REAL IQLATY(6)
C
C      NAME   DESCRIPTION OF LOCAL VARIABLES AND ARGUMENTS
C      CQLATY LABOR TYPE L REQUIRED FOR INNOCULATION
C      L      LOOP COUNTER FOR LABOR TYPES
C      MJ     TRANSLATES ARGUMENT TIME
C      S      STAGE ARGUE: CURRENT ALGAE STAGE
C      TIME   ARGUE: MONTH THAT CURRENT STAGE IS IN
C
C      TRANSLATE ARGUMENTS INTO LOCAL VARIABLES
C      S = STAGE
C      MJ = TIME
C
C      FIND QUANTITY OF LABOR TYPE L, FILE INTO ARRAY
C      ALLAB(S) = 0.
C      TALLAB(S) = 0.
C
DO 10 L = 1,NUMLAB
  TLABTY(L,S) = HRLABW(L)/7. *DAYSTG(S) /FLOAT(NUMSTG) +HRLABS(L,S)
C
C      ADD INNOCULATION - TEST FOR FIRST STAGE
C      IF (S .EQ. 1) THEN
C        IQLATY(L) = 0.
C        TTLYT(L,S) = TLABTY(L,S)
C      ELSE
C        IF (TLITRE(S-1) .LE. 0.) STOP 'ERROR LABOR; TLITRE < 0'
C        IQLATY(L) = INQTNK(S)*NUMTRK(S) * (TTLYT(L,S-1)/TLITRE(S-1))
C        TTLYT(L,S) = TLABTY(L,S) + IQLATY(L)
C      ENDIF
C
C      SUM ALL TYPES L
C      ALLAB(S) = TLABTY(L,S) + ALLAB(S)
C      TALLAB(S) = TTLYT(L,S) + TALLAB(S)
C
10    CONTINUE
C
C      TOTAL ALL LABOR USED PER LITRE
C      IF (TLITRE(S) .LE. 0.) STOP 'ERROR LABOR2; TLITRE = 0'
C      TALLAB(S) = TALLAB(S) / TLITRE(S)
C
RETURN
END
C
C      SUBROUTINE GOODS(STAGE,TIME)
C
C      COMMON BLOCK VARIABLES
C      INCLUDE (IPARI)
C      INCLUDE (ICNS$)
C
C      2. ARGUMENT VARIABLES
C      INTEGER STAGE, TIME
C      LOCAL VARIABLES
C      INTEGER G, S, MJ
C      REAL IQOTY(5)
C
C      NAME   DESCRIPTION OF LOCAL VARIABLES AND ARGUMENTS
C      CQLATY GOOD TYPE G REQUIRED FOR INNOCULATION
C      G      LOOP COUNTER FOR GOODS
C      MJ     TRANSLATES ARGUMENT TIME FOR USE IN SUBROUTINE
C      S      TRANSLATES ARGUMENT STAGE FOR USE IN SUBROUTINE
C      STAGE ARGUE: CURRENT STAGE OF ALGAE
C      TIME   ARGUE: MONTH THAT CURRENT STAGE IS IN
C
C      TRANSLATE ARGUMENTS INTO LOCAL VARIABLES
C      S = STAGE
C      MJ = TIME
C
3855
3856
3857
3858
3859
3860
3861
3862
3863
3864
3865
3866
3867
3868
3869
3870
3871
3872
3873
3874
3875
3876
3877
3878
3879
3880
3881
3882
3883
3884
3885
3886
3887
3888
3889
3890
3891
3892
3893
3894
3895
3896
3897
3898
3899
3900
3901
3902
3903
3904
3905
3906
3907
3908
3909
3910
3911
3912
3913
3914
3915
3916
3917
3918
3919
3920
3921
3922
3923
3924
3925
3926
3927
3928
3929
3930
3931
3932
3933
3934
3935
3936
3937
3938
3939
3940
3941
3942
3943
3944
3945
3946
3947
3948
3949
3950
3951
3952
3953
3954
3955
3956
3957
3958
3959
3960
3961
3962
C      ORGNSM:
C
C      SUBROUTINE VARIES WITH ORGANISM
C      VARIABLE CHANGE, RETURNED ONLY TO PRICES
C      CURRENT ORGANISM: BIVALVES
C
C      SUBROUTINE COST (STAGE,TIME)
C
C      ORGNSM:
C
C      SUBROUTINE VARIES WITH ORGANISM
C      VARIABLES VARY, BUT RETURNED ONLY TO PRICES
C      CURRENT ORGANISM: BIVALVES
C

```

```

4027 C SUBROUTINE: DETERMINES COST OF GIVEN ALGAE INPUT QUANT. FOR PRICE N=59963
4028 C
4029 C 1. COMMON BLOCK VARIABLES
4030 INCLUDE (IPAR1)
4031 INCLUDE (ICN5$)
4032 C 2. ARGUMENT VARIABLES
4033 INTEGER TIME, STAGE
4034 C 3. LOCAL VARIABLES
4035 INTEGER G, I, L, S
4036
4037 REAL CLABTY, TCGOTY, CGDODY
4038
4039 C NAME DESCRIPTION OF LOCAL VARIABLES AND ARGUMENTS
4040
4041 C C GODY COST OF GOOD, TYPE G, LESS INNOCULATION
4042 C CLABTY COST OF LABOR, TYPE L, LESS INNOCULATION
4043 C LOOP COUNTER FOR GOODS
4044 C I LOOP COUNTER FOR INPUTS
4045 C S LOOP COUNTER FOR LABOR TYPES
4046 C STAGE ARGUE: CURRENT STAGE OF ALGAE
4047 C TCGOTY TOTAL COST OF GOODS, TYPE G, LESS INNOCULATION
4048 C TIME ARGUE: MONTH THAT CURRENT STAGE IS IN
4049
4050
4051
4052
4053
4054
4055
4056
4057
4058
4059
4060
4061
4062
4063
4064
4065
4066
4067
4068
4069
4070
4071
4072
4073
4074
4075
4076
4077
4078
4079
4080
4081
4082
4083
4084
4085
4086
4087
4088
4089
4090
4091
4092
4093
4094
4095
4096
4097
4098
4099
4100
4101
4102
4103
4104
4105
4106
4107
4108
4109
4110
4111
4112
4113
4114
4115
4116
4117
4118
4119
4120
4121
4122
4123
4124
4125
4126
4127
4128
4129
4130
4131
4132
4133
4134
4135
4136
4137
4138
4139
4140
4141
4142
4143
4144
4145
4146
4147
4148
4149
4150
4151
4152
4153
4154
4155
4156
4157
4158
4159
4160
4161
4162
4163
4164
4165
4166
4167
4168
4169
4170
4171
4172
4173
4174
4175
4176
4177
4178
4179
4180
4181
4182
4183
4184
4185
4186
4187
4188
4189
4190
4191
4192
4193
4194
4195
4196
4197
4198
4199
4200
4201
4202
4203
4204
4205
4206
4207
4208
4209
4210
4211
4212
4213
4214
4215
4216
4217
4218
4219
4220
4221
4222
4223
4224
4225
4226
4227
4228
4229
4230
4231
4232
4233
4234
4235
4236
4237
4238
4239
4240
4241
4242
4243
4244
4245
4246
4247
4248
4249
4250
4251
4252
4253
4254
4255
4256
4257
4258
4259
4260
4261
4262
4263
4264
4265
4266
4267
4268
4269
4270
4271
4272
4273
4274
4275
4276
4277
4278
4279
4280
4281
4282
4283
4284
4285
4286
4287
4288
4289
4290
4291
4292
4293
4294
4295
4296
4297
4298
4299
4300
4301
4302
4303
4304
4305
4306
4307
4308
4309
4310
4311
4312
4313
4314
4315
4316
4317
4318
4319
4320
4321
4322
4323
4324
4325
4326
4327
4328
4329
4330
4331
4332
4333
4334
4335
4336
4337
4338
4339
4340
4341
4342
4343
4344
4345
4346
4347
4348
4349
4350
4351
4352
4353
4354
4355
4356
4357
4358
4359
4360
4361
4362
4363
4364
4365
4366
4367
4368
4369
4370
4371
4372
4373
4374
4375
4376
4377
4378
4379
4380
4381
4382
4383
4384
4385
4386
4387
4388
4389
4390
4391
4392
4393
4394
4395
4396
4397
4398
4399
4400
4401
4402
4403
4404
4405
4406
4407
4408
4409
4410
4411
4412
4413
4414
4415
4416
4417
4418
4419
4420
4421
4422
4423
4424
4425
4426
4427
4428
4429
4430
4431
4432
4433
4434
4435
4436
4437
4438
4439
4440
4441
4442
4443
4444
4445
4446
4447
4448
4449
4450
4451
4452
4453
4454
4455
4456
4457
4458
4459
4460
4461
4462
4463
4464
4465
4466
4467
4468
4469
4470
4471
4472
4473
4474
4475
4476
4477
4478
4479
4480
4481
4482
4483
4484
4485
4486
4487
4488
4489
4490
4491
4492
4493
4494
4495
4496
4497
4498
4499
4500
4501
4502
4503
4504
4505
4506
4507
4508
4509
4510
4511
4512
4513
4514
4515
4516
4517
4518
4519
4520
4521
4522
4523
4524
4525
4526
4527
4528
4529
4530
4531
4532
4533
4534
4535
4536
4537
4538
4539
4540
4541
4542
4543
4544
4545
4546
4547
4548
4549
4550
4551
4552
4553
4554
4555
4556
4557
4558
4559
4560
4561
4562
4563
4564
4565
4566
4567
4568
4569
4570
4571
4572
4573
4574
4575
4576
4577
4578
4579
4580
4581
4582
4583
4584
4585
4586
4587
4588
4589
4590
4591
4592
4593
4594
4595
4596
4597
4598
4599
4600
4601
4602
4603
4604
4605
4606
4607
4608
4609
4610
4611
4612
4613
4614
4615
4616
4617
4618
4619
4620
4621
4622
4623
4624
4625
4626
4627
4628
4629
4630
4631
4632
4633
4634
4635
4636
4637
4638
4639
4640
4641
4642
4643
4644
4645
4646
4647
4648
4649
4650
4651
4652
4653
4654
4655
4656
4657
4658
4659
4660
4661
4662
4663
4664
4665
4666
4667
4668
4669
4670
4671
4672
4673
4674
4675
4676
4677
4678
4679
4680
4681
4682
4683
4684
4685
4686
4687
4688
4689
4690
4691
4692
4693
4694
4695
4696
4697
4698
4699
4700
4701
4702
4703
4704
4705
4706
4707
4708
4709
4710
4711
4712
4713
4714
4715
4716
4717
4718
4719
4720
4721
4722
4723
4724
4725
4726
4727
4728
4729
4730
4731
4732
4733
4734
4735
4736
4737
4738
4739
4740
4741
4742
4743
4744
4745
4746
4747
4748
4749
4750
4751
4752
4753
4754
4755
4756
4757
4758
4759
4760
4761
4762
4763
4764
4765
4766
4767
4768
4769
4770
4771
4772
4773
4774
4775
4776
4777
4778
4779
4780
4781
4782
4783
4784
4785
4786
4787
4788
4789
4790
4791
4792
4793
4794
4795
4796
4797
4798
4799
4800
4801
4802
4803
4804
4805
4806
4807
4808
4809
4810
4811
4812
4813
4814
4815
4816
4817
4818
4819
4820
4821
4822
4823
4824
4825
4826
4827
4828
4829
4830
4831
4832
4833
4834
4835
4836
4837
4838
4839
4840
4841
4842
4843
4844
4845
4846
4847
4848
4849
4850
4851
4852
4853
4854
4855
4856
4857
4858
4859
4860
4861
4862
4863
4864
4865
4866
4867
4868
4869
4870
4871
4872
4873
4874
4875
4876
4877
4878
4879
4880
4881
4882
4883
4884
4885
4886
4887
4888
4889
4890
4891
4892
4893
4894
4895
4896
4897
4898
4899
4900
4901
4902
4903
4904
4905
4906
4907
4908
4909
4910
4911
4912
4913
4914
4915
4916
4917
4918
4919
4920
4921
4922
4923
4924
4925
4926
4927
4928
4929
4930
4931
4932
4933
4934
4935
4936
4937
4938
4939
4940
4941
4942
4943
4944
4945
4946
4947
4948
4949
4950
4951
4952
4953
4954
4955
4956
4957
4958
4959
4960
4961
4962
4963
4964
4965
4966
4967
4968
4969
4970
4971
4972
4973
4974
4975
4976
4977
4978
4979
4980
4981
4982
4983
4984
4985
4986
4987
4988
4989
4990
4991
4992
4993
4994
4995
4996
4997
4998
4999
4999

```

WRITE(W,1101) NAME, DATE  
 WRITE(W,101) DATE2  
 WRITE(W,102) MONBEG  
 WRITE(W,1103) (S,  
 S = 1,NUMSTG)  
 WRITE(W,103) (MONAVA( S), S = 1,NUMSTG)  
 WRITE(W,1031) (TLTRE( S), S = 1,LASSTG)  
 WRITE(W,104) (LFOOD( S), S = 1,LASSTG)  
 WRITE(W,105) (TCCELL( S), S = 1,LASSTG)  
 WRITE(W,108) (TCALIN( S), S = 1,LASSTG)  
 WRITE(W,109) (TCCLTR(S), S = 1,LASSTG)  
 WRITE(W,1092) (TCCELL( S), S = 1,LASSTG)  
 WRITE(W,110) (CALIN( S), S = 1,LASSTG)  
 WRITE(W,113) (TCINP (1,S), S = 1,LASSTG)  
 WRITE(W,114) (TCINP (1,S), S = 1,LASSTG)  
 WRITE(W,115) (CINP (1,S), S = 1,LASSTG)  
 WRITE(W,116) (CTKWHR( S), S = 1,LASSTG)  
 WRITE(W,117) (CTKWHL( S), S = 1,LASSTG)  
 WRITE(W,118) (TKWHR( S), S = 1,LASSTG)  
 WRITE(W,119) (TCINP (2,S), S = 1,LASSTG)  
 WRITE(W,120) (TCINPL(2,S), S = 1,LASSTG)  
 WRITE(W,121) (CINP (2,S), S = 1,LASSTG)  
 WRITE(W,122) (TTOIL( S), S = 1,LASSTG)  
 WRITE(W,123) (TTOILL( S), S = 1,LASSTG)  
 WRITE(W,124) (TTOIL( S), S = 1,LASSTG)  
 WRITE(W,125) (TCINP (3,S), S = 1,LASSTG)  
 WRITE(W,126) (TCINPL(3,S), S = 1,LASSTG)  
 WRITE(W,127) (CINP (3,S), S = 1,LASSTG)  
 WRITE(W,128) (TALLAB( S), S = 1,LASSTG)  
 WRITE(W,129) (TALLA( S), S = 1,LASSTG)  
 WRITE(W,130) (CALLAB( S), S = 1,LASSTG)  
 WRITE(W,131) (TCINP (4,S), S = 1,LASSTG)  
 WRITE(W,132) (TCINPL(4,S), S = 1,LASSTG)  
 WRITE(W,133) (CINP (4,S), S = 1,LASSTG)  
 1101 FORMAT ('1//', 'A10', 'COST SIMULATION MODEL; ALGAE PRICE ',  
 2 'COMPUTATIONS', /, '59(- -)', 'DATE: ', A25/  
 3 'OUTPUT GENERATED',  
 101 FORMAT ('0', 'DATE OF PRICE COMPUTATIONS: ', A11/)  
 102 FORMAT ('0', 'BATCH BEGIN: ', 2X, A1/)  
 1103 FORMAT ('0', '143,6(3X,12,'STAGE',5X)', /, 'T43,6(3X,12,'STAGE',5X)', /)  
 103 FORMAT ('0', 'ALGAE AVAILABLE FOR USE IN MONTH',  
 B T41, 6(4X,A9,2X), /, 'T41, 6(4X,A9,2X)')  
 1031 FORMAT ('0', '\*TOTAL LITRES PRODUCED',  
 B 'CELLS OF ALGAE PRODUCED',  
 108 FORMAT ('0', '\*TOTAL COST OF STAGE',  
 B T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 109 FORMAT ('0', 'LITRES OF ALGAE AVAILABLE',  
 B T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 105 FORMAT ('0', 'CELLS OF ALGAE PRODUCED',  
 B T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 1092 FORMAT ('0', '1X, 'TOTAL COST PER CELL OF ALGAE',  
 B T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 110 FORMAT ('0', '1X, 'TOTAL COST OF STAGE, LESS INNOCULATION',  
 B T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 113 FORMAT ('0', '\*TOTAL COST OF ELECTRICITY',  
 C -----)

4091 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4092 B FORMAT ('0', '1X, 'TOTAL COST OF ELECTRICITY/LITRE OF ALGAE',  
 4093 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4094 B FORMAT ('0', 'TOTAL COST OF ELEC. LESS INNOCULATION',  
 4095 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4096 B FORMAT ('0', '\*TOTAL KWH USED',  
 4097 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4098 B FORMAT ('0', '1X, 'TOTAL KWH USED PER LITRE OF ALGAE',  
 4099 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4100 B FORMAT ('0', '1X, 'TOTAL USED LESS INNOCULATION',  
 4101 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4102 B FORMAT ('0', '\*TOTAL COST OF FUEL OIL',  
 4103 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4104 B FORMAT ('0', '1X, 'TOTAL COST OF OIL/LITRE OF ALGAE',  
 4105 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4106 B FORMAT ('0', '1X, 'TOTAL COST OF OIL LESS INNOCULATION',  
 4107 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4108 B FORMAT ('0', '1X, 'TOTAL COST OF OIL/LITRE OF ALGAE',  
 4109 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4110 B FORMAT ('0', '1X, 'TOTAL LITRES OF FUEL OIL',  
 4111 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4112 B FORMAT ('0', '1X, 'TOTAL LITRES OIL USED/LITRE OF ALGAE',  
 4113 B FORMAT ('0', '1X, 'TOTAL L. OIL USED LESS INNOCULATION',  
 4114 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4115 B FORMAT ('0', '\*TOTAL COST OF ALL LABOR TYPES',  
 4116 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4117 B FORMAT ('0', '1X, 'TOTAL COST ALL LABOR/LITRE OF ALGAE',  
 4118 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4119 B FORMAT ('0', '1X, 'TOTAL COST ALL LABOR LESS INNOCULATN',  
 4120 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4121 B FORMAT ('0', '1X, 'TOTAL HOURS OF ALL LABOR TYPES',  
 4122 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4123 B FORMAT ('0', '1X, 'TOTAL HRS ALL LABOR USED/LITRE ALGAE',  
 4124 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4125 B FORMAT ('0', '1X, 'TOTAL HRS ALL LABOR LESS INNOCULATION',  
 4126 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4127 B FORMAT ('0', '1X, 'TOTAL HRS ALL LABOR LESS INNOCULATION',  
 4128 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4129 B FORMAT ('0', '\*TOTAL COST OF ALL MATERIALS',  
 4130 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4131 B FORMAT ('0', '1X, 'TOTAL COST ALL MATERIALS/LITRE OF ALGAE',  
 4132 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4133 B FORMAT ('0', '1X, 'TOTAL COST ALL MATER. LESS INNOCULATION',  
 4134 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4135 B FORMAT ('0', '1X, 'TOTAL COST OF STAGE',  
 4136 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4137 B FORMAT ('0', '1X, 'TOTAL COST OF STAGE',  
 4138 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4139 B FORMAT ('0', '1X, 'TOTAL COST OF STAGE',  
 4140 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4141 B FORMAT ('0', '1X, 'TOTAL COST OF STAGE',  
 4142 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4143 B FORMAT ('0', '1X, 'TOTAL COST OF STAGE',  
 4144 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4145 B FORMAT ('0', '1X, 'TOTAL COST OF STAGE',  
 4146 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4147 B FORMAT ('0', '1X, 'TOTAL COST OF STAGE',  
 4148 B FORMAT ('0', 'T41, 6(G13,4,2X), /, 'T41, 6(G13,4,2X)')  
 4149 B FORMAT ('0', '1X, 'TOTAL COST OF STAGE',  
 4150 B FORMAT ('0', '1X, 'TOTAL COST OF STAGE, LESS INNOCULATION',  
 4151 B FORMAT ('0', '1X, 'TOTAL COST OF STAGE',  
 4152 B FORMAT ('0', '1X, 'TOTAL COST OF STAGE',  
 4153 B FORMAT ('0', '\*TOTAL COST OF ELECTRICITY',  
 4154 C -----)

72

SUBROUTINE PRICE6 (WRT, RED, HREAD, RUN2, RUN3)  
 RETURN  
 END

SUBROUTINE PRICE6 (WRT, RED, HREAD, RUN2, RUN3)  
 RETURN  
 END





```

C      ARGUE: STAGE THAT CURRENT PERIOD IS IN          4539
C      J      TRANSLATES JNOW FOR CURRENT PERIOD        4540
C      ARGUE: CURRENT PERIOD BEING CALCULATED       4541
C      M      TRANSLATE MNOW FOR CURRENT MONTH        4542
C      MNOW   ARGUE: MONTH THAT CURRENT J IS IN        4543
C      TNO    SUM OF ALL ORGANISMS EXISTING EACH DAY OF PERIOD 4544
C      VALUE   VALUE TO DECREASE REIMANN SUM. 1ST TERM OF SUM 4545
C      WKS    TRANSLATE WKSNOW FOR ELAPSED WEEKS       4546
C      WKSNOW ARGUE: ELAPSED WEEKS FOR CURRENT J       4547
C
C      TRANSLATE LOCAL VARIABLES                      4548
C
C      I      = INOW                                     4549
C      J      = JNOW                                     4550
C      WKS   = WKSNOW                                   4551
C      M      = MNOW                                     4552
C      DAY    = DAYNOW                                  4553
C
C      ALGORITHM: DEPENDS ON TYPE OF ORGANISM          4554
C
C      IF (OTYPE .EQ. 1 .OR. OTYPE .EQ. 2) THEN          4555
C          WHEN 1=CONDITIONING, ASSUME NO DEATH RATE AND ANIMALS ALL AT 4556
C          TWO YEARS OLD SO AS NOT TO EXCEED MAXIMUM OF AGE/SIZE EQUATION 4557
C          IF (I .EQ. 1) THEN                           4558
C              AGEUPJ(J) = 365. * 2.                  4559
C              ONOOFJ(J) = NO011                      4560
C              ONOOFPJ(J) = NO011                     4561
C
C          ALL OTHER STAGES, COMPUTE NUMBER OF ORGANISMS SURVIVING AT EACH 4562
C          DAY OF THE PERIOD BY CALLING FUNCTION AGENO. THEN FIND AVERAGE 4563
C          SURVIVAL OF THE PERIOD BY SUMMING USING THE TRAPEZOIDAL RULE 4564
C          ELSEIF (I .GT. 1) THEN                      4565
C
C              DCNTR = 0                                4566
C              TNO = 0.                                 4567
C              DO 100 D = (DAY-DAYSJ(1)-DAYSII+1.), (DAY-DAYSII+.499999) 4568
C                  DWK = FLOAT(D) / 7.                 4569
C                  ONOUPJ(J) = AGENO(DWK)             4570
C                  TNO = TNO + ONOUPJ(J)            4571
C                  DCNTR = DCNTR + 1                4572
C
C              COMPUTE NUMBER DURING DAY BEFORE PERIOD BEGINS          4573
C              DWK = (DAY-DAYS(I)-DAYSII) / 7.           4574
C              VALUE = AGENO(DWK)                      4575
C
C              INCREASE TOTAL BY 1/2 VALUE, DECREASE BY 1/2 LAST SUM TERM 4576
C              TNO = TNO + ((VALUE - ONOUPJ(J)) / 2.)          4577
C
C              COMPUTE AVERAGE VALUE WHERE DELTA T IS ALWAYS = 1          4578
C              ONOOFJ(J) = TNO / FLOAT(DCNTR)            4579
C
C              AGE OF ANIMAL AT END OF PERIOD DOESNT INCLUDE CONDITIONING 4580
C              AGEUPJ(J) = DAY - DAYSII               4581
C
C              ENDIF                                     4582
C
C              ENDIF                                     4583
C
C              ELSE STOP 'ORGANISM TYPE INCORRECT IN SUBR. NUMBR.'      4584
C              ENDIF                                     4585
C
C              MNOW = MNOW + WKS                         4586
C              WKS = WKS - DWK                          4587
C              MNOW = MNOW - WKS                         4588
C
C              MNOW = MNOW - WKS                         4589
C              WKS = WKS - DWK                          4590
C              MNOW = MNOW - WKS                         4591
C
C              WKS = WKS - DWK                          4592
C
C              MNOW = MNOW - WKS                         4593
C              WKS = WKS - DWK                          4594
C
C              MNOW = MNOW - WKS                         4595
C
C              MNOW = MNOW - WKS                         4596
C
C              MNOW = MNOW - WKS                         4597
C
C              MNOW = MNOW - WKS                         4598
C
C              MNOW = MNOW - WKS                         4599
C
C              MNOW = MNOW - WKS                         4600
C
C              MNOW = MNOW - WKS                         4601
C
C              MNOW = MNOW - WKS                         4602
C
C      TRANSLATE LOCAL AND ARGUMENT VARIABLES          4603
C
C      NAME   DESCRIPTION OF LOCAL AND ARGUMENT VARIABLES
C
C      D      LOOP COUNTER FOR DAYS                  4604
C      DAY   TRANSLATE DAYNOW FOR ELAPSED DAYS OF PERIOD 4605
C      DCNTR LAST DAY OF PERIOD LOOED THROUGH        4606
C      DWK   TRANSLATE DAYS LOOED IN PERIOD INTO WEEKS 4607
C      INOW  TRANSLATE INOW INTO STAGE OF CURRENT PERIOD 4608
C      J      ARGUE: STAGE THAT CURRENT J IS IN        4609
C      JNOW  TRANSLATE JNOW FOR CURRENT PERIOD        4610
C      M      ARGUE: CURRENT PERIOD                  4611
C      MNOW  TRANSLATE MNOW FOR MONTH THAT CURRENT PERIOD OCCURS IN 4612
C      TSZE  SUM OF SIZE FOR ALL DAYS OVER THE PERIOD 4613
C      WKS   VALUE OF 1ST TERM TO SUBTRACT OFF OF THE REIMANN SUM 4614
C      WKS   TRANSLATE WKSNOW FOR ELAPSED WEEKS       4615
C      WKS   ARGUE: ELAPSED WEEKS TO END OF CURRENT PERIOD 4616
C
C      TRANSLATE ARGUMENTS INTO LOCAL VARIABLES          4617
C
C      I      = INOW                                     4618
C      J      = JNOW                                     4619
C      WKS   = WKSNOW                                   4620
C      M      = MNOW                                     4621
C      DAY   = DAYNOW                                  4622
C
C      ENDIF                                     4623
C
C      ENDIF                                     4624
C
C      ENDIF                                     4625
C
C      ENDIF                                     4626
C
C      ENDIF                                     4627
C
C      ENDIF                                     4628
C
C      ENDIF                                     4629
C
C      ENDIF                                     4630
C
C      ENDIF                                     4631
C
C      ENDIF                                     4632
C
C      ENDIF                                     4633
C
C      ENDIF                                     4634
C
C      ENDIF                                     4635
C
C      ENDIF                                     4636
C
C      ENDIF                                     4637
C
C      ENDIF                                     4638
C
C      ENDIF                                     4639
C
C      ENDIF                                     4640
C
C      ENDIF                                     4641
C
C      ENDIF                                     4642
C
C      ENDIF                                     4643
C
C      ENDIF                                     4644
C
C      ENDIF                                     4645
C
C      ENDIF                                     4646
C
C      ENDIF                                     4647
C
C      ENDIF                                     4648
C
C      ENDIF                                     4649
C
C      ENDIF                                     4650
C
C      ENDIF                                     4651
C
C      ENDIF                                     4652
C
C      ENDIF                                     4653
C
C      ENDIF                                     4654
C
C      ENDIF                                     4655
C
C      ENDIF                                     4656
C
C      ENDIF                                     4657
C
C      ENDIF                                     4658
C
C      ENDIF                                     4659
C
C      ENDIF                                     4660
C
C      ENDIF                                     4661
C
C      ENDIF                                     4662
C
C      ENDIF                                     4663
C
C      ENDIF                                     4664
C
C      ENDIF                                     4665
C
C      ENDIF                                     4666
C
C      ENDIF                                     4667
C
C      ENDIF                                     4668
C
C      ENDIF                                     4669
C
C      ENDIF                                     4670
C
C      ENDIF                                     4671
C
C      ENDIF                                     4672
C
C      ENDIF                                     4673
C
C      ENDIF                                     4674
C
C      ENDIF                                     4675
C
C      ENDIF                                     4676
C
C      ENDIF                                     4677
C
C      ENDIF                                     4678
C
C      ENDIF                                     4679
C
C      ENDIF                                     4680
C
C      ENDIF                                     4681
C
C      ENDIF                                     4682
C
C      ENDIF                                     4683
C
C      ENDIF                                     4684
C
C      ENDIF                                     4685
C
C      ENDIF                                     4686
C
C      ENDIF                                     4687
C
C      ENDIF                                     4688
C
C      ENDIF                                     4689
C
C      ENDIF                                     4690
C
C      ENDIF                                     4691
C
C      ENDIF                                     4692
C
C      ENDIF                                     4693
C
C      ENDIF                                     4694
C
C      ENDIF                                     4695
C
C      ENDIF                                     4696
C
C      ENDIF                                     4697
C
C      ENDIF                                     4698
C
C      ENDIF                                     4699
C
C      ENDIF                                     4700
C
C      ENDIF                                     4701
C
C      ENDIF                                     4702
C
C      ENDIF                                     4703
C
C      ENDIF                                     4704
C
C      ENDIF                                     4705
C
C      ENDIF                                     4706
C
C      ENDIF                                     4707
C
C      ENDIF                                     4708
C
C      ENDIF                                     4709
C
C      ENDIF                                     4710
C
C      ENDIF                                     4711
C
C      ENDIF                                     4712
C
C      ENDIF                                     4713
C
C      ENDIF                                     4714
C
C      ENDIF                                     4715
C
C      ENDIF                                     4716
C
C      ENDIF                                     4717
C
C      ENDIF                                     4718
C
C      ENDIF                                     4719
C
C      ENDIF                                     4720
C
C      ENDIF                                     4721
C
C      ENDIF                                     4722
C
C      ENDIF                                     4723
C
C      ENDIF                                     4724
C
C      ENDIF                                     4725
C
C      ENDIF                                     4726
C
C      ENDIF                                     4727
C
C      ENDIF                                     4728
C
C      ENDIF                                     4729
C
C      ENDIF                                     4730
C
C      ENDIF                                     4731
C
C      ENDIF                                     4732
C
C      ENDIF                                     4733
C
C      ENDIF                                     4734
C
C      ENDIF                                     4735
C
C      ENDIF                                     4736
C
C      ENDIF                                     4737
C
C      ENDIF                                     4738
C
C      ENDIF                                     4739
C
C      ENDIF                                     4740
C
C      ENDIF                                     4741
C
C      ENDIF                                     4742
C
C      ENDIF                                     4743
C
C      ENDIF                                     4744
C
C      ENDIF                                     4745
C
C      ENDIF                                     4746
C
C      ENDIF                                     4747
C
C      ENDIF                                     4748
C
C      ENDIF                                     4749
C
C      ENDIF                                     4750
C
C      ENDIF                                     4751
C
C      ENDIF                                     4752
C
C      ENDIF                                     4753
C
C      ENDIF                                     4754
C
C      ENDIF                                     4755
C
C      ENDIF                                     4756
C
C      ENDIF                                     4757
C
C      ENDIF                                     4758
C
C      ENDIF                                     4759
C
C      ENDIF                                     4760
C
C      ENDIF                                     4761
C
C      ENDIF                                     4762
C
C      ENDIF                                     4763
C
C      ENDIF                                     4764
C
C      ENDIF                                     4765
C
C      ENDIF                                     4766
C
C      ENDIF                                     4767
C
C      ENDIF                                     4768
C
C      ENDIF                                     4769
C
C      ENDIF                                     4770
C
C      ENDIF                                     4771
C
C      ENDIF                                     4772
C
C      ENDIF                                     4773
C
C      ENDIF                                     4774
C
C      ENDIF                                     4775
C
C      ENDIF                                     4776
C
C      ENDIF                                     4777
C
C      ENDIF                                     4778
C
C      ENDIF                                     4779
C
C      ENDIF                                     4780
C
C      ENDIF                                     4781
C
C      ENDIF                                     4782
C
C      ENDIF                                     4783
C
C      ENDIF                                     4784
C
C      ENDIF                                     4785
C
C      ENDIF                                     4786
C
C      ENDIF                                     4787
C
C      ENDIF                                     4788
C
C      ENDIF                                     4789
C
C      ENDIF                                     4790
C
C      ENDIF                                     4791
C
C      ENDIF                                     4792
C
C      ENDIF                                     4793
C
C      ENDIF                                     4794
C
C      ENDIF                                     4795
C
C      ENDIF                                     4796
C
C      ENDIF                                     4797
C
C      ENDIF                                     4798
C
C      ENDIF                                     4799
C
C      ENDIF                                     4800
C
C      ENDIF                                     4801
C
C      ENDIF                                     4802
C
C      ENDIF                                     4803
C
C      ENDIF                                     4804
C
C      ENDIF                                     4805
C
C      ENDIF                                     4806
C
C      ENDIF                                     4807
C
C      ENDIF                                     4808
C
C      ENDIF                                     4809
C
C      ENDIF                                     4810
C
C      ENDIF                                     4811
C
C      ENDIF                                     4812
C
C      ENDIF                                     4813
C
C      ENDIF                                     4814
C
C      ENDIF                                     4815
C
C      ENDIF                                     4816
C
C      ENDIF                                     4817
C
C      ENDIF                                     4818
C
C      ENDIF                                     4819
C
C      ENDIF                                     4820
C
C      ENDIF                                     4821
C
C      ENDIF                                     4822
C
C      ENDIF                                     4823
C
C      ENDIF                                     4824
C
C      ENDIF                                     4825
C
C      ENDIF                                     4826
C
C      ENDIF                                     4827
C
C      ENDIF                                     4828
C
C      ENDIF                                     4829
C
C      ENDIF                                     4830
C
C      ENDIF                                     4831
C
C      ENDIF                                     4832
C
C      ENDIF                                     4833
C
C      ENDIF                                     4834
C
C      ENDIF                                     4835
C
C      ENDIF                                     4836
C
C      ENDIF                                     4837
C
C      ENDIF                                     4838
C
C      ENDIF                                     4839
C
C      ENDIF                                     4840
C
C      ENDIF                                     4841
C
C      ENDIF                                     4842
C
C      ENDIF                                     4843
C
C      ENDIF                                     4844
C
C      ENDIF                                     4845
C
C      ENDIF                                     4846
C
C      ENDIF                                     4847
C
C      ENDIF                                     4848
C
C      ENDIF                                     4849
C
C      ENDIF                                     4850
C
C      ENDIF                                     4851
C
C      ENDIF                                     4852
C
C      ENDIF                                     4853
C
C      ENDIF                                     4854
C
C      ENDIF                                     4855
C
C      ENDIF                                     4856
C
C      ENDIF                                     4857
C
C      ENDIF                                     4858
C
C      ENDIF                                     4859
C
C      ENDIF                                     4860
C
C      ENDIF                                     4861
C
C      ENDIF                                     4862
C
C      ENDIF                                     4863
C
C      ENDIF                                     4864
C
C      ENDIF                                     4865
C
C      ENDIF                                     4866
C
C      ENDIF                                     4867
C
C      ENDIF                                     4868
C
C      ENDIF                                     4869
C
C      ENDIF                                     4870
C
C      ENDIF                                     4871
C
C      ENDIF                                     4872
C
C      ENDIF                                     4873
C
C      ENDIF                                     4874
C
C      ENDIF                                     4875
C
C      ENDIF                                     4876
C
C      ENDIF                                     4877
C
C      ENDIF                                     4878
C
C      ENDIF                                     4879
C
C      ENDIF                                     4880
C
C      ENDIF                                     4881
C
C      ENDIF                                     4882
C
C      ENDIF                                     4883
C
C      ENDIF                                     4884
C
C      ENDIF                                     4885
C
C      ENDIF                                     4886
C
C      ENDIF                                     4887
C
C      ENDIF                                     4888
C
C      ENDIF                                     4889
C
C      ENDIF                                     4890
C
C      ENDIF                                     4891
C
C      ENDIF                                     4892
C
C      ENDIF                                     4893
C
C      ENDIF                                     4894
C
C      ENDIF                                     4895
C
C      ENDIF                                     4896
C
C      ENDIF                                     4897
C
C      ENDIF                                     4898
C
C      ENDIF                                     4899
C
C      ENDIF                                     4900
C
C      ENDIF                                     4901
C
C      ENDIF                                     4902
C
C      ENDIF                                     4903
C
C      ENDIF                                     4904
C
C      ENDIF                                     4905
C
C      ENDIF                                     4906
C
C      ENDIF                                     4907
C
C      ENDIF                                     4908
C
C      ENDIF                                     4909
C
C      ENDIF                                     4910
C
C      ENDIF                                     4911
C
C      ENDIF                                     4912
C
C      ENDIF                                     4913
C
C      ENDIF                                     4914
C
C      ENDIF                                     4915
C
C      ENDIF                                     4916
C
C      ENDIF                                     4917
C
C      ENDIF                                     4918
C
C      ENDIF                                     4919
C
C      ENDIF                                     4920
C
C      ENDIF                                     4921
C
C      ENDIF                                     4922
C
C      ENDIF                                     4923
C
C      ENDIF                                     4924
C
C      ENDIF                                     4925
C
C      ENDIF                                     4926
C
C      ENDIF                                     4927
C
C      ENDIF                                     4928
C
C      ENDIF                                     4929
C
C      ENDIF                                     4930
C
C      ENDIF                                     4931
C
C      ENDIF                                     4932
C
C      ENDIF                                     4933
C
C      ENDIF                                     4934
C
C      ENDIF                                     4935
C
C      ENDIF                                     4936
C
C      ENDIF                                     4937
C
C      ENDIF                                     4938
C
C      ENDIF                                     4939
C
C      ENDIF                                     4940
C
C      ENDIF                                     4941
C
C      ENDIF                                     4942
C
C      ENDIF                                     4943
C
C      ENDIF                                     4944
C
C      ENDIF                                     4945
C
C      ENDIF                                     4946
C
C      ENDIF                                     4947
C
C      ENDIF                                     4948
C
C      ENDIF                                     4949
C
C      ENDIF                                     4950
C
C      ENDIF                                     4951
C
C      ENDIF                                     4952
C
C      ENDIF                                     4953
C
C      ENDIF                                     4954
C
C      ENDIF                                     4955
C
C      ENDIF                                     4956
C
C      ENDIF                                     4957
C
C      ENDIF                                     4958
C
C      ENDIF                                     4959
C
C      ENDIF                                     4960
C
C      ENDIF                                     4961
C
C      ENDIF                                     4962
C
C      ENDIF                                     4963
C
C      ENDIF                                     4964
C
C      ENDIF                                     4965
C
C      ENDIF                                     4966
C
C      ENDIF                                     4967
C
C      ENDIF                                     4968
C
C      ENDIF                                     4969
C
C      ENDIF                                     4970
C
C      ENDIF                                     4971
C
C      ENDIF                                     4972
C
C      ENDIF                                     4973
C
C      ENDIF                                     4974
C
C      ENDIF                                     4975
C
C      ENDIF                                     4976
C
C      ENDIF                                     4977
C
C      ENDIF                                     4978
C
C      ENDIF                                     4979
C
C      ENDIF                                     4980
C
C      ENDIF                                     4981
C
C      ENDIF                                     4982
C
C      ENDIF                                     4983
C
C      ENDIF                                     4984
C
C      ENDIF                                     4985
C
C      ENDIF                                     4986
C
C      ENDIF                                     4987
C
C      ENDIF                                     4988
C
C      ENDIF                                     4989
C
C      ENDIF                                     4990
C
C      ENDIF                                     4991
C
C      ENDIF                                     4992
C
C      ENDIF                                     4993
C
C      ENDIF                                     4994
C
C      ENDIF                                     4995
C
C      ENDIF                                     4996
C
C      ENDIF                                     4997
C
C      ENDIF                                     4998
C
C      ENDIF                                     4999
C
C      ENDIF                                     5000
C
C      ENDIF                                     5001
C
C      ENDIF                                     5002
C
C      ENDIF                                     5003
C
C      ENDIF                                     5004
C
C      ENDIF                                     5005
C
C      ENDIF                                     5006
C
C      ENDIF                                     5007
C
C      ENDIF                                     5008
C
C      ENDIF                                     5009
C
C      ENDIF                                     5010
C
C      ENDIF                                     5011
C
C      ENDIF                                     5012
C
C      ENDIF                                     5013
C
C      ENDIF                                     5014
C
C      ENDIF                                     5015
C
C      ENDIF                                     5016
C
C      ENDIF                                     5017
C
C      ENDIF                                     5018
C
C      ENDIF                                     5019
C
C      ENDIF                                     5020
C
C      ENDIF                                     5021
C
C      ENDIF                                     5022
C
C      ENDIF                                     5023
C
C      ENDIF                                     5024
C
C      ENDIF                                     5025
C
C      ENDIF                                     5026
C
C      ENDIF                                     5027
C
C      ENDIF                                     5028
C
C      ENDIF                                     5029
C
C      ENDIF                                     5030
C
C      ENDIF                                     5031
C
C      ENDIF                                     5032
C
C      ENDIF                                     5033
C
C      ENDIF                                     5034
C
C      ENDIF                                     5035
C
C      ENDIF                                     5036
C
C      ENDIF                                     5037
C
C      ENDIF                                     5038
C
C      ENDIF                                     5039
C
C      ENDIF                                     5040
C
C      ENDIF                                     5041
C
C      ENDIF                                     5042
C
C      ENDIF                                     5043
C
C      ENDIF                                     5044
C
C      ENDIF                                     5045
C
C      ENDIF                                     5046
C
C      ENDIF                                     5047
C
C      ENDIF                                     5048
C
C      ENDIF                                     5049
C
C      ENDIF                                     5050
C
C      ENDIF                                     5051
C
C      ENDIF                                     5052
C
C      ENDIF                                     5053
C
C      ENDIF                                     5054
C
C      ENDIF                                     5055
C
C      ENDIF                                     5056
C
C      ENDIF                                     5057
C
C      ENDIF                                     5058
C
C      ENDIF                                     5059
C
C      ENDIF                                     5060
C
C      ENDIF                                     5061
C
C      ENDIF                                     5062
C
C      ENDIF                                     5063
C
C      ENDIF                                     5064
C
C      ENDIF                                     5065
C
C      ENDIF                                     5066
C
C      ENDIF                                     5067
C
C      ENDIF                                     5068
C
C      ENDIF                                     5069
C
C      ENDIF                                     5070
C
C      ENDIF                                     5071
C
C      ENDIF                                     5072
C
C      ENDIF                                     5073
C
C      ENDIF                                     5074
C
C      ENDIF                                     5075
C
C      ENDIF                                     5076
C
C      ENDIF                                     5077
C
C      ENDIF                                     5078
C
C      ENDIF                                     5079
C
C      ENDIF                                     5080
C
C      ENDIF                                     5081
C
C      ENDIF                                     5082
C
C      ENDIF                                     5083
C
C      ENDIF                                     5084
C
C      ENDIF                                     5085
C
C      ENDIF                                     5086
C
C      ENDIF                                     5087
C
C      ENDIF                                     5088
C
C      ENDIF                                     5089
C
C      ENDIF                                     5090
C
C      ENDIF                                     5091
C
C      ENDIF                                     5092
C
C      ENDIF                                     5093
C
C      ENDIF                                     5094
C
C      ENDIF                                     5095
C
C      ENDIF                                     5096
C
C      ENDIF                                     5097
C
C      ENDIF                                     5098
C
C      ENDIF                                     5099
C
C      ENDIF                                     5100
C
C      ENDIF                                     5101
C
C      ENDIF                                     5102
C
C      ENDIF                                     5103
C
C      ENDIF                                     5104
C
C      ENDIF                                     5105
C
C      ENDIF                                     5106
C
C      ENDIF                                     5107
C
C      ENDIF                                     5108
C
C      ENDIF                                     5109
C
C      ENDIF                                     5110
C
C      ENDIF                                     5111
C
C      ENDIF                                     5112
C
C      ENDIF                                     5113
C
C      ENDIF                                     5114
C
C      ENDIF                                     5115
C
C      ENDIF                                     5116
C
C      ENDIF                                     5117
C
C      ENDIF                                     5118
C
C      ENDIF                                     5119
C
C      ENDIF                                     5120
C
C      ENDIF                                     5121
C
C      ENDIF                                     5122
C
C      ENDIF                                     5123
C
C      ENDIF                                     5124
C
C      ENDIF                                     5125
C
C      ENDIF                                     5126
C
C      ENDIF                                     5127
C
C      ENDIF                                     5128
C
C      ENDIF                                     5129
C
C      ENDIF                                     5130
C
C      ENDIF                                     5131
C
C      ENDIF                                     5132
C
C      ENDIF                                     5133
C
C      ENDIF                                     5134
C
C      ENDIF                                     5135
C
C      ENDIF                                     5136
C
C      ENDIF                                     5137
C
C      ENDIF                                     5138
C
C      ENDIF                                     5139
C
C      ENDIF                                     5140
C
C      ENDIF                                     5141
C
C      ENDIF                                     5142
C
C      ENDIF                                     5143
C
C      ENDIF                                     5144
C
C      ENDIF                                     5145
C
C      ENDIF                                     5146
C
C      ENDIF                                     5147
C
C      ENDIF                                     5148
C
C      ENDIF                                     5149
C
C      ENDIF                                     5150
C
C      ENDIF                                     5151
C
C      ENDIF                                     5152
C
C      ENDIF                                     5153
C
C      ENDIF                                     5154
C
C      ENDIF                                     5155
C
C      ENDIF                                     5156
C
C      ENDIF                                     5157
C
C      ENDIF                                     5158
C
C      ENDIF                                     5159
C
C      ENDIF                                     5160
C
C      ENDIF                                     5161
C
C      ENDIF                                     5162
C
C      ENDIF                                     5163
C
C      ENDIF                                     5164
C
C      ENDIF                                     5165
C
C      ENDIF                                     5166
C
C      ENDIF                                     5167
C
C      ENDIF                                     5168
C
C      ENDIF                                     5169
C
C      ENDIF                                     5170
C
C      ENDIF                                     5171
C
C      ENDIF                                     5172
C
C      ENDIF                                     5173
C
C      ENDIF                                     5174
C
C      ENDIF                                     5175
C
C      ENDIF                                     5176
C
C      ENDIF                                     5177
C
C      ENDIF                                     5178
C
C      ENDIF                                     5179
C
C      ENDIF                                     5180
C
C      ENDIF                                     5181
C
C      ENDIF                                     5182
C
C      ENDIF                                     5183
C
C      ENDIF                                     5184
C
C      ENDIF                                     5185
C
C      ENDIF                                     5186
C
C      ENDIF                                     5187
C
C      ENDIF                                     5188
C
C      ENDIF                                     5189
C
C      ENDIF                                     5190
C
C      ENDIF                                     5191
C
C      ENDIF                                     5192
C
C      ENDIF                                     5193
C
C      ENDIF                                     5194
C
C      ENDIF                                     5195
C
C      ENDIF                                     5196
C
C      ENDIF                                     5197
C
C      ENDIF                                     5198
C
C      ENDIF                                     5199
C
C      ENDIF                                     5200
C
C      ENDIF                                     5201
C
C      ENDIF                                     5202
C
C      ENDIF                                     5203
C
C      ENDIF                                     5204
C
C      ENDIF                                     5205
C
C      ENDIF                                     5206
C
C      ENDIF                                     5207
C
C      ENDIF                                     5208
C
C      ENDIF                                     5209
C
C      ENDIF                                     5210
C
C      ENDIF                                     5211
C
C      ENDIF                                     5212
C
C      ENDIF                                     5213
C
C      ENDIF                                     5214
C
C      ENDIF                                     5215
C
C      ENDIF                                     5216
C
C      ENDIF                                     5217
C
C      ENDIF                                     5218
C
C      ENDIF                                     5219
C
C      ENDIF                                     5220
C
C      ENDIF                                     5221
C
C      ENDIF                                     5222
C
C      ENDIF                                     5223
C
C      ENDIF                                     5224
C
C      ENDIF                                     5225
C
C      ENDIF                                     5226
C
C      ENDIF                                     5227
C
C      ENDIF                                     5228
C
C      ENDIF                                     5229
C
C      ENDIF                                     5230
C
C      ENDIF                                     5231
C
C      ENDIF                                     5232
C
C      ENDIF                                     5233
C
C      ENDIF                                     5234
C
C      ENDIF                                     5235
C
C      ENDIF                                     5236
C
C      ENDIF                                     5237
C
C      ENDIF                                     5238
C
C      ENDIF                                     5239
C
C      ENDIF                                     5240
C
C      ENDIF                                     5241
C
C      ENDIF                                     5242
C
C      ENDIF                                     5243
C
C      ENDIF                                     5244
C
C      ENDIF                                     5245
C
C      ENDIF                                     5246
C
C      ENDIF                                     5247
C
C      ENDIF                                     5248
C
C      ENDIF                                     5249
C
C      ENDIF                                     5250
C
C      ENDIF                                     5251
C
C      ENDIF                                     5252
C
C      ENDIF                                     5253
C
C      ENDIF                                     5254
C
C      ENDIF                                     5255
C
C      ENDIF                                     5256
C
C      ENDIF                                     5257
C
C      ENDIF                                     5258
C
C      ENDIF                                     5259
C
C      ENDIF                                     5260
C
C      ENDIF                                     5261
C
C      ENDIF                                     5262
C
C      ENDIF                                     5263
C
C      ENDIF                                     5264
C
C      ENDIF                                     5265
C
C      ENDIF                                     5266
C
C      ENDIF                                     5267
C
C      ENDIF                                     5268
C
C      ENDIF                                     5269
C
C      ENDIF                                     5270
C
C      ENDIF                                     5271
C
C      ENDIF                                     5272
C
C      ENDIF                                     5273
C
C      ENDIF                                     5274
C
C      ENDIF                                     5275
C
C      ENDIF                                     5276
C
C      ENDIF                                     5277
C
C      ENDIF                                     5278
C
C      ENDIF                                     5279
C
C      ENDIF                                     5280
C
C      ENDIF                                     5281
C
C      ENDIF                                     5282
C
C      ENDIF                                     5283
C
C      ENDIF                                     5284
C
C      ENDIF                                     5285
C
C      ENDIF                                     5286
C
C      ENDIF                                     5287
C
C      ENDIF                                     5288
C
C      ENDIF                                     5289
C
C      ENDIF                                     5290
C
C      ENDIF                                     5291
C
C      ENDIF                                     5292
C
C      ENDIF                                     5293
C
C      ENDIF                                     5294
C
C      ENDIF                                     5295
C
C      ENDIF                                     5296
C
C      ENDIF                                     5297
C
C      ENDIF                                     5298
C
C      ENDIF                                     5299
C
C      ENDIF                                     5300
C
C      ENDIF                                     5301
C
C      ENDIF                                     5302
C
C      ENDIF                                     5303
C
C      ENDIF                                     5304
C
C      ENDIF                                     5305
C
C      ENDIF                                     5306
C
C      ENDIF                                     5307
C
C      ENDIF                                     5308
C
C      ENDIF                                     5309
C
C      ENDIF                                     5310
C
C      ENDIF                                     5311
C
C      ENDIF                                     5312
C
C      ENDIF                                     5313
C
C      ENDIF                                     5314
C
C      ENDIF                                     5315
C
C      ENDIF                                     5316
C
C      ENDIF                                     5317
C
C      ENDIF                                     5318
C
C      ENDIF                                     5319
C
C      ENDIF                                     5320
C
C      ENDIF                                     5321
C
C      ENDIF                                     5322
C
C      ENDIF                                     5323
C
C      ENDIF                                     5324
C
C      ENDIF                                     5325
C
C      ENDIF                                     5326
C
C      ENDIF                                     5327
C
C      ENDIF                                     5328
C
C      ENDIF                                     5329
C
C      ENDIF                                     5330
C
C      ENDIF                                     5331
C
C      ENDIF                                     5332
C
C      ENDIF                                     5333
C
C      ENDIF                                     5334
C
C      ENDIF                                     5335
C
C      ENDIF                                     5336
C
C      ENDIF                                     5337
C
C      ENDIF                                     5338
C
C      ENDIF                                     5339
C
C      ENDIF                                     5340
C
C      ENDIF                                     5341
C
C      ENDIF                                     5342
C
C      ENDIF                                     5343
C
C      ENDIF                                     5344
C
C      ENDIF                                     5345
C
C      ENDIF                                     5346
C
C      ENDIF                                     5347
C
C      ENDIF                                     5348
C
C      ENDIF                                     5349
C
C      ENDIF                                     5350
C
C      ENDIF                                     5351
C
C      ENDIF                                     5352
C
C      ENDIF                                     5353
C
C      ENDIF                                     5354
C
C      ENDIF                                     5355
C
C      ENDIF                                     5356
C
C      ENDIF                                     5357
C
C      ENDIF                                     5358
C
C      ENDIF                                     5359
C
C      ENDIF                                     5360
C
C      ENDIF                                     5361
C
C      ENDIF                                     5362
C

```

```

4667 ALGORITHM DEPENDS ON TYPE OF ORGANISM
4668 IF (OTYPE .EQ. 1 .OR. OTYPE .EQ. 2) THEN
4669   CONDITIONING STAGE, ASSUME SIZE CONSTANT AND ALL AT 3 YEARS
4670   IF (I .EQ. 1) THEN
4671     DWK = AGESZ(J) / 7.
4672     SZDUP(J) = AGESZE(DWK)
4673     SZRF(J) = SZEUP(J)
4674   ENDIF
4675
4676 OTHERWISE, COMPUTE SIZE AT EACH DAY OF PERIOD BY CALLING
4677 FUNCTION AGESZE, THEN FIND AVG. SIZE OF PERIOD BY SUMMING
4678 USING THE TRAPEZOIDAL RULE
4679 ELSEIF (I .GT. 1) THEN
4680
4681   DCNTR = 0
4682   TSZE = 0.
4683   DO 100 D = (DAY-DAYSJ(I)-DAYS1+1.), (DAY-DAYS1+.49999)
4684     DWK = FLOAT(D) / 7.
4685     SZEUP(J) = AGESZE(DWK)
4686     TSZE = TSZE + SZEUP(J)
4687     DCNTR = DCNTR + 1
4688   CONTINUE
4689   COMPUTE SIZE DURING DAY BEFORE PERIOD BEGINS
4690   DWK = (DAY-DAYSJ(I)-DAYS1) / 7.
4691   VALUE = AGESZE(DWK)
4692
4693   INCREASE TOTAL BY 1/2 VALUE, DECREASE BY 1/2 LAST SUM TERM
4694   TSZE = TSZE + ((VALUE - SZEUP(J)) / 2.)
4695
4696   COMPUTE AVERAGE SIZE DURING J, WHERE DELTA T ALWAYS = 1
4697   SZRF(J) = TSZE / FLOAT(DCNTR)
4698   ENDIF
4699
4700   ELSE
4701     STOP 'ORGANISM TYPE WRONG IN SUBROUTINE SIZE'
4702   ENDIF
4703
4704   SUBROUTINE QUANT1(CINOW, JNOW, WKSNOW, MNOW, DAYNOW)
4705   RETURN
4706 END
4707
4708   SUBROUTINE QUANT2(CINOW, JNOW, WKSNOW, MNOW, DAYNOW)
4709   RETURN
4710
4711   SUBROUTINE QUANT3(CINOW, JNOW, WKSNOW, MNOW, DAYNOW)
4712   RETURN
4713
4714   SUBROUTINE QUANT4(CINOW, JNOW, WKSNOW, MNOW, DAYNOW)
4715   RETURN
4716
4717   SUBROUTINE ARGUE(MNOW, CINOW, CINOW, CINOW, CINOW, CINOW)
4718   ARGUE: MONTH THAT CURRENT PERIOD IS IN
4719   NUMBER OF BIVALE. EXISTING AT EACH DAY OF PERIOD
4720   SIZE OF BIVALE AT EACH DAY OF PERIOD
4721   LOOP COUNTER FOR TYPES OF INPUT N
4722
4723   TRANSLATE MNOW FOR CURRENT MONTH
4724   ARGUE: MONTH THAT CURRENT PERIOD IS IN
4725   NUMBER OF BIVALE. EXISTING AT EACH DAY OF PERIOD
4726   SIZE OF BIVALE AT EACH DAY OF PERIOD
4727   LOOP COUNTER FOR TYPES OF INPUT N
4728
4729   SUM OF TCDWK OVER ENTIRE PERIOD
4730   CELLS EATEN FOR ALL BIVALES AT THAT DAY OF PERIOD
4731   TRANSLATE WKSNOW FOR ELAPSED WEEKS
4732
4733
4734
4735
4736
4737
4738
4739
```

```

4731 C WKNOW ARGUE: ELAPSED WEEKS TO END OF PERIOD
4732 C -----
4733 C TRANSLATE ARGUMENTS INTO LOCAL VARIABLES
4734 C
4735 I = INNOW
4736 J = JNOW
4737 WKS = WKNOW
4738 M = MNOW
4739 DAY = DAYNOW

C ALGORITHM USED SHOULD DEPEND ON ORGANISM TYPE
IF (OTYPE .EQ. 1 .OR. OTYPE .EQ. 2) THEN
4740
C CONDITIONING STAGE REQUIRES DIFFERENT ALGORITHM
IF (I .EQ. 1) THEN
4741 STOP 'ERROR SUBR. QUANTS. CANT COMPUTE N5 FOR I=1'
4742
C OTHERWISE, COMPUTE TOTAL CELLS USED OVER PERIOD BY SUMMING
OVER THE PERIOD THE AMOUNT REQUIRED EACH DAY OF THE PERIOD.
4743 USE THE TRAPEZOIDAL RULE
4744 ELSEIF (I .GT. 1) THEN
4745
4746 DAY = DAYST1+4.99
4747
4748 C CELLS/DAY IS FUNCTION OF NUMBER OF ANIMALS AT THAT
4749 C DAY. ALSO MULTIPLY BY 10000. FOR PROPER EQUATION
4750 C
4751 DO 100 D = (DAY-DAYSJ(I)-DAYST1+1.), (DAY-DAYST1+.499)
4752 DWK = FLOAT(D) / 7.
4753
4754 DCNTR = 0.
4755 DO 100 D = (DAY-DAYSJ(I)-DAYST1+1.), (DAY-DAYST1+.499)
4756 DWK = FLOAT(D) / 7.
4757
4758 USE THE ALGORITHM CHOSEN IN READNS TO FIND CELLS/BIVALV
4759 IF (QEQQ5 .EQ. 1) THEN
4760 SZEDWK = AGENZE(DWK)
4761 CDWK = ((SZEDWK)**(1/EQQ5B(1))) *
4762 (EXP ((-EQQ5A(1)) / EQQ5B(1)) )
4763 ELSEIF (QEQQ5 .EQ. 2) THEN
4764 CDWK = ((DWK)**(1/EQQ5B(2))) *
4765 (EXP ((-EQQ5A(2)) / EQQ5B(2)) )
4766
4767 CONTINUE
4768
4769 TOTAL CELLS/DAY IS FUNCTION OF NUMBER OF ANIMALS AT THAT
4770 C DAY. ALSO MULTIPLY BY 10000. FOR PROPER EQUATION
4771 C
4772 NODWK = AGENO(DWK)
4773 TCDWK = CDWK * NODWK * 10000.
4774 TCELLS = TCELLS + TCDWK
4775 DCNTR = DCNTR + 1
4776
4777 DETERMINE QUANTITY OF DAY BEFORE PERIOD BEGINS, TO MORE
4778 CLOSELY APPROXIMATE THE INTEGRAL
4779 DNR = (DAY-DAYSJ(I)-DAYST1) / 7.
4780 IF (QEQQ5 .EQ. 1) THEN
4781 SZEDWK = AGENZE(DWK)
4782 VALUE = ((SZEDWK)**(1/EQQ5B(1))) *
4783 (EXP ((-EQQ5A(1)) / EQQ5B(1)) )
4784 ELSEIF (QEQQ5 .EQ. 2) THEN
4785 VALUE = ((DWK)**(1/EQQ5B(2))) *
4786 (EXP ((-EQQ5A(2)) / EQQ5B(2)) )
4787 ENDIF
4788 NODWK = AGENO(DWK)
4789 VALUE = VALUE * NODWK * 10000.
4790
4791 INCREASE SUM BY 1/2 VALUE, DECREASE BY 1/2 LAST SUM TERM
4792 TCELLS = TCELLS + ((VALUE - TCDWK) / 2.)
4793
4794 FILTERING RATE 100%, THEN INCREASE AMOUNT/PERIOD

```

```

FCELLS = TCELLSFILTER(I)
DO 200 T=1,NOFT(5)
    COMPUTE CELLS EACH TYPE
    CELLS = FCELLS * N5REQ
    COMPUTE QUANT/TYPE MUST
    AVAIL. FROM SEAWATER.
    TQUANJ(J,5,T) = CELLS

    COMPUTE AVERAGE QUANTILE
    PRINTED OUT IN ADDITION
    QUAOF(J,J,T) = TQUANJ(J

CONTINUE

        ENDIF
    ELSE
        STOP 'ORGANISM TYPE INCORRECT'
    ENDIF

RETURN
END

SUBROUTINE QUANT6(INOW,JNOW,WKSNN)
RETURN
END

SUBROUTINE COSTJ(INOW,JNOW,WKSNN)
-----
```

ORGNSM:

SUBROUTINE DOES NOT VARY WITH ORGANISM  
 VARIABLES DO NOT CHANGE

STRUCTURE: COMPUTES COSTS OF PERIOD  
 COSTS ACCUMULATED IN J  
 COSTS ACCUMULATED UP TO  
 COSTS PER ORGANISM UP TO

PARAMETERS  
 INCLUDE (IPAR1)  
 ARGUMENTS  
 INTEGER INOW, JNOW, MNOW  
 REAL WKSNOW, DAYNOWN  
 COMMON BLOCKS  
 INCLUDE (ICMAIN)  
 INCLUDE (ICBLK1)  
 INCLUDE (ICBLK0)  
 INCLUDE (ICBLK1)  
 LOCAL VARIABLES

```

INTEGER I, J, M, N, T
REAL WKS, DAY, CSTOFJ
C NAME DESCRIPTION OF LOCAL VARIAB AND ARGUMENTS
C -----
C COST OF TYPE T IN J (WRITTEN OVER)
C DAY TRANSLATE DAYNOW FOR ELAPSED DAYS
C DAYNOW ARGUE: ELAPSED DAYS TO END OF PERIOD
C I TRANSLATE JNOW FOR CURRENT STAGE
C INOW ARGUE: CURRENT STAGE THAT J IS IN
C J NOW TRANSLATE JNOW FOR CURRENT PERIOD
C JNOW ARGUE: CURRENT PERIOD
C M TRANSLATE MNOW FOR CURRENT MONTH
C MNOW ARGUE: MONTH THAT CURRENT PERIOD IS IN
C N LOOP COUNTER FOR INPUTS
C T LOOP COUNTER FOR TYPES
C WKS TRANSLATE WKNOW FOR ELAPSED WEEKS
C WKNOW ARGUE: ELAPSED WEEKS TO END OF PERIOD
C

C COMPUTE COST OF EACH INPUT, N, ONE AT A TIME
C
C I = INOW
C J = JNOW
C WKS = WKNOW
C M = MNOW
C DAY = DAYNOW
C
C IF (OPTION .EQ. 1 .AND. J .EQ. 1) THEN
C   CSUPJ(J) = CSOP1
C   CSUPJ(J) = CSOP1
C   CSNORJ(J,N) = 0.
C   CSNUPJ(J,N) = 0.
C ENDIF
C
C COST OF EACH TYPE, T, OF EACH INPUT, N, DURING A PERIOD, J
C IF (I .GE. BEGIN) THEN
C   CSNORJ(J,N) = 0.
C   DO 200 T=1,NOOFT(N)
C     CSTOFJ = TQUANJ(J,N,T) * PRIGHT(N,T,MOFJ(J))
C     CSNORJ(J,N) = CSNORJ(J,N) + CSTOFJ
C   CONTINUE
C
C IE, LET THIS VARIABLE EQUAL CALCULATIONS FOR LAST DAY
C COST OF ALL INPUTS USED IN A PERIOD
C   CSTOFJ(J) = CSOFJ(J) + CSNORJ(J,N)
C
C IF (J .EQ. 1 ) THEN
C   CSNUPJ(J,N) = CSNORJ(J,N)
C   CSUPJ(J) = CSOFJ(J)
C ELSEIF (J .NE. 1 ) THEN
C   CSNUPJ(J,N) = CSNORJ(J,N) + CSNUPJ(J-1,N)
C   CSUPJ(J) = CSOFJ(J) + CSUPJ (J-1)
C ENDIF
C
C 100 CONTINUE
C
C COMPUTE COST PER ORGANISM
C   AVERAGE PER DAY OVER THE PERIOD
C   CSOOFJ(J) = (CSOFJ(J) / ONOOFJ(J)) / DAYS(J,IOFJ(J))
C
C UP TO END OF PERIOD
C   IF (ONOOFJ(J) .EQ. 0) THEN
C     CSOUPJ(J) = 0.
C   ELSE
C     CSOUPJ(J) = CSUPJ(J) / ONOUPJ(J)
C   ENDIF
C
C JNOW ARGUE: CURRENT PERIOD AND UP TO END OF PERIOD
C DAYNOW ARGUE: CURRENT MONTH AND UP TO END OF PERIOD
C MNOW ARGUE: CURRENT STAGE AND UP TO END OF PERIOD
C T ARGUE: CURRENT TYPE AND UP TO END OF PERIOD
C WKS ARGUE: CURRENT WEEKS AND UP TO END OF PERIOD
C
C SUBROUTINE WRTOUT(MBEGIN, LASTJ,WRT)
C
C I = INOW
C J = JNOW
C WKS = WKNOW
C M = MNOW
C DAY = DAYNOW
C
C IF (OPTION .EQ. 1 .AND. J .EQ. 1) THEN
C   CSUPJ(J) = CSOP1
C   CSUPJ(J) = CSOP1
C   CSNORJ(J,N) = 0.
C   CSNUPJ(J,N) = 0.
C ENDIF
C
C COST OF EACH INPUT USED IN EACH PERIOD
C   -STAGE AND PERIOD INFORMATION
C   -QUANTITY OF EACH INPUT USED DURING EACH PERIOD AND UP TO A PERIOD
C   -COST OF EACH INPUT USED DURING EACH PERIOD AND UP TO A PERIOD
C   -COST OF PRODUCING AN ORGANISM UP TO THE END OF A PERIOD
C   -COST OF PRODUCING AN ORGANISM UP TO THE END OF A PERIOD
C
C STRUCTURE: WRITES OUT INFORMATION COMPUTED FOR A BATCH BEGUN ON THE
C FIRST OF EACH MONTH OF THE PRODUCTION SEASON.
C
C THIS INFORMATION INCLUDES:
C   -STAGE AND PERIOD INFORMATION
C   -QUANTITY OF EACH TYPE OF EACH INPUT USED IN EACH PERIOD
C   -COST OF EACH INPUT USED DURING EACH PERIOD AND UP TO A PERIOD
C   -COST OF PRODUCING AN ORGANISM UP TO THE END OF A PERIOD
C
C 1. PARAMETERS
C   INCLUDE (IPAR1)
C 2. ARGUMENTS
C   INTEGER LASTJ, WRT, WRT2
C   CHARACTER * 10 MBEGUN
C 3. COMMON BLOCKS
C   INCLUDE (ICMAIN)
C   INCLUDE (ICBLK1)
C   INCLUDE (ICBLK0)
C   INCLUDE (ICUNIT)
C 4. LOCAL VARIABLES
C   INTEGER I, J, M, N, T, W
C
C NAME DESCRIPTION OF LOCAL VARIABLES AND ARGUMENTS
C -----
C I LOOP COUNTER FOR STAGE
C J LOOP COUNTER FOR PERIODS
C LASTJ LAST J PERIOD SUCCESSFULLY COMPUTED
C M LOOP COUNTER FOR MONTHS
C MBEGUN MONTH THAT CURRENT BATCH WAS BEGIN
C N LOOP COUNTER FOR INPUTS
C T LOOP COUNTER FOR TYPES
C WRT ARGUMENT: UNIT NUMBER
C W TRANSLATE ARGUMENT UNIT NUMBER TO LOCAL WRITE VARIABLE
C WRT2 ARGUE: TRANSLATE UNIT WRITE TO NEXT LEVEL SUBROUTINE
C
C -----
```



4 T46,'IN PERIOD/DAY',T61,'AT END PERIOD')  
 36000 FORMAT(' ',I3,T11,I1,T16,G14.7,T31,G14.7,T46,G14.7,T61,G14.7)  
 33000 FORMAT(' ',I2,T8,I1,T11,A10,T25,I2,T31,G14.7,T46,G14.7,T61,  
 33100 FORMAT(' ',I2,T8,I1,T11,A10,T26,I2,T31,G14.7,T46,'TIME PAST ',  
 2 'MONTHS OF ',  
 3 'PRODUCTION. CANT PRODUCE ',A,' OF THIS AGE IF BEGUN IN ',A)  
 40000 FORMAT('0'/' ','COST OF INPUTS IN A PERIOD:'//+',27(' '\_'))  
 41000 FORMAT(' ','PERIOD',T9,'STAGE',T16,6(A10,5X))  
 42000 FORMAT(' ',I3,T11,I1,4(T16,7(G14.7,1X)/' '))  
 42100 FORMAT(' ',I3,T11,I1,T16,'TIME PAST PRODUCTION MONTHS')  
 50000 FORMAT('0'/' ','COST OF INPUTS UP TO END OF A PERIOD',//+,37(' '\_'))  
 70000 FORMAT('0'/' ','QUANTITY OF INPUT USED PER PERIOD://+,34(' '\_'))  
 71000 FORMAT(' '/',A10,IX,A,:'/ ',17(' '-'))  
 72000 FORMAT(' ','PERIOD',T9,'STAGE',T16,10(A10,5X))  
 72100 FORMAT(' ','PERIOD',T9,'STAGE')  
 73000 FORMAT(' ',I3,T11,I1,T16,10(G14.7,1X))  
 90000 FORMAT(' ',3(' -'),T6,4(' ',''),T11,14(',-'),T26,4(',-'),T31,14(',-'),  
 2 T46,14(' -'),T6,14(' -'),T76,14(' -'),T91,14(' -'),T106,14(' -'))  
 93000 FORMAT(' ',6(' -'),T9,6(' -'),T16,14(' -'),T31,14(' -'),T46,14(' -'),  
 2 T61,14(' -'))  
 95000 FORMAT(' ',6(' -'),T9,6(' -'),T16,14(' -'),T31,14(' -'),T46,14(' -'),  
 2 T61,14(' -'),T76,14(' -'),T91,14(' -'))  
 5154 NAME DESCRIPTION OF LOCAL VARIABLES AND ARGUMENTS  
 5155 C  
 5156 C  
 5157 C J LOOP COUNTER FOR PERIODS  
 5158 C LASTJ ARGUE: LAST PERIOD THAT COMPUTATIONS WERE DONE FOR  
 5159 C MBEGUN ARGUE: MONTH IN WHICH CURRENT BATCH IS BEGUN  
 5160 C WRT ARGUE: UNIT NUMBER FOR WRITING  
 5161 C W UNIT NUMBER TO WRITE  
 5162 C  
 5163  
 5164 W = WRT  
 5165 SUBROUTINE WOUTN1(NBEGUN,LASTJ,WRT)  
 5166 RETURN  
 5167 END  
 5168 C WRITE INFORMATION SPECIFIC TO THIS PROGRAM  
 5169 DO 100 J = 1,LASTJ  
 5170 WRITE(W,10000) J, MOF(J), DAYUP(J), AGEUP(J), ONOOFJ(J),  
 5171 2 SZEOPJ(J), CSOOFJ(J), CSUPJ(J), CSOUPJ(J), CSNDFJ(J),  
 5172 3 CSNDPJ(J), TQUANJ(J,5,5), TQUANJ(J,5,6),  
 5173 4 QUAOFJ(J,5), QUAOFJ(J,6)  
 5174 100 CONTINUE  
 5175  
 5176 10000 FORMAT(' ',I2,1X,I2,T10,SE12.5/' ',T10,SE12.5)  
 5177  
 5178

```

5243   C FUNCTION RETURNED TO SUBROUTINES QUANTS, SIZE
5244   C CURRENT ORGANISM: BIVALVES
5245   C STRUCTURE: RELATES SIZE TO AGE IN TERMS OF WEEKS
5246
5247   C 1. PARAMETERS
5248   C     INCLUDE (IPAR1)
5249   C 2. ARGUMENTS
5250   C     REAL WEEKS
5251   C 3. COMMON BLOCKS
5252   C     INCLUDE (ICMAIN)
5253   C     INCLUDE (ICBLKO)
5254
5255   C NAME DESCRIPTION OF LOCAL VARIABLES AND ARGUMENTS
5256
5257   C
5258   C WEEKS ARGUE: ELAPSED WEEKS UP TO THE END OF PERIOD J
5259   C
5260
5261   C ORGNSM:
5262   C FUNCTION VARIES WITH ORGANISM
5263   C FUNCTION REQUIRED FOR SUBROUTINE NUMBRB
5264   C CURRENT ORGANISM: BIVALVES
5265
5266   C STRUCTURE: RELATES SIZE TO AGE IN TERMS OF WEEKS
5267   C 1. PARAMETERS
5268   C     INCLUDE (IPAR1)
5269   C     INCLUDE (ICBLKO)
5270   C     INCLUDE (ICBLKO)
5271   C     RETURN
5272   C     END
5273
5274
5275
5276
5277
5278
5279
5280
5281
5282
5283
5284
5285
5286
5287
5288
5289
5290
5291
5292
5293
5294
5295
5296
5297
5298
5299
5300
5301
5302
5303
5304
5305
5306
5307
5308
5309
5310
5311
5312
5313
5314
5315
5316
5317
5318
5319
5320
5321
5322
5323
5324
5325
5326
5327
5328
5329
5330
5331
5332
5333
5334
5335
5336
5337
5338
5339
5340
5341
5342
5343
5344
5345
5346
5347
5348
5349
5350
5351
5352
5353
5354
5355
5356
5357
5358
5359
5360
5361
5362
5363
5364
5365
5366
5367
5368
5369
5370

```

81

```

5435 C K LOOP COUNTER FOR PROGRAM RUNNING OPTIONS
5436 C M LOOP COUNTER FOR MONTHS
5437 C N LOOP COUNTER FOR INPUTS
5438 C T LOOP COUNTER FOR TYPES OF INPUT N
5439 C

C VARIABLES THAT SHOULD NOT CHANGE
C -----
C NAME OF MONTHS
C DATA (NAMEM(N), N=1,12) /'JANUARY','FEBRUARY','MARCH','APRIL',
C   2   /'MAY','JUNE','JULY','AUGUST','SEPTEMBER',/ 5374
C   3   /'OCTOBER','NOVEMBER','DECEMBER',/ 5375

C UNIT NUMBERS FOR ALL BUT OUTPUT FILES/ AND TO INITIATE VARIABLES
C -----
C DATA WF, RL, WI, RF / 3,5,6,8 /
C DATA (QUTQN(N), N=1,NOOFN) / NOOFN * 'N' /
C
C VARIABLES THAT CHANGE WITH CAPABILITIES OF THE PROGRAM'S OPTIONS
C -----
C NUMBER OF TYPES,T, OF EACH INPUT,N
C DATA (NOOFT(N), N=1,NOOFN) / 1,1,1,1,1,1,1 /
C
C OPTION INFORMATION: 0=AVAILABLE, 1=NOT AVAILABLE
C   OPTIONS 1-3 FOR RUNNING PROGRAM
C   DATA (OPQOK(K), K=1,3) / 0,0,1 /
C
C TO COMPUTE QUANTITY FOR EACH INPUT,N, IN EACH STAGE,J
C   DATA (OPNOK(I,1), I=1,NOOFI) / NOOFI * 1 /
C   DATA (OPNOK(I,2), I=1,NOOFI) / NOOFI * 1 /
C   DATA (OPNOK(I,3), I=1,NOOFI) / NOOFI * 1 /
C   DATA (OPNOK(I,4), I=1,NOOFI) / NOOFI * 1 /
C   DATA (OPNOK(I,5), I=1,NOOFI) / 1,0,0,0,0 /
C   DATA (OPNOK(I,6), I=1,NOOFI) / NOOFI * 1 /
C
C TO COMPUTE THE PRICE OF INPUT,N
C   DATA (OPNSOK(N), N=1,NOOFN) / 1,1,1,1,0,1 /
C
C TO CHANGE THE NUMBER OF TYPES,T, OF EACH INPUT,N
C   DATA (OPTOK(N), N=1,NOOFN) / 1,1,0,1,0,1 /
C
C TO PRINT OUT ADDITIONAL INFORMATION ABOUT PRODUCTION INPUTS
C   DATA (OUTON(N), N=1,NOOFN) / 1,1,1,1,0,1 /
C
C ORGNSM:
C -----
C SUBROUTINE VARIES WITH ORGANISM
C SUBROUTINES THAT MUST BE DATA INITIALIZED:
C   C NAME(N)
C   C NAMEU(N)
C   C NAMEI(N)
C
5371 C NAMET(N,T)
5372 C NAME(O)
5373 C CURRENT ORGANISM: BIVALVES
5374 C STRUCTURE: INITIALIZE VARIABLES IN COMMON BLOCK, ICBLKO
5375 C

5376 C 1. PARAMETERS
5377 C   INCLUDE ('IPARI')
5378 C   3. COMMON BLOCKS
5379 C   INCLUDE ('ICBLKO')
5380 C   4. LOCAL VARIABLES
5381 C   INTEGER EQ, I, N, O, T
5382 C
5383 C   NAME      DESCRIPTION OF LOCAL VARIABLES
5384 C   EQ       LOOP COUNTER FOR FEEDING EQUATIONS, INPUT 5
5385 C   I        LOOP COUNTER FOR STAGES
5386 C   N        LOOP COUNTER FOR INPUTS
5387 C   O        LOOP COUNTER FOR ORGANISM TYPES
5388 C   T        LOOP COUNTER FOR TYPES OF INPUT N
5389 C
5390 C
5391 C
5392 C
5393 C   VARIABLES REQUIRED BUT THAT CHANGE WITH ORGANISM:
5394 C   DATA NAME / 'BIVALVE' /
5395 C
5396 C   DATA (NAME(N), N=1,NOOFN) / 'ELECTRICITY','FUEL OIL','LABOR',
5397 C   2   /'MATERIALS','ALGAE','CUCH' /
5398 C   DATA (NAMEU(N), N=1,NOOFN) / 'KWHR','LITRE','HOUR','UNIT',
5399 C   2   /'CELL','LITRE' /
5400 C   DATA (NAMEU(N), N=1,NOOFN) / 'CONDITION','SPAWN','LARVAE',
5401 C   2   /'SETTING','JUVENILE' /
5402 C   DATA (NAME(O), O=1,NOOFO) / 'OYSTERS','CLAMS' /
5403 C
5404 C   VARIABLES RELATING TO ORGANISM DEPENDENT FUNCTIONS:
5405 C   DATA EQNO(1) /'POPULATION = POPULATION(INITIAL) * [(WEEKS+1)**A]' /
5406 C   DATA EQNO(1) /'POPULATION = POPULATION(INITIAL) * [((WEEKS+1)**A)**B]' /
5407 C   DATA EQNOAF / - .255068 /
5408 C
5409 C
5410 C   DATA (EQ05(EO), EQ=1,2) /
5411 C   2   /'NATURAL LOG(SIZE IN MM) = A + B * NAT LOG(10,000 CELLS)',/
5412 C   2   /'NATURAL LOG(AGE IN WKS) = A + B * NAT LOG(10,000 CELLS)' /
5413 C
5414 C   DATA (EQ05AF(EO), EQ=1,2) / -.631048, -.330940 /
5415 C   DATA (EQ05BF(EO), EQ=1,2) / .338426, .344505 /
5416 C
5417 C   DATA EQSZE(1) / 'SIZE = (A * AGE) + (B * AGE SQUARED)' /
5418 C
5419 C   DATA EQSZAF / .737 /
5420 C   DATA EQSZBF / -.002532 /
5421 C
5422 C
5423 C
5424 C
5425 C
5426 C
5427 C
5428 C
5429 C
5430 C
5431 C
5432 C
5433 C
5434 C
5435 C
5436 C
5437 C
5438 C
5439 C
5440 C
5441 C
5442 C
5443 C
5444 C
5445 C
5446 C
5447 C
5448 C
5449 C
5450 C
5451 C
5452 C
5453 C
5454 C
5455 C
5456 C
5457 C
5458 C
5459 C
5460 C
5461 C
5462 C
5463 C
5464 C
5465 C
5466 C
5467 C
5468 C
5469 C
5470 C
5471 C
5472 C
5473 C
5474 C
5475 C
5476 C
5477 C
5478 C
5479 C
5480 C
5481 C
5482 C
5483 C
5484 C
5485 C
5486 C
5487 C
5488 C
5489 C
5490 C
5491 C
5492 C
5493 C
5494 C
5495 C
5496 C
5497 C
5498 C
5499 C
5490 C
5491 C
5492 C
5493 C
5494 C
5495 C
5496 C
5497 C
5498 C
5499 C

```

```

0001 C FILE NAME : ICM5S
0002 C ORGANISM: VARIABLES SPECIFIC TO ORGANISM
0003 C CURRENT ORGANISM: BIVALVES
0004 C TO INCLUDE COMMONS USED IN PRICES SUBROUTINE
0005 C VARIABLE NAMES ARE IDENTICAL IN ALL PRICES SUBROUTINES.
0006 C PARAMETERS USED IN PRICES
0007 INTEGER NUMGOD, MAXLAB
0008 PARAMETER (NUMGOD=2, MAXLAB=6)
0009 COMMON / CN55 /
0010      I BEGPRO, ENDPRO, NUMLAB, NUMSTIC,
0011      R BLBLIF, COMEFF, HPCOM, HPPMP, HRSCLM, NUMTUB,
0012      R PKWHR, PMPGPM, POIL, WATAIR, WATBLB,
0013      R ALLAB, BBLRM(MAXT), CALIN(MAXT), TCELL(MAXT), CPPRML(MAXT),
0014      R DAYSTG(MAXT), DISSTG(MAXT) HRSAUT(MAXT), H20TNIK(MAXT), 0014
0015      R INQNIK(MAXT), LFOOD(MAXT), LITWHR(MAXT), NUMAIR(MAXT), NUMTNIK(MAXT),
0016      R STGRML(MAXT), TALLAB(MAXT), TALLL(MAXT), TCALIN(MAXT),
0017      R TKWHL(MAXT), TLITRE(MAXT), TOLL(MAXT), TTKWHR(MAXT),
0018      R T20TNIK(MAXT), TTIOIL(MAXT), NUTLTR(MAXT), TCKEL(MAXT),
0019      R ALFOOU(MAXT), TCELL(MAXT),
0020      R HRLABS(6,MAXT), PLABOR(6), TCLAYT(6), TLATAY(6,MAXT),
0021      R CINP(4,MAXT), PGOD(5), TGOTY(5,MAXT), TIGOTY(5,MAXT),
0022      R HRSAT(MAXM,MAXT), TCLITR(MAXT),
0023      C NAMAG(MAXT), MONAVA(MAXT),
0024      C DATE2, QWRITS$5
0025      INTEGER BEGPRO, ENDPRO, NUMLAB, NUMSTIC,
0026      REAL BLBLIF, COMEFF, HPCOM, HPPMP, HRSCLM, NUMTUB, PKWHR,
0027      R PMPGPM, POIL, WATAIR, WATBLB,
0028      R ALLAB, BBLRM, CALIN, TCELL, CPPRML, DAYSTG, DISSTG, HRSAUT,
0029      R HRSBLB, H20TNIK, INQNIK, LFOOD, LITWHR, NUMAIR, NUMTNIK,
0030      R STGRML, TALLAB, TALLL, TCALIN, TKWHL, TLITRE, TOIL,
0031      R TKWHL, TTWKHR, TTIOIL, TNLTR, TUBTNK,
0032      R ALFOOU, TCELL,
0033      R HRLABW, PLABOR, TCLAYT, TLATAY, TLITR,
0034      R PGOD, TGOTY, TIGOTY, CIMP, TCIMP, TCINPL, HRLABS,
0035      R HSAIR, TCLITR
0036      CHARACTER*10 NAMALG, MONAVA
0037      CHARACTER*10 DATE2
0038      CHARACTER*40 QWRITS$5
0039
0040      C FILE NAME : ICBLK1
0041      C ORGANISM: VARIABLES DON'T CHANGE WITH ORGANISM, BUT WITH MACHINE CAPABILITY
0042      C TO INCLUDE COMMONS NECESSARY TO RUN BIVALVE PROGRAM SKELETON
0043      COMMON / CBLK1 /
0044      I ANYNOK, BEGINJ, BEGINJ, OTYPE, ENDJ,
0045      I ENDM, IOFJ, JPERI, MOFJ, M1, M2, OPTION, PASEDM, TOTALI,
0046      I AGEPJ, ONOBEG, ONOOFJ, ONOUPJ, CSOOFJ, CSNOFJ, 0007
0047      R CSNUPJ, CSOOFJ, CSUPJ, CSUP1, CSUPJ, 0008
0048      R DAYOJ, DAYP1, DAYSI, DAYSJ, PRICET, SZEOFJ, 0009
0049      R TQUANI, TQUANJ, WKSUPJ, 0010
0050      C DATE, NAMEF1, NAMES, NQBYM, OPNNEW, QOPNS$, QOUTQN
0051      C INTEGERT
0052      I ANYNOK, BEGINJ, BEGINJ, OTYPE, ENDJ,
0053      I ENDM, IOFJ, JPERI, MOFJ, M1, M2, OPTION, PASEDM, TOTALI
0054      I REAL
0055      R AGEUPJ, ONOBEG, ONOOFJ, ONOUPJ, CSOOFJ, CSNOFJ, 0013
0056      R CSNUPJ, CSOOFJ, CSUP1, CSUPJ, DAYUPJ, 0014
0057      R DAYOJ, DAYSI, DAYSJ, PRICET, SZEOFJ, SZEUPJ, 0015
0058      R TQUANI, TQUANJ, WKSUPJ, 0016
0059      C CHARACTER * 10 NAMEF1
0060      C CHARACTER * 1 NQBYM, N$BYM, QOPNS$, OPNNEW, QOUTQN
0061      C DIMENSION
0062      I ANYNOK(NOOFN), BEGINJ(NOOFN), IOFJ(MAXJI), JPERI(NOOFI),
0063      I MOFJ(MAXJI), PASEDM(MAXJI),
0064      I AGEUPJ(MAXJI), ONOOFJ(MAXJI), ONOUPJ(MAXJI), CSOOFJ(MAXJI),
0065      I CSNUPJ(MAXJI), CSNOFJ(MAXJI, NOOFN), 0024
0066      R CSOOFJ(MAXJI), DAYUPJ(MAXJI), DAYSI(NOOFT),
0067      R CSNUPJ(MAXJI), CSUP1(NOOFN), DAYOJ(NOOFT), 0025
0068      R CSUPJ(MAXJI), DAYUPJ(MAXJI), DAYSI(NOOFN, MAXT, MAXM),
0069      R DAYOJ(NOOFT), ENDJ(NOOFT), PRICET(NOOFN, MAXT, MAXM),
0070      R SZEOFJ(MAXJI), SZEUPJ(MAXJI), 0026
0071      R TQUANI(NOOFT), NOOFN, MAXT, MAXM, 0027
0072      R TQUANJ(MAXJI, NOOFN), TQUANJ(MAXJI, NOOFN, MAXT),
0073      R NQBYM(NOOFN), N$BYM(NOOFN), OPNNEW(NOOFI, NOOFN), 0028
0074      C QOUTQN(NOOFN)
0075      C
0076      C FILE NAME : ICBLK2
0077      C ORGANISM: VARIABLES DON'T CHANGE WITH ORGANISM, BUT WITH MACHINE CAPABILITY
0078      C USED FOR INCLUSION OF COMMON BLOCKS VARIABLES INITIALIZED IN BLOCK
0079      C DATA SUBPROGRAM
0080      COMMON / CBLK2 /
0081      I NOOFT, OPTOK, OPNSOK, OPQOK, OPNOK, OUTQN,
0082      C NAME
0083      C CHARACTER * 10 NAMEM
0084      C DIMENSION
0085      I NOOFT(NOOFN), OPTOK(NOOFN), OPNSOK(NOOFN), OPQOK(3),
0086      I OPNOK(NOOFI, NOOFN), OUTQN(NOOFN),
0087      C NAMEM(MAXM)
0088
0089      C FILE NAME : ICBLK3
0090      C ORGANISM: VARIABLE DON'T CHANGE
0091      C FOR INCLUSION OF COMMON BLOCK DATA INITIALIZED, ORGANISM SPECIFIC
0092      COMMON / CBLK3 /
0093      I EQNOA, EQNOAF, EQQ5AF, EQQ5BF, EQQ5BF,
0094      R EQSZA, EQSZA, EQSZA, EQSZA, EQSZA, EQSZA, 0001
0095      C EQNO, EQQ5, EQSZE,
0096      C NAME, NAMEO, NAMEI, NAMEM, NAMEU
0097      REAL EQNOA, EQNOAF, EQQ5AF, EQQ5BF, EQQ5BF,
0098      R EQSZA, EQSZA, EQSZA, EQSZA, EQSZA, EQSZA, 0002
0099      CHARACTER * 5 NAMEU
0100      CHARACTER * 10 NAME, NAMEO, NAMEI, NAMEM, NAMEU
0101      CHARACTER * 62 EQNO, EQQ5, EQSZE
0102      C EQQA(2), EQQ5AF(2), EQQ5BF(2), EQQ5BF(2),
0103      C EQNO(1), EQQ5(2), EQSZE(1),
0104      C NAMEO(NOOFO), NAMEI(NOOFI), NAMEU(NOOFN),
0105      C NAMEU(NOOFN, MAXT),
0106      C
0107      C FILE NAME : ICBLK4
0108      C ORGANISM: VARIABLES DON'T CHANGE WITH ORGANISM, BUT WITH MACHINE CAPABILITY
0109      C ALL UNIT NUMBER FOR READ, WRITE, AND OUTPUT FILES
0110      COMMON / CBLK4 /
0111      I RF, RI, WF, WI, WO, WON
0112      C
0113      C
0114      C
0115      C
0116      C
0117      C
0118

```

```

C FILE NAME: ICORG
C ORGNSM: VARIABLES SPECIFIC TO ORGANISM
C CURRENT ORGANISM: BIVALVE
C TO INCLUDE COMMONS SPECIFIC TO ORGANISM
C COMMON / CORG /
I
I QEQQ5,
R DASY11, FILTER, NO011, N5AVAL, N5REQ, QUAOFJ, SEACEL,
C N5ABYM, SEABTM
INTEGER QEQQ5
REAL DASY11, FILTER, NO011, N5AVAL, N5REQ, QUAOFJ, SEACEL,
CHARACTER*1 N5ABYM, SEABYM
DIMENSION
R FILTER(MAXJ), N5AVAL(MAXJ,MAXM),
R N5REQ(MAXJ,MAXJ), QUAOFJ(MAXJ,MAXJ), SEACEL(MAXJ,MAXM)

```

## APPENDIX B

This appendix contains the data used in the base run of the simulation model and the output for the base run. All or portions of this information is printed at the user's request. However, what appears in the following pages is an edited version of what is actually printed by the program. Data were not available to estimate all component costs of bivalve seed. Thus, to save space, only that information pertaining to algae and total feed costs is reported. In addition, the algae price computations are reported only for one month, although algae cost differences by month are reflected in the bivalve feed costs reported in subsequent sections of the output.

BIVALVE COST SIMULATION MODEL: INPUTS ENTERED

DATE: MARCH 30, 1988

OPTION DATA ENTERED

NAME OF FILE USED: 1201 8DATA

OPTION CHOSEN: 1

COST UP TO FIRST STAGE OF INTEREST IS ENTERED  
USER CHOOSES INPUTS TO COMPUTE IN REMAINING STAGES

THE FIRST STAGE OF INTEREST IS THE SPAWN STAGE

IF C, THE PROGRAM COMPUTES THE QUANTITY

IF E, THE USERS ENTERS THE QUANTITY

IF -, THE QUANTITY IS NEITHER ENTERED NOR COMPUTED

INPUT	CONDITION	SPAWN	LARVAE	SETTING	JUVENILE
ELECTRICITY	-	E	E	E	E
FUEL OIL	-	E	E	E	E
LABOR	-	E	E	E	E
MATERIALS	-	E	E	E	E
ALGAE	-	C	C	C	C
CULCH	-	E	E	E	E

THE NUMBER OF EACH SECTION FOR WHICH DATA MUST BE ENTERED

SECTION 1: GENERAL DATA

SECTION 2: BIVALVE DATA

SECTION 7: ALGAE DATA

BIVALVE COST SIMULATION MODEL: ALGAE PRICE COMPUTATIONS

DATE: MARCH 30, 1988 INPUTS ENTERED

1. ALGAE THESIS RESULTS

\*PRICE INFORMATION

2. OF ELECTRICITY	\$ 0.1000	PER KWHR
3. OF FUEL OIL	\$ 0.2990	PER LITR
4. OF LIGHTBULBS	\$ 2.0000	PER BULB
4. OF NUTRIENT SOL.	\$ 1.0000	PER LITR
5. # OF LABOR TYPES:	3	
6. PRICE OF TYPE1	\$ 6.7310	PER HR
6. PRICE OF TYPE2	\$ 5.5290	PER HR
6. PRICE OF TYPE3	\$ 4.8080	PER HR
8. WATT RATING OF AIR CONDIT.	1050.0	WATT
9. WATT RATING OF LITEBULBS	40.0	WATT
10. LIFETIME OF LITEBULBS	15000.0	HRS
11. WATT RATING OF AUTOCLAVE	2000.0	WATT
12. COMPRESSOR HORSEPOWER	1.0	HP
13. HOURS COMPRESSOR ON/DAY	24.0	HRS
14. TOTAL # COMPRESSOR TUBES	74.	TUBE
15. EFF. RATING OF COMPRESSOR	0.850	
16. EFF. RATING OF WATER PUMP	0.450	GPM
17. GPM RATING OF WATER PUMP	40.0	HP
18. HP RATING OF WATER PUMP	2.5	HP
19. HOURS OF LABOR/WEEK		
OF TYPE1	11.5	HRS
OF TYPE2	3.0	HRS
OF TYPE3	0.0	HRS

## \*STAGE INFORMATION

20. NUMBER OF STAGES: 6

	1STAGE	2STAGE	3STAGE	4STAGE	5STAGE	6STAGE
21. NUTRIENT/LITRE H2O	0.100E-02	0.225E-02	0.101E-02	0.269E-03	0.147E-03	0.587E-04
22. NUMBER OF TANKS	6.0	5.0	5.0	5.0	5.0	2.0
23. NUMBER OF STAGES PER ROOM	3.0	3.0	3.0	1.0	1.0	1.0
24. DAYS TO REACH Maturity	14.0	7.0	7.0	7.0	7.0	7.0
25. CELLS PER MILLITRE - DENSITY	0.300E+06	0.300E+06	0.300E+06	0.400E+06	0.650E+06	0.600E+06
26. LITRES WATER ADDED PER TANK	0.500E-02	1.24	13.8	743.	0.341E+04	0.341E+05
27. LITRES OF INNOCULATION ADDED/TANK	0.000E+00	0.500E-02	0.200	14.0	379.	0.379E+04
28. AVG. LITRES UNCONTROLLED DISCARD	0.500E-02	2.08	0.000E+00	379.	0.170E+04	0.852E+04
29. NUMBER OF AIR TUBES PER TANK	1.00	1.00	1.00	2.00	4.00	5.00
30. NUMBER OF LIGHTBULBS PER ROOM	10.0	10.0	10.0	12.0	8.0	8.0
31. HOURS LIGHTS ON PER DAY PER ROOM	24.00	24.00	24.00	24.00	2.00	0.50
32. HOURS AUTOCLAVE USED	0.17	0.50	0.84	0.00	0.00	0.00
33. NUMBER OF AIR CONDIT. PER ROOM	1.0	1.0	1.0	1.0	0.0	0.0
34. HRS OF LABOR PER STAGE OF TYPE1 OF TYPE2 OF TYPE3	1.50 0.00 0.00	1.50 0.00 0.00	1.50 0.00 0.00	4.00 0.00 0.00	2.00 2.00 0.00	2.00 2.00 2.00
35. NAME OF EACH ALGAE STAGE	PRIMARY1	PRIMARY2	PRIMARY3	SECONDARY	TERTIARY	SOLAR
*MONTH INFORMATION						
36. MONTH BEGIN PRODUCTION	1 JANUARY					
37. MONTH END PRODUCTION	12 DECEMBER					
38. HOURS AIR CONDIT. ON PER DAY/ROOM	1STAGE	2STAGE	3STAGE	4STAGE	5STAGE	6STAGE
1 JANUARY	0.00	0.00	0.00	0.00	0.00	0.00
2 FEBRUARY	0.00	0.00	0.00	0.00	0.00	0.00
3 MARCH	0.00	0.00	0.00	0.00	0.00	0.00
4 APRIL	4.40	4.40	4.40	4.40	0.00	0.00
5 MAY	8.80	8.80	8.80	8.80	0.00	0.00
6 JUNE	11.50	11.50	11.50	11.50	0.00	0.00
7 JULY	16.00	16.00	16.00	16.00	0.00	0.00
8 AUGUST	15.00	15.00	15.00	15.00	0.00	0.00
9 SEPTEMBER	11.50	11.50	11.50	11.50	0.00	0.00
10 OCTOBER	8.40	8.40	8.40	8.40	0.00	0.00
11 NOVEMBER	4.40	4.40	4.40	4.40	0.00	0.00
12 DECEMBER	0.00	0.00	0.00	0.00	0.00	0.00

BIVALVE COST SIMULATION MODEL: ALGAE PRICE COMPUTATIONS (cont.)

DATE: MARCH 30, 1988

OUTPUT GENERATED

DATE OF PRICE COMPUTATIONS: ALGAE THESIS RESULTS

BATCH BEGUN: APRIL	1STAGE	2STAGE	3STAGE	4STAGE	5STAGE	6STAGE
ALGAE AVAILABLE FOR USE IN MONTH	APRIL	APRIL	APRIL	MAY	MAY	MAY
*TOTAL LITRES PRODUCED	0.2500E-01	4.170	70.00	3406.	0.1722E+05	0.6718E+05
LITRES OF ALGAE AVAILABLE	0.0000E+00	3.170	0.0000E+00	1511.	9652.	0.6718E+05
CELLS OF ALGAE PRODUCED	0.7500E+07	0.1251E+10	0.2100E+11	0.1362E+13	0.1119E+14	0.4031E+14
*TOTAL COST OF STAGE	50.64	80.90	49.79	110.3	109.3	107.1
TOTAL COST PER LITRE OF ALGAE	2025.	19.40	0.7113	0.3240E-01	0.6347E-02	0.1594E-02
TOTAL COST PER CELL OF ALGAE	0.672E-05	0.6467E-07	0.2371E-08	0.8099E-10	0.9764E-11	0.2656E-11
TOTAL COST OF STAGE, LESS INNOCULATION	50.64	30.26	30.39	60.55	47.91	59.01
*TOTAL COST OF ELECTRICITY	9.060	13.47	7.716	24.41	18.79	13.47
TOTAL COST OF ELECTRICITY/LITRE OF ALGAE	362.4	3.231	0.1102	0.7166E-02	0.1091E-02	0.2005E-03
TOTAL COST OF ELEC. LESS INNOCULATION	9.060	4.414	4.484	16.69	5.208	5.210
*TOTAL KWH USED	90.60	134.7	77.16	244.1	187.9	134.7
TOTAL KWH USED PER LITRE OF ALGAE	3624.	32.31	1.102	0.7166E-01	0.1091E-01	0.2005E-02
TOTAL KWH USED LESS INNOCULATION	90.60	44.14	44.84	166.9	52.08	52.10
*TOTAL COST OF FUEL OIL	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00
TOTAL COST OF OIL/LITRE OF ALGAE	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00
TOTAL COST OF OIL LESS INNOCULATION	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00
*TOTAL LITRES OF FUEL OIL	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00
TOTAL LITRES OIL USED/LITRE OF ALGAE	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00
TOTAL L. OIL USED LESS INNOCULATION	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00
*TOTAL COST OF ALL LABOR TYPES	41.43	67.19	41.87	84.46	87.18	88.12
TOTAL COST ALL LABOR/LITRE OF ALGAE	1657.	16.11	0.5982	0.2480E-01	0.5062E-02	0.1312E-02
TOTAL COST ALL LABOR LESS INNOCULATN	41.43	25.76	25.76	42.59	40.19	49.80
*TOTAL HOURS OF ALL LABOR TYPES	6.333	10.25	6.375	12.79	13.53	14.37
TOTAL HRS ALL LABOR USED/LITRE ALGAE	253.3	2.458	0.9107E-01	0.3756E-02	0.7858E-03	0.2138E-03
TTL HRS ALL LABOR LESS INNOCULATN	6.333	3.917	3.917	6.417	6.417	8.417
*TOTAL COST OF ALL MATERIALS	0.1494	0.2380	0.2017	1.470	3.333	5.468
TTL COST ALL MATERIALS/LITRE OF ALGAE	5.975	0.5708E-01	0.2882E-02	0.4315E-03	0.1935E-03	0.8139E-04
TTL COST ALL MATER. LESS INNOCULATION	0.1494	0.8867E-01	0.1446	1.268	2.515	4.003

BIVALE COST SIMULATION MODEL: INPUT ENTERED (CONT.) MARCH 30, 1988  
GENERAL DATA

1. MONTH THAT PRODUCTION SEASON BEGINS: JANUARY (-1)  
2. MONTH THAT PRODUCTION SEASON ENDS: DECEMBER (12)  
2 STAGE 3 STAGE 4 STAGE 5 STAGE

3. NUMBER OF DAYS IN EACH STAGE:

2.00 12.00 5.00 371.00

4. NUMBER OF PERIODS IN EACH STAGE:

1 3 1 53

5. FOR OPTION 1, THE NUMBER OF DAYS UP TO STAGE 2: 30.00

6. FOR OPTION 1, THE COST UP TO STAGE 2: 0.00

INFORMATION PERTAINING TO PRICE

INPUT: ELECTRICITY

7. PRICE COMPUTED BY PROGRAM?: N

INPUT: FUEL OIL

8. PRICE COMPUTED BY PROGRAM?: N

INPUT: LABOR

9. PRICE COMPUTED BY PROGRAM?: N

INPUT: MATERIALS

10. PRICE COMPUTED BY PROGRAM?: N

INPUT: ALGAE

11. PRICE COMPUTED BY PROGRAM?: Y

INPUT: CULCH

12. PRICE COMPUTED BY PROGRAM?: N

INPUT: ALGAE PER CELL (COMPUTED BY THE PROGRAM):

	JANUARY	FEBRUARY	MARCH	APRIL	MAY	JUNE	DECEMBER
PRIMARY1 :	0.6464135E-05	0.6464135E-05	0.6464135E-05	0.6751600E-05	0.7033068E-05	0.7215468E-05	0.6464135E-05
	0.7509438E-05	0.7444134E-05	0.7215468E-05	0.7012933E-05	0.6751600E-05	0.6464135E-05	
PRIMARY2 :	0.6208467E-07	0.6208467E-07	0.6208467E-07	0.6466979E-07	0.6725492E-07	0.6884125E-07	
	0.7148515E-07	0.7089761E-07	0.6884125E-07	0.6701993E-07	0.6466979E-07	0.6208467E-07	
PRIMARY3 :	0.2282785E-08	0.2282785E-08	0.2282785E-08	0.2371048E-08	0.2459312E-08	0.2513474E-08	
	0.2603744E-08	0.2583683E-08	0.2513474E-08	0.2451288E-08	0.2371048E-08	0.2282785E-08	
SECONDARY :	0.748333E-10	0.7488353E-10	0.7488353E-10	0.7725728E-10	0.8099153E-10	0.8380865E-10	
	0.8707120E-10	0.8792311E-10	0.872570E-10	0.8297109E-10	0.7938458E-10	0.7624403E-10	
TERTIARY :	0.9350308E-11	0.9350308E-11	0.9350308E-11	0.9511042E-11	0.9763898E-11	0.9954654E-11	
	0.1017557E-10	0.1023322E-10	0.1008446E-10	0.9897940E-11	0.9686914E-11	0.9442431E-11	
SOLAR :	0.2605360E-11	0.2605360E-11	0.2605360E-11	0.2624981E-11	0.2655846E-11	0.2679430E-11	
	0.2706097E-11	0.2713139E-11	0.2694976E-11	0.2672208E-11	0.2646449E-11	0.2616660E-11	

BIVALVE COST SIMULATION MODEL: INPUT ENTERED (CONT.)  
 BIVALVE DATA  
 1. TYPE OF BIVALVE 1 = OYSTERS  
 2. NAME OF SYSTEM UPWELL  
 3. NUMBER DAYS IN STAGE CONDITION 30.000  
 4. NUMBER BIVALVES IN STAGE CONDITION 2.0000  
 5. NUMBER OF OYSTERS BATCH BEGUN WITH: 1000000.

EQUATION TO DETERMINE SURVIVAL:

$$\text{POPULATION} = \text{POPULATION}(\text{INITIAL}) * [(\text{WEEKS}+1)^{\star\star A}]$$

6. PARAMETER A = -0.2550680

EQUATION TO EQUATE AGE TO SIZE:

$$\text{SIZE} = (\text{A} * \text{AGE}) + (\text{B} * \text{AGE SQUARED})$$

7. PARAMETER A = 0.7370000

8. PARAMETER B = -0.252000E-02

BIVALVE COST SIMULATION MODEL: INPUT ENTERED (CONT.)  
 ALGAE DATA

1. AGE/ALGAE EQUATION CHOSEN: 1  
 NATURAL LOG(SIZE IN MM) = A + B \* NAT LOG(10,000 CELLS)
2. PARAMETER A = -0.6510480
3. PARAMETER B = 0.3384260

IN THE FOLLOWING, WHERE QUANTITY=0, QUANTITY IS ENTERED

4. FILTERING RATE

5. FRACTION OF EACH ALGAE TYPE REQUIRED PER STAGE:

TYPE PRIMARY1	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
TYPE PRIMARY2	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
TYPE PRIMARY3	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
TYPE SECONDARY	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
TYPE TERTIARY	1.0	1.0	1.0	0.00E+00	0.00E+00
TYPE SOLAR	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.0

6. FRACTION FROM SEA VARIES BY MONTH: N

7. FRACTION OF DIET FROM SEA ALGAE IN EACH STAGE, EACH MONTH:

JANUARY	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.12
FEBRUARY	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.12
MARCH	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.12
APRIL	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.12
MAY	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.12
JUNE	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.12
JULY	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.12
AUGUST	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.12
SEPTEMBER	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.12
OCTOBER	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.12
NOVEMBER	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.12
DECEMBER	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.12

BIVALVE COST SIMULATION MODEL: OUTPUT GENERATED  
 DATE: MARCH 30, 1988  
 FILE: 1201 8DATA

BIVALVE TYPE: OYSTERS  
 NUMBER BIVALVE BEGUN: .10000E+07  
 SYSTEM USED: UPWELL  
 OPTION CHOSEN: 1, FIRST STAGE OF INTEREST IS THE SPAWN STAGE  
 (\*NOTE: STAGE 1 INCLUDES ALL STAGES UP TO THAT STAGE)

PRICE OF INPUTS PER MONTH

MONTH	JANUARY	FEBRUARY	MARCH	APRIL	OCTOBER	NOVEMBER	MAY	JUNE	DECEMBER
	JULY	AUGUST	SEPTEMBER						
ELECTRICITY/KWHR	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000
FUEL OIL /LITRE	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000
LABOR /HOUR	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000
MATERIALS /UNIT	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000
PRIMARY1 /CELL	0.6464135E-05	0.6464135E-05	0.6464135E-05	0.6464135E-05	0.6751600E-05	0.7039068E-05	0.7215468E-05	0.7215468E-05	0.6464135E-05
PRIMARY2 /CELL	0.7509468E-05	0.744134E-05	0.7215468E-05	0.701233E-05	0.6751600E-05	0.6751600E-05	0.6751600E-05	0.6751600E-05	0.6464135E-05
PRIMARY3 /CELL	0.6208467E-07	0.6208467E-07	0.6208467E-07	0.6208467E-07	0.6466979E-07	0.6725492E-07	0.6725492E-07	0.6725492E-07	0.6834125E-07
SECONDARY /CELL	0.7148515E-07	0.7089761E-07	0.6884125E-07	0.6884125E-07	0.6701993E-07	0.6466979E-07	0.6466979E-07	0.6466979E-07	0.6208467E-07
TERTIARY /CELL	0.2282785E-08	0.2282785E-08	0.2282785E-08	0.2371048E-08	0.2459312E-08	0.2459312E-08	0.2459312E-08	0.2459312E-08	0.2513474E-08
SOLAR /CELL	0.2603744E-08	0.2583683E-08	0.2513474E-08	0.2451288E-08	0.2371048E-08	0.2282785E-08	0.2282785E-08	0.2282785E-08	0.2282785E-08
CULCH /LITRE	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000

COST SIMULATION MODEL: OUTPUT GENERATED (CONT.)				MARCH 30, 1988				BATCH BEGUN: APRIL			
NAME OF BIVALVE		TIME UP TO MNTN END OF PERIOD		AGE AT END OF PERIOD		BIVALV NO. IN PERIOD		AVG SIZE IN PERIOD		SIZE BIVALV AT END PERIOD	
PRD *STG	STAGE										
1	1	CONDITION	4	30.00000	730.00000	2.0000000	2.0000000	49.45227	49.45227		
2	2	SPAWN	5	32.00000	2.0000000	967734.0	937909.1	0.1052085	0.2103657		
3	3	LARVAE	5	36.00000	6.0000000	892905.5	853937.6	0.3939469	0.6298628		
4	3	LARVAE	5	40.00000	10.000000	872521.6	797460.8	0.7601836	1.47713		
5	3	LARVAE	5	44.00000	14.000000	7756712.3	755616.8	1.2499980	1.463920		
6	4	SETTING	5	49.00000	19.000000	734727.4	715555.1	1.676703	1.981861		
7	5	JUVENILE	5	56.00000	26.000000	693416.0	673338.0	2.272559	2.702662		
8	5	JUVENILE	6	63.00000	33.000000	656581.6	641096.2	3.01759	3.18421		
9	5	JUVENILE	6	70.00000	40.000000	6227751.0	615260.1	3.744302	4.129142		
10	5	JUVENILE	6	77.00000	47.000000	604251.8	593853.4	4.473164	4.834821		
11	5	JUVENILE	6	84.00000	54.000000	584336.4	575671.4	5.1255461	5.535461		
12	5	JUVENILE	7	91.00000	61.000000	567532.6	559942.1	5.845417	6.231061		
13	5	JUVENILE	7	98.00000	68.000000	552892.6	54611.6	6.560244	6.921621		
14	5	JUVENILE	7	105.0000	75.000000	539864.1	533832.3	7.198956	7.607141		
15	5	JUVENILE	7	112.0000	82.000000	52819.8	52273.8	7.904419	8.287622		
16	5	JUVENILE	7	119.0000	89.000000	517716.3	512794.6	8.605203	8.963061		
17	5	JUVENILE	8	126.0000	96.000000	508146.7	503671.1	9.229816	9.633462		
18	5	JUVENILE	8	133.0000	103.00000	499626.1	495294.5	9.921291	10.29882		
19	5	JUVENILE	8	140.0000	110.00000	491378.7	487561.6	10.60814	10.95914		
20	5	JUVENILE	8	147.0000	117.00000	483931.8	480388.5	11.21867	11.61442		
21	5	JUVENILE	9	154.0000	124.00000	477009.6	473706.4	11.89805	12.26466		
22	5	JUVENILE	9	161.0000	131.00000	470548.7	467438.2	12.56816	12.90986		
23	5	JUVENILE	9	168.0000	138.00000	464496.7	461595.6	13.16335	13.55002		
24	5	JUVENILE	9	175.0000	145.00000	458809.6	456077.8	13.82870	14.18514		
25	5	JUVENILE	10	182.0000	152.00000	453449.7	450810.2	14.48736	14.81521		
26	5	JUVENILE	10	189.0000	159.00000	448384.1	445942.6	15.06990	15.44025		
27	5	JUVENILE	10	196.0000	166.00000	443385.7	441659.1	15.71921	16.06024		
28	5	JUVENILE	10	203.0000	173.00000	439029.6	436827.1	16.36383	16.67520		
29	5	JUVENILE	10	210.0000	180.00000	434695.1	432396.8	16.93234	17.28513		
30	5	JUVENILE	11	217.0000	187.00000	430563.1	428560.8	17.56760	17.89000		
31	5	JUVENILE	11	224.0000	194.00000	426651.7	424703.5	18.12675	18.48984		
32	5	JUVENILE	11	231.0000	201.00000	422844.1	421011.3	18.75266	19.08464		
33	5	JUVENILE	11	238.0000	208.00000	419229.0	417471.7	19.37389	19.67441		
34	5	JUVENILE	12	245.0000	215.00000	415761.1	41473.9	19.91889	20.25912		
35	5	JUVENILE	12	252.0000	222.00000	412430.1	410808.1	20.53085	20.83881		
36	5	JUVENILE	12	259.0000	229.00000	409226.6	40765.1	21.06662	21.41344		
37	5	JUVENILE	12	266.0000	236.00000	404614.7	404337.0	21.66911	21.98305		
38	5	JUVENILE	1	273.0000	243.00000	403167.8	40116.5	22.19551	22.54761		
39	5	JUVENILE	1	280.0000	250.00000	400298.1	398896.8	22.78865	23.10712		
40	5	JUVENILE	1	287.0000	257.00000	397526.6	39672.0	23.37712	23.66161		
41	5	JUVENILE	1	294.0000	264.00000	394846.6	393336.4	23.88947	24.21104		
42	5	JUVENILE	2	301.0000	271.00000	392253.4	39084.8	24.46858	24.75545		
43	5	JUVENILE	2	308.0000	278.00000	389742.0	38812.6	24.97157	25.29480		
44	5	JUVENILE	2	315.0000	285.00000	387307.6	386115.4	25.54131	26.82912		
45	5	JUVENILE	2	322.0000	292.00000	384946.1	383789.4	26.03494	26.35840		
46	5	JUVENILE	2	329.0000	299.00000	382654.3	381530.7	26.59534	26.88266		
47	5	JUVENILE	3	336.0000	306.00000	380427.6	379335.9	27.017961	27.40186		
48	5	JUVENILE	3	343.0000	313.00000	378263.6	377701.9	27.63063	27.91602		
49	5	JUVENILE	3	350.0000	320.00000	376158.6	375125.7	28.10555	28.42513		
50	5	JUVENILE	3	357.0000	327.00000	374110.3	373104.5	28.64720	28.92921		
51	5	JUVENILE	4	364.0000	334.00000	37115.6	370112.5	29.11276	29.42825		
52	5	JUVENILE	4	371.0000	341.00000	36917.2	36917.0	29.64507	29.92226		
53	5	JUVENILE	4	378.0000	348.00000	368277.6	367346.4	30.10126	30.41121		
54	5	JUVENILE	4	385.0000	355.00000	366430.0	365521.3	30.62421	30.89913		
55	5	JUVENILE	5	392.0000	362.00000	364626.7	363740.1	31.07103	31.37401		
56	5	JUVENILE	5	399.0000	369.00000	362866.6	362000.8	31.38463	31.8485		
57	5	JUVENILE	5	406.0000	376.00000	361147.4	360301.5	32.02209	32.31665		
58	5	JUVENILE	5	413.0000	383.00000	359467.6	358640.9	32.56329	32.95444		
59	5	JUVENILE	5	420.0000	390.00000	357825.6	357017.2	32.95444	33.23914		

PERIOD	STAGE	NAME OF	MNTH	COST OF	COST UP TO END AVG COST BIV			COST BIVALV		
					PERIOD	OF PERIOD	IN PERIOD/DAY	IN PERIOD	AT END PERIOD	
1	1	CONDITION	4	0.000000E+00	0.000000E+00	0.000000E+00	0.000000E+00	0.000000E+00	0.000000E+00	0.000000E+00
1	2	SPAWN	5	0.7474519E-02	0.7474519E-02	0.7474519E-02	0.386864E-08	0.7963341E-08	0.386864E-08	0.7963341E-08
3	3	LARVAE	5	0.4317153	0.4394898	0.4394898	0.12085137E-06	0.5143113E-06	0.12085137E-06	0.5143113E-06
4	4	LARVAE	5	2.615227	3.054416	3.054416	0.793467E-06	0.3830176E-06	0.793467E-06	0.3830176E-06
5	5	LARVAE	5	7.857286	10.9170	10.9170	0.253412E-05	0.1444079E-04	0.253412E-05	0.1444079E-04
6	6	SETTING	5	23.55299	34.46469	34.46469	0.6411356E-05	0.4816496E-04	0.6411356E-05	0.4816496E-04
7	7	JUVENILE	5	18.45863	52.92332	52.92332	0.3802835E-05	0.7859844E-04	0.3802835E-05	0.7859844E-04
8	8	JUVENILE	6	38.56452	91.46985	91.46985	0.838842E-05	0.1426773E-03	0.838842E-05	0.1426773E-03
9	9	JUVENILE	6	68.17392	159.6438	159.6438	0.155432E-04	0.2594735E-03	0.155432E-04	0.2594735E-03
10	10	JUVENILE	6	108.8479	268.4915	268.4915	0.4521173E-04	0.4521173E-03	0.4521173E-04	0.4521173E-03
11	11	JUVENILE	6	161.7548	430.2461	430.2461	0.395188E-04	0.7473773E-03	0.395188E-04	0.7473773E-03
12	12	JUVENILE	7	230.2422	660.4883	660.4883	0.579545E-04	0.1179565E-02	0.579545E-04	0.1179565E-02
13	13	JUVENILE	7	311.4697	971.9580	971.9580	0.804794E-04	0.1779747E-02	0.804794E-04	0.1779747E-02
14	14	JUVENILE	7	407.9121	1079.870	1079.870	0.1079404E-03	0.2584838E-02	0.1079404E-03	0.2584838E-02
15	15	JUVENILE	7	520.3247	1900.195	1900.195	0.140219E-03	0.3634693E-02	0.140219E-03	0.3634693E-02
16	16	JUVENILE	7	649.3794	2549.574	2549.574	0.179879E-03	0.4971918E-02	0.179879E-03	0.4971918E-02
17	17	JUVENILE	8	797.7424	3347.317	3347.317	0.224634E-03	0.6645836E-02	0.224634E-03	0.6645836E-02
18	18	JUVENILE	8	962.2297	4309.543	4309.543	0.2753387E-03	0.8700971E-02	0.2753387E-03	0.8700971E-02
19	19	JUVENILE	8	1144.994	5454.535	5454.535	0.3328808E-03	0.1118737E-01	0.3328808E-03	0.1118737E-01
20	20	JUVENILE	8	1346.435	6800.969	6800.969	0.397687E-03	0.1415722E-01	0.397687E-03	0.1415722E-01
21	21	JUVENILE	9	1556.416	8357.383	8357.383	0.4666228E-03	0.1764254E-01	0.4666228E-03	0.1764254E-01
22	22	JUVENILE	9	1794.602	10151.98	10151.98	0.5443356E-03	0.2171741E-01	0.5443356E-03	0.2171741E-01
23	23	JUVENILE	9	2052.225	12204.21	12204.21	0.6311669E-03	0.2643918E-01	0.6311669E-03	0.2643918E-01
24	24	JUVENILE	9	2329.478	14533.68	14533.68	0.1308051E-03	0.3186667E-01	0.1308051E-03	0.3186667E-01
25	25	JUVENILE	10	2604.325	17138.01	17138.01	0.8204796E-03	0.3801095E-01	0.8204796E-03	0.3801095E-01
26	26	JUVENILE	10	2918.578	20036.59	20036.59	0.9298711E-03	0.4497571E-01	0.9298711E-03	0.4497571E-01
27	27	JUVENILE	10	3252.629	23309.21	23309.21	0.1047512E-02	0.5282313E-01	0.1047512E-02	0.5282313E-01
28	28	JUVENILE	10	3616.510	26915.72	26915.72	0.117532E-02	0.6161642E-01	0.117532E-02	0.6161642E-01
29	29	JUVENILE	10	3980.225	30895.95	30895.95	0.1308051E-02	0.7141972E-01	0.1308051E-02	0.7141972E-01
30	30	JUVENILE	11	4331.574	35227.52	35227.52	0.1431779E-02	0.8219957E-01	0.1431779E-02	0.8219957E-01
31	31	JUVENILE	11	4740.824	39948.34	39948.34	0.1587512E-02	0.9410876E-01	0.1587512E-02	0.9410876E-01
32	32	JUVENILE	11	5169.527	45157.87	45157.87	0.1746516E-02	0.1072130	0.1746516E-02	0.1072130
33	33	JUVENILE	11	5617.547	50155.42	50155.42	0.1911244E-02	0.1215781	0.1911244E-02	0.1215781
34	34	JUVENILE	12	6016.117	56771.54	56771.54	0.2067161E-02	0.1371048	0.2067161E-02	0.1371048
35	35	JUVENILE	12	6496.813	63268.35	63268.35	0.2253359E-02	0.1540095	0.2253359E-02	0.1540095
36	36	JUVENILE	12	6996.082	70264.38	70264.38	0.2442266E-02	0.1725580	0.2442266E-02	0.1725580
37	37	JUVENILE	12	7513.711	77778.06	77778.06	0.2642889E-02	0.19222169	0.2642889E-02	0.19222169
38	38	JUVENILE	12	8014.852	85792.88	85792.88	0.283955E-02	0.2135657	0.283955E-02	0.2135657
39	39	JUVENILE	1	8566.055	94338.88	94338.88	0.3057026E-02	0.2363496	0.3057026E-02	0.2363496
40	40	JUVENILE	1	9134.746	103493.6	103493.6	0.328708E-02	0.2612339	0.328708E-02	0.2612339
41	41	JUVENILE	1	9720.613	113214.1	113214.1	0.3516958E-02	0.2878840	0.3516958E-02	0.2878840
42	42	JUVENILE	2	10323.35	123537.4	123537.4	0.3755722E-02	0.3156648	0.3755722E-02	0.3156648
43	43	JUVENILE	2	10942.60	134480.0	134480.0	0.4010931E-02	0.3461406	0.4010931E-02	0.3461406
44	44	JUVENILE	2	11578.04	146058.0	146058.0	0.4270520E-02	0.3782755	0.4270520E-02	0.3782755
45	45	JUVENILE	2	12299.26	158387.3	158387.3	0.453391E-02	0.4124326	0.453391E-02	0.4124326
46	46	JUVENILE	2	12895.91	171183.1	171183.1	0.4814457E-02	0.4486746	0.4814457E-02	0.4486746
47	47	JUVENILE	3	13577.59	184760.7	184760.7	0.509619E-02	0.4870635	0.509619E-02	0.4870635
48	48	JUVENILE	3	14273.87	199034.5	199034.5	0.5390748E-02	0.5276604	0.5390748E-02	0.5276604
49	49	JUVENILE	3	14984.34	214018.8	214018.8	0.5690739E-02	0.5705256	0.5690739E-02	0.5705256
50	50	JUVENILE	3	15708.59	229727.4	229727.4	0.5998455E-02	0.6157185	0.5998455E-02	0.6157185
51	51	JUVENILE	4	16570.00	246297.4	246297.4	0.6361309E-02	0.6636314	0.6361309E-02	0.6636314
52	52	JUVENILE	4	17326.07	263623.4	263623.4	0.6688494E-02	0.7140064	0.6688494E-02	0.7140064
53	53	JUVENILE	4	18094.64	281718.1	281718.1	0.7019021E-02	0.7669003	0.7019021E-02	0.7669003
54	54	JUVENILE	4	18875.22	300593.3	300593.3	0.7358730E-02	0.82235686	0.7358730E-02	0.82235686
55	55	JUVENILE	5	19898.57	320491.8	320491.8	0.7766064E-02	0.8811012	0.7766064E-02	0.8811012
56	56	JUVENILE	5	20711.18	341202.9	341202.9	0.8153796E-02	0.9423476	0.8153796E-02	0.9423476
57	57	JUVENILE	5	21534.45	362737.4	362737.4	0.8518264E-02	1.006761	0.8518264E-02	1.006761
58	58	JUVENILE	5	22367.88	385105.3	385105.3	0.8889288E-02	1.073791	0.8889288E-02	1.073791
59	59	JUVENILE	5	23210.93	408316.1	408316.1	0.9266656E-02	1.143687	0.9266656E-02	1.143687

## COST OF INPUTS IN A PERIOD:

PRD	*STAGE	MNTH	AIGAE	COST OF INPUTS UP TO END OF A PERIOD	AIGAE	QUANTITY OF INPUT USED PER PERIOD:
2	2 SPAWN	5	0.7474519E-02	0.7474519E-02	TERTIARY	SOLAR
3	3 LARVAE	5	0.4391898	0.4391898	0.765265E+09	0.0000000E+00
4	3 LARVAE	5	2.6515227	2.6515227	0.4421547E+11	0.0000000E+00
5	3 LARVAE	5	7.837286	7.837286	0.2678467E+12	0.0000000E+00
6	4 SETTING	5	23.55299	23.55299	0.8047285E+12	0.0000000E+00
7	5 JUVENILE	5	18.55863	18.55863	0.2412253E+13	0.0000000E+00
8	5 JUVENILE	6	38.54652	38.54652	0.6950191E+13	0.0000000E+00
9	5 JUVENILE	6	91.46985	91.46985	0.0000000E+00	0.1438777E+14
10	5 JUVENILE	6	159.6438	159.6438	0.0000000E+00	0.2544629E+14
11	5 JUVENILE	6	108.8479	108.8479	0.0000000E+00	0.4062808E+14
12	5 JUVENILE	7	161.7548	161.7548	0.6037587E+14	0.0000000E+00
13	5 JUVENILE	7	230.2422	230.2422	0.0000000E+00	0.8508278E+14
14	5 JUVENILE	7	311.4697	311.4697	0.0000000E+00	0.1150993E+15
15	5 JUVENILE	7	407.9121	407.9121	0.0000000E+00	0.1507383E+15
16	5 JUVENILE	7	520.3247	520.3247	0.0000000E+00	0.1922787E+15
17	5 JUVENILE	8	649.3794	649.3794	0.0000000E+00	0.2398691E+15
18	5 JUVENILE	8	797.7424	797.7424	0.0000000E+00	0.2904294E+15
19	5 JUVENILE	8	962.2297	962.2297	0.0000000E+00	0.3545556E+15
20	5 JUVENILE	8	1144.994	1144.994	0.0000000E+00	0.4220181E+15
21	5 JUVENILE	9	1346.435	1346.435	0.0000000E+00	0.4962647E+15
22	5 JUVENILE	9	1556.416	1556.416	0.0000000E+00	0.5752495E+15
23	5 JUVENILE	9	1794.602	1794.602	0.0000000E+00	0.6639065E+15
24	5 JUVENILE	9	2052.225	2052.225	0.0000000E+00	0.7615004E+15
25	5 JUVENILE	10	2329.478	2329.478	0.0000000E+00	0.8633780E+15
26	5 JUVENILE	10	2604.325	2604.325	0.0000000E+00	0.9755968E+15
27	5 JUVENILE	10	2918.578	2918.578	0.0000000E+00	0.1022197E+16
28	5 JUVENILE	10	3252.629	3252.629	0.0000000E+00	0.1217207E+16
29	5 JUVENILE	10	3606.510	3606.510	0.0000000E+00	0.1396377E+16
30	5 JUVENILE	11	3980.225	3980.225	0.0000000E+00	0.1489489E+16
31	5 JUVENILE	11	4331.574	4331.574	0.0000000E+00	0.1636750E+16
32	5 JUVENILE	11	4740.824	4740.824	0.0000000E+00	0.1791391E+16
33	5 JUVENILE	11	5169.527	5169.527	0.0000000E+00	0.1933383E+16
34	5 JUVENILE	12	5617.547	5617.547	0.0000000E+00	0.2122674E+16
35	5 JUVENILE	12	6016.117	6016.117	0.0000000E+00	0.2289207E+16
36	5 JUVENILE	12	6496.813	6496.813	0.0000000E+00	0.24422916E+16
37	5 JUVENILE	12	6996.082	6996.082	0.0000000E+00	0.2637326E+16
38	5 JUVENILE	12	7513.711	7513.711	0.0000000E+00	0.281549E+16
39	5 JUVENILE	1	8014.852	8014.852	0.0000000E+00	0.3076293E+16
40	5 JUVENILE	1	8566.055	8566.055	0.0000000E+00	0.3287859E+16
41	5 JUVENILE	1	9134.746	9134.746	0.0000000E+00	0.356136E+16
42	5 JUVENILE	1	9720.613	9720.613	0.0000000E+00	0.371006E+16
43	5 JUVENILE	2	10323.35	10323.35	0.0000000E+00	0.392351E+16
44	5 JUVENILE	2	10942.60	10942.60	0.0000000E+00	0.4200033E+16
45	5 JUVENILE	2	11578.04	11578.04	0.0000000E+00	0.4433929E+16
46	5 JUVENILE	2	12229.26	12229.26	0.0000000E+00	0.4693884E+16
47	5 JUVENILE	3	12895.91	12895.91	0.0000000E+00	0.4949761E+16
48	5 JUVENILE	3	13577.59	13577.59	0.0000000E+00	0.5211406E+16
49	5 JUVENILE	4	14273.87	14273.87	0.0000000E+00	0.5478656E+16
50	5 JUVENILE	3	14984.34	14984.34	0.0000000E+00	0.571352E+16
51	5 JUVENILE	4	15708.59	15708.59	0.0000000E+00	0.6029335E+16
52	5 JUVENILE	4	16570.00	16570.00	0.0000000E+00	0.6312429E+16
53	5 JUVENILE	4	17326.07	17326.07	0.0000000E+00	0.6600459E+16
54	5 JUVENILE	4	18094.64	18094.64	0.0000000E+00	0.6833246E+16
55	5 JUVENILE	5	18875.22	18875.22	0.0000000E+00	0.7190613E+16
56	5 JUVENILE	5	19898.57	19898.57	0.0000000E+00	0.7492369E+16
57	5 JUVENILE	5	20711.18	20711.18	0.0000000E+00	0.7798338E+16
58	5 JUVENILE	5	21534.45	21534.45	0.0000000E+00	0.8108318E+16
59	5 JUVENILE	5	22367.38	22367.38	0.0000000E+00	0.8422130E+16
		23210.93	23210.93	0.0000000E+00	0.8739563E+16	